



Acid deposition to semi-natural ecosystems - 2000



Percentage of the area of semi-natural ecosystems receiving acid deposition above the critical loads for the emissions of the year 2000. Calculation results for the meteorological conditions of 1997, using ecosystem-specific deposition.

Source: IIASA


Acid deposition to semi-natural ecosystems – Baseline 2020



Percentage of the area of semi-natural ecosystems receiving acid deposition above the critical loads for the emissions of the current legislation case of the “Climate policy” scenario in 2020. Calculation results for the meteorological conditions of 1997, using ecosystem-specific deposition.

Source: IIASA


Acid deposition to semi-natural ecosystems – Scenario A



Percentage of the area of semi-natural ecosystems receiving acid deposition above the critical loads for the emissions of the D23 (A) scenario in 2020. Calculation results for the meteorological conditions of 1997, using ecosystem-specific deposition.

Source: IIASA


Acid deposition to semi-natural ecosystems – Scenario B



Percentage of the area of semi-natural ecosystems receiving acid deposition above the critical loads for the emissions of the D23 (B) scenario in 2020. Calculation results for the meteorological conditions of 1997, using ecosystem-specific deposition.

Source: IIASA


Acid deposition to semi-natural ecosystems – Scenario C



Percentage of the area of semi-natural ecosystems receiving acid deposition above the critical loads for the emissions of the D23 (C) scenario in 2020. Calculation results for the meteorological conditions of 1997, using ecosystem-specific deposition.

Source: IIASA

Acid deposition to semi-natural ecosystems – Thematic Strategy 2020



Percentage of the area of semi-natural ecosystems receiving acid deposition above the critical loads for the emissions of the Thematic Strategy on Air Pollution in 2020. Calculation results for the meteorological conditions of 1997, using ecosystem-specific deposition.

Source: IIASA