

Notifications of the co-ordinated programme (specific exercise) to the European Commission for RICE (husked and polished)											
	Number of samples analysed	Without detectable residues	%	With residues at or below MRL or without MRL	%	With residues above MRL	%	With residues above EC-MRL	%	With residues above national MRL	%
Country											
Belgium	41	41	100	0	0.0	0	0.0	0	0.0	0	0.0
Denmark	12	11	92	1	8.3	0	0.0	0	0.0	0	0.0
Germany	287	256	89	24	8.4	7	2.4	7	2.4	0	0.0
Greece	22	21	95	0	0.0	1	4.5	1	4.5	0	0.0
Spain	43	42	98	1	2.3	0	0.0	0	0.0	0	0.0
France	38	22	58	16	42.1	0	0.0	0	0.0	0	0.0
Ireland	12	12	100	0	0.0	0	0.0	0	0.0	0	0.0
Italy	76	68	89	8	10.5	0	0.0	0	0.0	0	0.0
Luxembourg	15	14	93	1	6.7	0	0.0	0	0.0	0	0.0
The Netherlands	43	43	100	0	0.0	0	0.0	0	0.0	0	0.0
Austria	12	12	100	0	0.0	0	0.0	0	0.0	0	0.0
Portugal	70	58	83	7	10.0	5	7.1	5	7.1	0	0.0
Finland	22	18	82	4	18.2	0	0.0	0	0.0	0	0.0
Sweden	41	39	95	2	4.9	0	0.0	0	0.0	0	0.0
United Kingdom	96	84	88	11	11.5	1	1.0	1	1.0	0	0.0
Norway	27	26	96	1	3.7	0	0.0	0	0.0	0	0.0
Iceland	0	0	0	0	0.0	0	0.0	0	0.0	0	0.0
Liechtenstein	12	11	92	0	0.0	1	8.3	1	8.3	0	0.0
TOTAL	869	778	90	76	8.7	15	1.7	15	1.7	0	0.0

Notifications of the co-ordinated programme (specific exercise) to the European Commission for CUCUMBER											
	Number of samples analysed	Without detectable residues	%	With residues at or below MRL or without MRL	%	With residues above MRL	%	With residues above EC-MRL	%	With residues above national MRL	%
Country											
Belgium	44	36	82	3	6.8	5	11.4	5	11.4	0	0.0
Denmark	47	44	94	2	4.3	1	2.1	1	2.1	0	0.0
Germany	173	126	73	44	25.4	3	1.7	1	0.6	2	1.2
Greece	85	53	62	27	31.8	5	5.9	5	5.9	0	0.0
Spain	44	38	86	6	13.6	0	0.0	0	0.0	0	0.0
France	96	86	90	10	10.4	0	0.0	0	0.0	0	0.0
Ireland	12	11	92	1	8.3	0	0.0	0	0.0	0	0.0
Italy	76	68	89	8	10.5	0	0.0	0	0.0	0	0.0
Luxembourg	15	14	93	0	0.0	1	6.7	1	6.7	0	0.0
The Netherlands	88	79	90	9	10.2	0	0.0	0	0.0	0	0.0
Austria	13	9	69	4	30.8	0	0.0	0	0.0	0	0.0
Portugal	61	47	77	12	19.7	2	3.3	2	3.3	0	0.0
Finland	115	92	80	23	20.0	0	0.0	0	0.0	0	0.0
Sweden	141	129	91	12	8.5	0	0.0	0	0.0	0	0.0
United Kingdom	59	43	73	16	27.1	0	0.0	0	0.0	0	0.0
Norway	84	76	90	8	9.5	0	0.0	0	0.0	0	0.0
Iceland	11	11	100	0	0.0	0	0.0	0	0.0	0	0.0
Liechtenstein	12	10	83	2	16.7	0	0.0	0	0.0	0	0.0
TOTAL	1176	972	83	187	15.9	17	1.4	15	1.3	2	0.2

Notifications of the co-ordinated programme (specific exercise) to the European Commission for HEAD CABBAGE											
	Number of samples analysed	Without detectable residues	%	With residues at or below MRL or without MRL	%	With residues above MRL	%	With residues above EC-MRL	%	With residues above national MRL	%
Country											
Belgium	44	27	61	11	25.0	6	13.6	4	9.1	2	4.5
Denmark	41	41	100	0	0.0	0	0.0	0	0.0	0	0.0
Germany	300	100	33	168	56.0	32	10.7	32	10.7	0	0.0
Greece	19	14	74	1	5.3	4	21.1	4	21.1	0	0.0
Spain	45	31	69	13	28.9	1	2.2	1	2.2	0	0.0
France	64	52	81	8	12.5	4	6.3	4	6.3	0	0.0
Ireland	13	10	77	3	23.1	0	0.0	0	0.0	0	0.0
Italy	102	97	95	5	4.9	0	0.0	0	0.0	0	0.0
Luxembourg	15	15	100	0	0.0	0	0.0	0	0.0	0	0.0
The Netherlands	31	26	84	5	16.1	0	0.0	0	0.0	0	0.0
Austria	12	10	83	2	16.7	0	0.0	0	0.0	0	0.0
Portugal	69	64	93	5	7.2	0	0.0	0	0.0	0	0.0
Finland	20	19	95	1	5.0	0	0.0	0	0.0	0	0.0
Sweden	53	53	100	0	0.0	0	0.0	0	0.0	0	0.0
United Kingdom	72	61	85	11	15.3	0	0.0	0	0.0	0	0.0
Norway	44	44	100	0	0.0	0	0.0	0	0.0	0	0.0
Iceland	0	0	#DIV/0!	0	#DIV/0!	0	#DIV/0!	0	#DIV/0!	0	#DIV/0!
Liechtenstein	12	12	100	0	0.0	0	0.0	0	0.0	0	0.0
TOTAL	956	676	71	233	24.4	47	4.9	45	4.7	2	0.2

Notifications of the co-ordinated programme (specific exercise) to the European Commission for PEAS (frozen or fresh, without pods)

	Number of samples analysed	Without detectable residues	%	With residues at or below MRL or without MRL	%	With residues above MRL	%	With residues above EC-MRL	%	With residues above national MRL	%
Country											
Belgium	34	24	71	7	20.6	3	8.8	3	8.8	0	0.0
Denmark	7	7	100	0	0.0	0	0.0	0	0.0	0	0.0
Germany	258	146	57	98	38.0	14	5.4	14	5.4	0	0.0
Greece	47	38	81	6	12.8	3	6.4	3	6.4	0	0.0
Spain	42	41	98	1	2.4	0	0.0	0	0.0	0	0.0
France	42	32	76	9	21.4	1	2.4	1	2.4	0	0.0
Ireland	12	12	100	0	0.0	0	0.0	0	0.0	0	0.0
Italy	60	57	95	3	5.0	0	0.0	0	0.0	0	0.0
Luxembourg	15	5	33	9	60.0	1	6.7	1	6.7	0	0.0
The Netherlands	16	13	81	3	18.8	0	0.0	0	0.0	0	0.0
Austria	12	11	92	1	8.3	0	0.0	0	0.0	0	0.0
Portugal	66	65	98	1	1.5	0	0.0	0	0.0	0	0.0
Finland	21	21	100	0	0.0	0	0.0	0	0.0	0	0.0
Sweden	14	14	100	0	0.0	0	0.0	0	0.0	0	0.0
United Kingdom	62	59	95	3	4.8	0	0.0	0	0.0	0	0.0
Norway	10	9	90	1	10.0	0	0.0	0	0.0	0	0.0
Iceland	0	0	0	0	0.0	0	0.0	0	0.0	0	0.0
Liechtenstein	12	12	100	0	0.0	0	0.0	0	0.0	0	0.0
TOTAL	730	566	78	142	19.5	22	3.0	22	3.0	0	0.0

Notifications of the co-ordinated programme (specific exercise) to the European Commission for ALL FOUR PRODUCTS											
	Number of samples analysed	Without detectable residues	%	With residues at or below MRL or without MRL	%	With residues above MRL	%	With residues above EC-MRL	%	With residues above national MRL	%
Country											
Belgium	163	128	79	21	12.9	14	8.6	12	7.4	2	1.2
Denmark	107	103	96	3	2.8	1	0.9	1	0.9	0	0.0
Germany	1018	628	62	334	32.8	56	5.5	54	5.3	2	0.2
Greece	173	126	73	34	19.7	13	7.5	13	7.5	0	0.0
Spain	174	152	87	21	12.1	1	0.6	1	0.6	0	0.0
France	240	192	80	43	17.9	5	2.1	5	2.1	0	0.0
Ireland	49	45	92	4	8.2	0	0.0	0	0.0	0	0.0
Italy	314	290	92	24	7.6	0	0.0	0	0.0	0	0.0
Luxembourg	60	48	80	10	16.7	2	3.3	2	3.3	0	0.0
The Netherlands	178	161	90	17	9.6	0	0.0	0	0.0	0	0.0
Austria	49	42	86	7	14.3	0	0.0	0	0.0	0	0.0
Portugal	266	234	88	25	9.4	7	2.6	7	2.6	0	0.0
Finland	178	150	84	28	15.7	0	0.0	0	0.0	0	0.0
Sweden	249	235	94	14	5.6	0	0.0	0	0.0	0	0.0
United Kingdom	289	247	85	41	14.2	1	0.3	1	0.3	0	0.0
Norway	165	155	94	10	6.1	0	0.0	0	0.0	0	0.0
Iceland	11	11	100	0	0.0	0	0.0	0	0.0	0	0.0
Liechtenstein	48	45	94	2	4.2	1	2.1	1	2.1	0	0.0
TOTAL	3731	2992	80	638	17.1	101	2.7	97	2.6	4	0.1

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Head cabbage	
Reporting country: <u>Belgium</u>		Year of sampling: <u>2000</u>	Remark:
Total number of samples analysed:	44	With residues above MRL (EC+national):	6
Without detectable residues:	27	With residues above EC-MRL:	4
With detectable residues at or below MRL or without MRL:	11	With residues above national MRL:	2

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	44	44	0,02-0,2																		
Benomyl group(#)	38	37	0,1-0,2					1										0.2	0	3	E
Chlorpyrifos	44	44	0.05																		
Chlorpyrifos-methyl	44	44	0.05																		
Deltamethrin	44	44	0,05-0,1																		
Diazinon	44	44	0,02-0,05																		
Endosulfan	44	44	0.05																		
Imazalil	44	43	0.1							1								0.7	1	0.02	E
Iprodione	44	44	0,1-0,2																		
Lambda-cyhalothrin	44	44	0.05																		
Maneb-group(##)	38	25	1							2	9	2						2.9	4	2-1	N-E
Mecarbam	44	44	0.05																		
Metalaxyl	44	40	0,05-0,1				3		1									0.3	0	1	E
Methamidophos	44	44	0.01																		
Methidathion	44	44	0,02-0,05																		
Permethrin	44	44	0,05-0,1																		
Pirimiphos-methyl	44	44	0.05																		
Thiabendazol	38	38	0,05-0,1																		
Triazophos	0																				
Vinclozolin	44	43	0,02-0,05				1											0.06	1	0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg
 (**) in alphabetical order of the English name
 (***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).
 (##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Legume vegetables (fresh)		Food item: Peas (frozen or fresh, analysed without pods)	
Reporting country:	<u>Belgium</u>	Year of sampling:	<u>2000</u>
Remark:		Only add information regarding peas and the 20 pesticides and do not change or delete rows or columns	
Total number of samples analysed:	34	With residues above MRL (EC+national):	3
Without detectable residues:	24	With residues above EC-MRL:	3
With detectable residues at or below MRL or without MRL:	7	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	34	34	0,02-0,2																	
Benomyl group(#)	28	21	0,1-0,2				5	2									0.2	2	0.1	E
Chlorpyrifos	34	34	0.05																	
Chlorpyrifos-methyl	34	34	0.05																	
Deltamethrin	34	34	0,05-0,1																	
Diazinon	34	34	0,02-0,05																	
Endosulfan	34	34	0.05																	
Imazalil	34	34	0.1																	
Iprodione	34	34	0,1-0,2																	
Lambda-cyhalothrin	34	34	0.05																	
Maneb-group(##)	34	33	0,5-0,6									1					2.5	1	0.1	E
Mecarbam	34	34	0.05																	
Metalaxyl	34	34	0,05-0,1																	
Methamidophos	34	34	0.01																	
Methidathion	34	34	0,02-0,05																	
Permethrin	34	34	0,05-0,1																	
Pirimiphos-methyl	34	34	0.05																	
Thiabendazol	28	28	0,05-0,1																	
Triazophos	0																			
Vinclozolin	34	32	0,02-0,05					1	1								0.22	0	0.3	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg
(**) in alphabetical order of the English name
(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).
(##) Sum of dithiocarbamates, expressed as CS₂

Table B:		Notifications of the co-ordinated programme (specific exercise) to the European Commission																					
Product group: Cereals		Food item: Rice (husked or polished)																					
Reporting country: Denmark		Year of sampling: 2000										Remark:											
Total number of samples analysed:		12		With residues above MRL (EC+national):										0								Only add information regarding rice and the 20 pesticides and do not change or delete rows or columns	
Without detectable residues:		11		With residues above EC-MRL:										0									
With detectable residues at or below MRL or without MRL:		1		With residues above national MRL:										0									
		Samples with quantifiable residues in classes up to and including (in mg/kg) (*)																					
Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50	Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)			
Acephate	0																						
Benomyl group(#)	0																						
Chlorpyrifos	12	12	0.005	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E			
Chlorpyrifos-methyl	12	12	0.005	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3.00	E			
Deltamethrin	12	11	0.01	0	1	0	0	0	0	0	0	0	0	0	0	0	0.018	0	1.00	E			
Diazinon	12	12	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05 (a)	E			
Endosulfan	0																						
Imazalil	0																						
Iprodione	0																						
Lambda-cyhalothrin	12	12	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.02	E			
Maneb-group(##)	0																						
Mecarbam	0																						
Metalaxyl	0																						
Methamidophos	0																						
Methidathion	0																						
Permethrin	0																						
Pirimiphos-methyl	12	12	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5.00	E			
Thiabendazol	0																						
Triazophos	0																						
Vinclozolin	12	12	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E			
(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg				(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).																			
(**) in alphabetical order of the English name				(##) Sum of dithiocarbamates, expressed as CS ₂																			
(***) E=EC-MRL, N=National MRL, W=without MRL				(a) 0.02 mg/kg from 1. juli 2000																			

Table B:		Notifications of the co-ordinated programme (specific exercise) to the European Commission																		
Product group: Fruiting vegetables				Food item: Cucumber																
Reporting country: Denmark		Year of sampling: 2000				Remark:														
Total number of samples analysed:		47		With residues above MRL (EC+national):												1		Only add information regarding cucumber and the 20 pesticides and do not change or delete rows or columns		
Without detectable residues:		44		With residues above EC-MRL:												1				
With detectable residues at or below MRL or without MRL:		2		With residues above national MRL:												0				
Samples with quantifiable residues in classes up to and including (in mg/kg) (*)																				
Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50	Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
Acephate	47	47	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.02	E
Benomyl group(#)	47	46	0.05	0	0	0	1	0	0	0	0	0	0	0	0	0	0.085	0	0.50	E
Chlorpyrifos	47	47	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E
Chlorpyrifos-methyl	47	47	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E
Deltamethrin	47	47	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.10	E
Diazinon	47	47	0.009	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.5 (a)	E
Endosulfan	47	47	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1 (c)	E
Imazalil	47	47	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.20	E
Iprodione	47	47	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2.00	E
Lambda-cyhalothrin	47	47	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.10	E
Maneb-group(##)	47	46	0.10	0	0	0	0	0	0	1	0	0	0	0	0	0	0.65	1	0.50	E
Mecarbam	47	47	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E
Metalaxyl	47	47	0.03	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	(c)	W / E
Methamidophos	47	45	0.006	0	1	1	0	0	0	0	0	0	0	0	0	0	0.039	0	1.00	E
Methidathion	47	47	0.03	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.02	E
Permethrin	47	47	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.10	E
Pirimiphos-methyl	47	47	0.007	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	(c)	W / E
Thiabendazol	47	47	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	(c)	W / E
Triazophos	47	47	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	(a)	W / E
Vinclozolin	47	47	0.03	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1.00	E
(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg																			#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).	
(**) in alphabetical order of the English name																			##) Sum of dithiocarbamates, expressed as CS ₂	
(***) E=EC-MRL, N=National MRL, W=without MRL																			(a) 0.02 mg/kg from 1. juli 2000	
																			(c) 0.05 mg/kg from 1. juli 2000	

Table B:		Notifications of the co-ordinated programme (specific exercise) to the European Commission																			
Product group: Legume vegetables (fresh)				Food item: Peas (frozen or fresh, analysed without pods)																	
Reporting country:		Denmark		Year of sampling:				2000				Remark:									
Total number of samples analysed:		7		With residues above MRL (EC+national):				0				Only add information regarding peas and the 20 pesticides and do not change or delete rows or columns									
Without detectable residues:		7		With residues above EC-MRL:				0													
With detectable residues at or below MRL or without MRL:		0		With residues above national MRL:				0													
Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	7	7	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.02	E
Benomyl group(#)	7	7	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.10	E
Chlorpyrifos	7	7	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E
Chlorpyrifos-methyl	7	7	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E
Deltamethrin	7	7	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E
Diazinon	7	7	0.009	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.02	E
Endosulfan	7	7	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1 (c)	E
Imazail	0																				
Iprodione	7	7	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.20	E
Lambda-cyhalothrin	7	7	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.02	E
Maneb-group(##)	0																				
Mecarbam	7	7	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E
Metalaxyl	7	7	0.03	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E
Methamidophos	7	7	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.01	E
Methidathion	7	7	0.03	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.02	E
Permethrin	7	7	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E
Pirimiphos-methyl	7	7	0.007	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E
Thiabendazol	7	7	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E
Triazophos	7	7	0.006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	(a)	W / E
Vinclozolin	7	7	0.03	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.30	E
(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg				(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).																	
(**) in alphabetical order of the English name				(##) Sum of dithiocarbamates, expressed as CS ₂																	
(*** E=EC-MRL, N=National MRL, W=without MRL				(a) 0.02 mg/kg from 1. juli 2000																	
				(c) 0.05 mg/kg from 1. juli 2000																	

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Rice (husked or polished)	
Reporting country: <u>Germany</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed:	287	With residues above MRL (EC+national):	7
Without detectable residues:	256	With residues above EC-MRL:	7
With detectable residues at or below MRL or without MRL:	24	With residues above national MRL:	0

Only add information regarding rice and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	249	248	0.02	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0.06	1	0.02	E
Benomyl group(#)	222	222	0.1	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0.1	E
Chlorpyrifos	261	260	0.02	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0.01		0.05	E
Chlorpyrifos-methyl	273	273	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0			3	E
Deltamethrin	261	261	0.1	0	0	0	0	0	0	0	0	0	0	0	0	0	0			1	E
Diazinon	265	265	0.04	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0.05	E
Endosulfan	273	273	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0.1	E
Imazalil	247	247	0.1	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0.02	E
Iprodione	261	260	0.04	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0.045		3	E
Lambda-cyhalothrin	236	236	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0.02	E
Maneb-group(##)	239	228	0.05	0	2	3	6	0	0	0	0	0	0	0	0	0	0	0.08	6	0.05	E
Mecarbam	273	273	0.04	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0.05	E
Metalaxyl	255	255	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0.05	E
Methamidophos	245	245	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0.01	E
Methidathion	273	273	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0.02	E
Permethrin	273	273	0.1	0	0	0	0	0	0	0	0	0	0	0	0	0	0			2	E
Pirimiphos-methyl	267	249	0.02	8	5	4	1	0	0	0	0	0	0	0	0	0	0	0.059		5	E
Thiabendazol	234	234	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0.2	N
Triazophos	277	277	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0.02	E
Vinclozolin	273	272	0.02	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0.1			

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Cucumber	
Reporting country: <u>Germany</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed:	173	With residues above MRL (EC+national):	3
Without detectable residues:	126	With residues above EC-MRL:	1
With detectable residues at or below MRL or without MRL:	44	With residues above national MRL:	2

Only add information regarding cucumber and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	137	137	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.02	E	
Benomyl group(#)	129	118	0.05	1	0	3	4	2	0	1	0	0	0	0	0	0	0.78	1	0.5	E
Chlorpyrifos	173	172	0.01	0	0	1	0	0	0	0	0	0	0	0	0	0	0.03		0.05	E
Chlorpyrifos-methyl	173	173	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0			0.05	E
Deltamethrin	161	161	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0			0.1	E
Diazinon	146	146	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0			0.5	E
Endosulfan	173	162	0.005	2	2	3	4	0	0	0	0	0	0	0	0	0.098		1	E	
Imazalil	131	128	0.1	3	0	0	0	0	0	0	0	0	0	0	0	0.002		0.2	E	
Iprodione	162	158	0.02	1	1	0	0	2	0	0	0	0	0	0	0	0.19		2	E	
Lambda-cyhalothrin	149	149	0.02	0	0	0	0	0	0	0	0	0	0	0	0			0.1	E	
Maneb-group(##)	114	95	0.02	1	3	3	4	7	1	0	0	0	0	0	0	0.22		0.5	E	
Mecarbam	173	173	0.02	0	0	0	0	0	0	0	0	0	0	0	0			0.05	E	
Metalaxyl	157	150	0.05	1	0	4	1	0	1	0	0	0	0	0	0	0.23	2	0.05	N	
Methamidophos	135	134	0.05	0	0	0	0	1	0	0	0	0	0	0	0	0.19		1	E	
Methidathion	173	173	0.02	0	0	0	0	0	0	0	0	0	0	0	0			0.02	E	
Permethrin	173	173	0.05	0	0	0	0	0	0	0	0	0	0	0	0			0.1	E	
Pirimiphos-methyl	174	174	0.01	0	0	0	0	0	0	0	0	0	0	0	0			0.5	N	
Thiabendazol	135	133	0.05	1	0	1	0	0	0	0	0	0	0	0	0	0.04		0.05	E	
Triazophos	171	171	0.02	0	0	0	0	0	0	0	0	0	0	0	0			0.02	N	
Vinclozolin	172	172	0.01	0	0	0	0	0	0	0	0	0	0	0	0			1	E	

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Head cabbage	
Reporting country: <u>Germany</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed:	300	With residues above MRL (EC+national):	32
Without detectable residues:	100	With residues above EC-MRL:	32
With detectable residues at or below MRL or without MRL:	168	With residues above national MRL:	0

Only add information regarding head cabbage and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	253	253	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	E
Benomyl group(#)	224	220	0.1	0	0	1	1	1	0	1	0	0	0	0	0	0	0	0.52	3	E
Chlorpyrifos	277	275	0.01	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0.085	1	E
Chlorpyrifos-methyl	278	278	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E
Deltamethrin	263	262	0.05	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0.01	0.1	E
Diazinon	278	278	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.5	E
Endosulfan	279	276	0.005	0	1	1	0	0	1	0	0	0	0	0	0	0	0	0.36	1	E
Imazalil	238	238	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.02	E
Iprodione	263	257	0.02	1	0	2	1	2	0	0	0	0	0	0	0	0	0	0.14	5	E
Lambda-cyhalothrin	248	248	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.2	E
Maneb-group(##)	266	73	0.043	0	2	20	28	49	42	33	15	4	0	0	0	0	2.7	19	1	E
Mecarbam	278	278	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E
Metalaxyl	238	238	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	E
Methamidophos	278	277	0.05	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0.13	0.5	E
Methidathion	278	278	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.02	E
Permethrin	278	276	0.05	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0.091	1	E
Pirimiphos-methyl	278	278	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	N
Thiabendazol	227	218	0.05	0	0	0	0	1	1	0	1	5	1	0	0	0	6	9	0.05	E
Triazophos	253	253	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.2	N
Vinclozolin	278	270	0.01	1	1	1	0	0	1	3	0	1	0	0	0	0	1.5	5	0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(**) in alphabetical order of the English name

(##) Sum of dithiocarbamates, expressed as CS₂

(***) E=EC-MRL, N=National MRL, W=without MRL

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Legume vegetables (fresh)		Food item: Peas (frozen or fresh, analysed without pods)	
Reporting country:	<u>Germany</u>	Year of sampling:	<u>2000</u>
Remark:			
Only add information regarding peas and the 20 pesticides and do not change or delete rows or columns			
Total number of samples analysed:	258	With residues above MRL (EC+national):	14
Without detectable residues:	146	With residues above EC-MRL:	14
With detectable residues at or below MRL or without MRL:	98	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	251	251	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.02	E	
Benomyl group(#)	203	195	0.1	1	2	1	0	2	2	0	0	0	0	0	0	0	0.32	4	0.1	E
Chlorpyrifos	247	247	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E		
Chlorpyrifos-methyl	255	255	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E		
Deltamethrin	254	254	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E		
Diazinon	255	255	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0.02	E		
Endosulfan	255	254	0.005	1	0	0	0	0	0	0	0	0	0	0	0	0	0.0009	0.05	E	
Imazalil	255	255	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0.02	E		
Iprodione	254	247	0.02	2	2	0	3	0	0	0	0	0	0	0	0	0	0.1	0.2	E	
Lambda-cyhalothrin	238	238	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0.2	E		
Maneb-group(##)	246	210	0.02	1	4	15	8	8	0	0	0	0	0	0	0	0	0.2	8	0.1	E
Mecarbam	255	255	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E		
Metalaxyl	255	255	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E		
Methamidophos	251	251	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0.01	E		
Methidathion	255	255	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0.02	E		
Permethrin	254	254	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E		
Pirimiphos-methyl	255	255	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E		
Thiabendazol	250	250	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05	E		
Triazophos	255	255	0.02	0	0	0	0	0	0	0	0	0	0	0	0	0	0.2	N		
Vinclozolin	255	172	0.01	7	6	22	21	20	7	0	0	0	0	0	0	0	0.41	2	0.3	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Cucumber	
Reporting country:	<u>Greece</u>	Year of sampling:	<u>2000</u>
Remark: Only add information regarding cucumbers and the 20 pesticides and do not change or delete rows or columns			
Total number of samples analysed:	85	With residues above MRL (EC+national):	5
Without detectable residues:	53	With residues above EC-MRL:	5
With detectable residues at or below MRL or without MRL:	27	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	23	21	0.02		1	1												0.05	1	0.02*	E
Benomyl group(#)	85	83						2										0.15		0.50	E
Chlorpyrifos	23	23	0.01																	0.05*	E
Chlorpyrifos-methyl	23	22	0.01		1													0.02		0.05*	E
Deltamethrin	23	22	0.02			1												0.04		0.10	E
Diazinon	23	23	0.01																	0.02*	E
Endosulfan	23	16	0.01	5		2												0.05		1.00	E
Imazalil	23	23	0.05																	0.20	E
Iprodione	23	23	0.02																	2.00	E
Lambda-cyhalothrin	23	23	0.01																	0.10	E
Maneb-group(##)	85	64					1	4	13	3								0.85	3	0.50	E
Mecarbam	23	23	0.01																	0.05*	E
Metalaxyl	23	22	0.02						1									0.22	1	0.05*	E
Methamidophos	23	19	0.02			2			2									0.39		1.00	E
Methidathion	23	23	0.01																	0.02*	E
Permethrin	23	22	0.02				1											0.07		0.10	E
Pirimiphos-methyl	23	23	0.01																	0.05*	E
Thiabendazol	43	43	0.05																	0.05*	E
Triazophos	23	23	0.02																	0.02*	E
Vinclozolin	23	23	0.01																	1.00	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Head cabbage	
Reporting country: <u>Greece</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed:	19	With residues above MRL (EC+national):	4
Without detectable residues:	14	With residues above EC-MRL:	4
With detectable residues at or below MRL or without MRL:	1	With residues above national MRL:	

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	19	19	0.02																2.00	E
Benomyl group(#)	19	19																	3.00	E
Chlorpyrifos	19	19	0.01																1.00	E
Chlorpyrifos-methyl	19	19	0.01																0.05*	E
Deltamethrin	19	19	0.02																0.10	E
Diazinon	19	19	0.01																0.02*	E
Endosulfan	19	19	0.01																0.05*	E
Imazalil	19	19	0.05																0.02*	E
Iprodione	19	19	0.02																5.00	E
Lambda-cyhalothrin	19	19	0.01																0.20	E
Maneb-group(##)	19	15								1	3						1.76	3	1.00	E
Mecarbam	19	19	0.01																0.05*	E
Metalaxyl	19	19	0.02																1.00	E
Methamidophos	19	19	0.02																0.50	E
Methidathion	19	18	0.01			1											0.025	1	0.02*	E
Permethrin	19	19	0.02																1.00	E
Pirimiphos-methyl	19	19	0.01																0.05*	E
Thiabendazol	19	19	0.05																0.05*	E
Triazophos	19	19	0.02																0.02*	E
Vinclozolin	19	19	0.01																0.05*	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Legume vegetables (fresh)		Food item: Peas (frozen or fresh, analysed without pods)	
Reporting country:	<u>Greece</u>	Year of sampling:	<u>2000</u>
Remark:		Only add information regarding peas and the 20 pesticides and do not change or delete rows or columns	
Total number of samples analysed:	<u>47</u>	With residues above MRL (EC+national):	<u>3</u>
Without detectable residues:	<u>38</u>	With residues above EC-MRL:	<u>3</u>
With detectable residues at or below MRL or without MRL:	<u>6</u>	With residues above national MRL:	<u> </u>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	25	25	0.02															0.02*		E	
Benomyl group(#)	47	44						3										0.18	3	0.10	E
Chlorpyrifos	25	25	0.01															0.05*		E	
Chlorpyrifos-methyl	25	25	0.01															0.05*		E	
Deltamethrin	25	25	0.02															0.05*		E	
Diazinon	25	25	0.01															0.02*		E	
Endosulfan	25	25	0.01															0.05*		E	
Imazalil	25	25	0.05															0.02*		E	
Iprodione	25	25	0.02															0.20		E	
Lambda-cyhalothrin	25	25	0.01															0.20		E	
Maneb-group(##)	37	37																0.10		E	
Mecarbam	25	25	0.01															0.05*		E	
Metalaxyl	25	21	0.02			4												0.04		E	
Methamidophos	25	25	0.02															0.01*		E	
Methidathion	25	25	0.01															0.02*		E	
Permethrin	25	25	0.02															0.05*		E	
Pirimiphos-methyl	25	25	0.01															0.05*		E	
Thiabendazol	35	35	0.05															0.05*		E	
Triazophos	25	25	0.02															0.02*		E	
Vinclozolin	25	23	0.01		2													0.02		0.30	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg
 (**) in alphabetical order of the English name
 (***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).
 (##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Rice (husked or polished)	
Reporting country: <u>Spain</u>	Year of sampling: <u>2000</u>	Remark: Only add information regarding rice and the 20 pesticides and do not change or delete rows or columns	
Total number of samples analysed:	<input type="text" value="43"/>	With residues above MRL (EC+national):	<input type="text" value="0"/>
Without detectable residues:	<input type="text" value="42"/>	With residues above EC-MRL:	<input type="text"/>
With detectable residues at or below MRL or without MRL:	<input type="text" value="1"/>	With residues above national MRL:	<input type="text"/>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	43	43	0.02																0.02	E
Benomyl group(#)	43	43	0.10																0.10	E
Chlorpyrifos	43	42	0.05		1													0.05	0.05	E
Chlorpyrifos-methyl	43	43	0.05																3.00	E
Deltamethrin	43	43	0.05																1.00	E
Diazinon	43	43	0.02																0.02	E
Endosulfan	43	43	0.05																0.05	E
Imazalil	43	43	0.02																0.02	E
Iprodione	43	43	0.02																3.00	E
Lambda-cyhalothrin	43	43	0.02																0.02	E
Maneb-group(##)	43	43	0.05																0.05	E
Mecarbam	43	43	0.05																0.05	E
Metalaxyl	43	43	0.05																0.05	E
Methamidophos	43	43	0.01																0.01	E
Methidathion	43	43	0.02																0.02	E
Permethrin	43	43	0.05																0.05	E
Pirimiphos-methyl	43	43	0.05																5.00	E
Thiabendazol	43	43	0.05																0.10	E
Triazophos	43	43	0.02																0.02	E
Vinclozolin	43	43	0.05																0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Cucumber	
Reporting country: <u>Spain</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed:	<input type="text" value="44"/>	With residues above MRL (EC+national):	<input type="text" value="0"/>
Without detectable residues:	<input type="text" value="38"/>	With residues above EC-MRL:	<input type="text"/>
With detectable residues at or below MRL or without MRL:	<input type="text" value="6"/>	With residues above national MRL:	<input type="text"/>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)		
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50						
Acephate	44	44	0.02																	0.02	E	
Benomyl group(#)	44	44	0.10																		0.50	E
Chlorpyrifos	44	44	0.05																		0.05	E
Chlorpyrifos-methyl	44	44	0.05																		0.05	E
Deltamethrin	44	44	0.05																		0.10	E
Diazinon	44	44	0.02																		0.50	E
Endosulfan	44	42	0.05				1	1									0.14				1.00	E
Imazalil	44	44	0.02																		0.20	E
Iprodione	44	44	0.02																		2.00	E
Lambda-cyhalothrin	44	44	0.02																		0.02	E
Maneb-group(##)	44	41	0.05				1			2								0.48			0.50	E
Mecarbam	44	44	0.05																		0.02	E
Metalaxyl	44	43	0.05				1										0.05				0.05	E
Methamidophos	44	44	0.01																		1.00	E
Methidathion	44	44	0.02																		0.02	E
Permethrin	44	44	0.05																		0.10	E
Pirimiphos-methyl	44	44	0.05																		0.20	E
Thiabendazol	44	44	0.05																		1.00	N
Triazophos	44	44	0.02																		0.02	E
Vinclozolin	44	44	0.05																		1.00	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Head cabbage	
Reporting country: <u>Spain</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed:	45	With residues above MRL (EC+national):	1
Without detectable residues:	31	With residues above EC-MRL:	1
With detectable residues at or below MRL or without MRL:	13	With residues above national MRL:	

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	45	44	0.02					1									0.13	1	0.02	E
Benomyl group(#)	45	45	0.10																0.50	E
Chlorpyrifos	45	45	0.05																1.00	E
Chlorpyrifos-methyl	45	45	0.05																0.05	E
Deltamethrin	45	44	0.05		1														0.10	E
Diazinon	45	45	0.02																0.50	E
Endosulfan	45	45	0.05																1.00	E
Imazalil	45	45	0.02																0.02	E
Iprodione	45	45	0.02																5.00	E
Lambda-cyhalothrin	45	44	0.02					1									0.14		0.20	E
Maneb-group(##)	45	35	0.05				2	3	4	1							0.8		1.00	E
Mecarbam	45	45	0.05																0.02	E
Metalaxyl	45	45	0.05																1.00	E
Methamidophos	45	44	0.01	1													0.01		0.01	E
Methidathion	45	45	0.02																0.02	E
Permethrin	45	45	0.05																1.00	E
Pirimiphos-methyl	45	44	0.05						1								0.3		1.00	E
Thiabendazol	45	45	0.05																0.05	E
Triazophos	45	45	0.02																0.02	E
Vinclazolin	45	45	0.05																0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Legume vegetables (fresh)		Food item: Peas (frozen or fresh, analysed without pods)	
Reporting country: <u>Spain</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed: <input type="text" value="42"/>	With residues above MRL (EC+national): <input type="text" value="0"/>	Only add information regarding peas and the 20 pesticides and do not change or delete rows or columns	
Without detectable residues: <input type="text" value="41"/>	With residues above EC-MRL: <input type="text"/>		
With detectable residues at or below MRL or without MRL: <input type="text" value="1"/>	With residues above national MRL: <input type="text"/>		

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	42	42	0.02																3.00	E
Benomyl group(#)	42	42	0.10																0.10	E
Chlorpyrifos	42	42	0.05																0.05	E
Chlorpyrifos-methyl	42	42	0.05																0.05	E
Deltamethrin	42	42	0.05																0.10	E
Diazinon	42	42	0.02																0.50	E
Endosulfan	42	42	0.05																1.00	E
Imazalil	42	42	0.02																0.02	E
Iprodione	42	42	0.02																0.20	E
Lambda-cyhalothrin	42	42	0.02																0.20	E
Maneb-group(##)	42	41	0.05							1							0.99		1.00	E
Mecarbam	42	42	0.05																0.02	E
Metalaxyl	42	42	0.05																0.05	E
Methamidophos	42	42	0.01																0.50	E
Methodathion	42	42	0.02																0.02	E
Permethrin	42	42	0.05																0.10	E
Pirimiphos-methyl	42	42	0.05																0.50	E
Thiabendazol	42	42	0.05																0.05	E
Triazophos	42	42	0.02																0.02	E
Vinclozolin	42	42	0.05																2.00	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg
(**) in alphabetical order of the English name
(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).
(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Cucumber	
Reporting country: <u>France</u>	Year of sampling: <u>2000</u>	Remark: Only add information regarding cucumbers and the 20 pesticides and do not change or delete rows or columns	
Total number of samples analysed:	96	With residues above MRL (EC+national):	0
Without detectable residues:	86	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	10	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	96	96	0.01															0.02	0	0.02	EC
Benomyl group(#)	96	95	0.02		1													0.02	0	1	EC
Chlorpyrifos	96	96	0.01																	0.05	EC
Chlorpyrifos-methyl	96	96	0.01																	0.05	EC
Deltamethrin	96	96	0.01																	0.10	EC
diazinon	96	95	0.01			1												0.02	0	0.02	EC
Endosulfan	96	96	0.01																	0.3/1	EC
Imazalil	12	12	0.05																	0.20	EC
Iprodione	96	93	0.01		1		1		1									0.47	0	2.00	EC
Lambda-cyhalothrin	96	96	0.01																	0.10	EC
Maneb-group(##)	96	94	0.05				1		1									0.34	0	0.5	EC
Mecarbam	96	96	0.01																	0.05	EC
Metalaxyl	96	95	0.02					1										0.11	0	0.5	EC
Methamidophos	96	96	0.01																	1.00	EC
Methidathion	96	96	0.01																	0.02	EC
Permethrin	96	95	0.02		1													0.02	0	0.10	EC
Pirimiphos-methyl	96	96	0.01																	0.05	EC
Thiabendazol	96	96	0, 02																	0.05	EC
Triazophos	96	96	0.01																	0.02	EC
Vinclazolin	96	95	0.01		1													0.01	0	1.00	EC

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(**) in alphabetical order of the English name

(##) Sum of dithiocarbamates, expressed as CS₂

(***) E=EC-MRL, N=National MRL, W=without MRL

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Head cabbage	
Reporting country: <u>France</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed:	64	With residues above MRL (EC+national):	4
Without detectable residues:	52	With residues above EC-MRL:	4
With detectable residues at or below MRL or without MRL:	8	With residues above national MRL:	0

Only add information regarding head cabbage and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	64	64	0.01														0.01	0	2.00	EC
Benomyl group(##)	64	64	0.02														0.02	0	3.00	EC
Chlorpyrifos	64	64	0.01														0.01	0	1.00	EC
Chlorpyrifos-methyl	64	64	0.01														0.01	0	0.05	EC
Deltamethrin	64	64	0.01														0.01	0	0.10	EC
Diazinon	64	64	0.01														0.01	0	0.02	EC
Endosulfan	64	64	0.01														0.01	0	1/0,05	EC
Imazalil	64	64	0.05															0	0.02	EC
Iprodione	64	63	0.01					1									0.18	0	5	EC
Lambda-cyhalothrin	64	64	0.01														0.01	0	0.20	EC
Maneb-group(##)	64	54	0.05					4	2	1	3						2.5	4	1.00	EC
Mecarbam	64	64	0.01														0.01	0	0.05	EC
Metalaxyl	64	64	0.01														0.01	0	1.00	EC
Methamidophos	64	64	0.01														0.01	0	0.50	EC
Methidathion	64	64	0.01														0.01	0	0.02	EC
Permethrin	64	64	0.05														0.05	0	1.00	EC
Pririmphos-methyl	64	64	0.01														0.01	0	1.00	EC
Thiabendazol	64	64	0.02														0.02	0	0.05	EC
Triazophos	64	64	0.01														0.01	0	0.02	EC
Vinclozolin	64	63	0.01	1													0.01	0	0.05	EC

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(##) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(**) in alphabetical order of the English name

(##) Sum of dithiocarbamates, expressed as CS₂

(***) E=EC-MRL, N=National MRL, W=without MRL

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Rice (husked or polished)	
Reporting country: Ireland		Year of sampling: 2000	Remark: Only add information regarding rice and the 20 pesticides and do not change or delete rows or columns
Total number of samples analysed:	12	With residues above MRL (EC+national):	0
Without detectable residues:	12	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	0	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	12	12	0.05																	
Benomyl group(#)	12	12	0.05																	
Chlorpyrifos	12	12	0.06																	
Chlorpyrifos-methyl	12	12	0.05																	
Deltamethrin	12	12	0.05																	
Diazinon	12	12	0.05																	
Endosulfan	12	12	0.02-0.03																	
Imazalil	0	0																		
Iprodione	12	12	0.10																	
Lambda-cyhalothrin	12	12	0.02																	
Maneb-group(##)	0	0																		
Mecarbam	0	0																		
Metalaxyl	0	0																		
Methamidophos	12	12	0.04																	
Methodathion	12	12	0.04																	
Permethrin	12	12	0.20																	
Pirimiphos-methyl	12	12	0.05																	
Thiabendazol	12	12	0.05																	
Triazophos	12	12	0.03																	
Vinclozolin	12	12	0.02																	

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS2

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables

Food item: Cucumber

Reporting country: <u>Ireland</u>	Year of sampling: <u>2000</u>	Remark:		
Total number of samples analysed: <table border="1"><tr><td>12</td></tr></table>	12	With residues above MRL (EC+national): <table border="1"><tr><td>0</td></tr></table>	0	Only add information regarding cucumbers and the 20 pesticides and do not change or delete rows or columns
12				
0				
Without detectable residues: <table border="1"><tr><td>11</td></tr></table>	11	With residues above EC-MRL: <table border="1"><tr><td>0</td></tr></table>	0	
11				
0				
With detectable residues at or below MRL or without MRL: <table border="1"><tr><td>1</td></tr></table>	1	With residues above national MRL: <table border="1"><tr><td>0</td></tr></table>	0	
1				
0				

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	12	12	0.05																		
Benomyl group(#)	12	11	0.05			1												0.05	0	0.50	E
Chlorpyrifos	12	12	0.06																		
Chlorpyrifos-methyl	12	12	0.05																		
Deltamethrin	12	12	0.05																		
Diazinon	12	12	0.05																		
Endosulfan	12	12	0.02-0.03																		
Imazalil																					
Iprodione	12	12	0.10																		
Lambda-cyhalothrin	12	12	0.02																		
Maneb-group(##)																					
Mecarbam																					
Metalaxyl																					
Methamidophos	12	11	0.04				1											0.1	0	1.00	E
Methidathion	12	12	0.04																		
Permethrin	12	12	0.20																		
Pirimiphos-methyl	12	12	0.05																		
Thiabendazol	12	12	0.05																		
Triazophos	12	12	0.03																		
Vinclozolin	12	12	0.02																		

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS2

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Rice (husked or polished)	
Reporting country: <u>Italy</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed:	<input type="text" value="76"/>	With residues above MRL (EC+national):	<input type="text"/>
Without detectable residues:	<input type="text" value="68"/>	With residues above EC-MRL:	<input type="text"/>
With detectable residues at or below MRL or without MRL:	<input type="text" value="8"/>	With residues above national MRL:	<input type="text"/>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Accephate	34	34																n.d.		0,02	E
Benomyl group(#)	8	8																n.d.		0,1	E
Chlorpyrifos	64	64																n.d.		0,05	E
Chlorpyrifos-methyl	44	44																n.d.		0,1	E
Deltamethrin	46	46																n.d.		1	E
Diazinon	70	70																n.d.		0,05	E
Endosulfan	56	56																n.d.		0,01	N
Imazalil	17	17																n.d.		0,02	E
Iprodione	40	39							1									0,35		3	E
Lambda-cyhalothrin																					
Maneb-group(##)	5	5																n.d.		0,05	E
Mecarbam	11	11																n.d.		0,05	E
Metalaxyl	44	44																n.d.		0,05	E
Methamidophos	36	36																n.d.		0,01	E
Methodathion	29	29																n.d.		0,02	E
Permethrin	41	41																n.d.		2	E
Pirimiphos-methyl	64	57		2	1	2			1	1								0,58		5	E
Thiabendazol	11	11																n.d.		0,2	N
Triazophos	7	7																n.d.		0,02	E
Vinclozolin	25	25																n.d.		0,05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Cucumber	
Reporting country: <u>Italy</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed:	<u>102</u>	With residues above MRL (EC+national):	<input type="text"/>
Without detectable residues:	<u>97</u>	With residues above EC-MRL:	<input type="text"/>
With detectable residues at or below MRL or without MRL:	<u>5</u>	With residues above national MRL:	<input type="text"/>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	64	64																n.d.	0,02	E
Benomyl group(#)	68	68																n.d.	0,5	E
Chlorpyrifos	88	87			1													0,03	0,05	E
Chlorpyrifos-methyl	84	83			1													0,04	0,05	E
Deltamethrin	72	72																n.d.	0,1	E
Diazinon	92	92																n.d.	0,5	E
Endosulfan	86	86																n.d.	1	E
Imazalil	79	79																n.d.	0,2	E
Iprodione	83	83																n.d.	2	E
Lambda-cyhalothrin																				
Maneb-group(##)	78	76		1			1											0,06	0,5	E
Mecarbam	58	58																n.d.	0,05	E
Metalaxyl	76	75				1												0,08	0,2	N
Methamidophos	85	85																n.d.	1	E
Methidathion	87	87																n.d.	0,02	E
Permethrin	72	72																n.d.	0,1	E
Pirimiphos-methyl	88	88																n.d.	0,5	N
Thiabendazol	77	77																n.d.	0,05	N
Triazophos	54	54																n.d.	0,02	N
Vinclozolin	89	89																n.d.	1	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Legume vegetables (fresh)		Food item: Head cabbage	
Reporting country: <u>Italy</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed:	<input type="text" value="60"/>	With residues above MRL (EC+national):	<input type="text"/>
Without detectable residues:	<input type="text" value="57"/>	With residues above EC-MRL:	<input type="text"/>
With detectable residues at or below MRL or without MRL:	<input type="text" value="3"/>	With residues above national MRL:	<input type="text"/>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	24	23							1								0,3	2	E	
Benomyl group(##)	19	19															n.d.	3	E	
Chlorpyrifos	51	51															n.d.	0,05	E	
Chlorpyrifos-methyl	46	46															n.d.	0,05	E	
Deltamethrin	32	32															n.d.	0,1	E	
Diazinon	49	49															n.d.	0,5	E	
Endosulfan	45	45															n.d.	1	E	
Imazalil	34	34															n.d.	0,02	E	
Iprodione	42	42															n.d.	0,02	E	
Lambda-cyhalothrin																				
Maneb-group(##)	29	27						1	1								0,83	1	E	
Mecarbam	13	13															n.d.	0,05	E	
Metalaxyl	36	36															n.d.	0,05	E	
Methamidophos	36	36															n.d.	0,5	E	
Methidathion	47	47															n.d.	0,02	E	
Permethrin	36	36															n.d.	0,05	E	
Pirimiphos-methyl	47	47															n.d.	0,5	N	
Thiabendazol	30	30															n.d.	0,05	E	
Triazophos	16	16															n.d.	0,02	N	
Vinclozolin	51	51															n.d.	0,05	E	

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg
 (**) in alphabetical order of the English name
 (***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).
 (##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Legume vegetables (fresh)		Food item: Peas (frozen or fresh, analysed without pods)	
Reporting country:	<u>Italy</u>	Year of sampling:	<u>2000</u>
Total number of samples analysed: <input type="text" value="41"/>		With residues above MRL (EC+national): <input type="text"/>	
Without detectable residues: <input type="text" value="41"/>		With residues above EC-MRL: <input type="text"/>	
With detectable residues at or below MRL or without MRL: <input type="text"/>		With residues above national MRL: <input type="text"/>	
Remark: Only add information regarding peas and the 20 pesticides and do not change or delete rows or columns			

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	15	15															n.d.		0,02	E
Benomyl group(##)	16	16															n.d.		0,1	E
Chlorpyrifos	31	31															n.d.		0,05	E
Chlorpyrifos-methyl	27	27															n.d.		0,05	E
Deltamethrin	14	14															n.d.		0,05	E
Diazinon	35	35															n.d.		0,5	E
Endosulfan	23	23															n.d.		1	E
Imazalil	16	16															n.d.		0,02	E
Iprodione	26	26															n.d.		0,02	E
Lambda-cyhalothrin																				
Maneb-group(##)	21	21															n.d.		0,1	E
Mecarbam	12	12															n.d.		0,05	E
Metalaxyl	15	15															n.d.		0,05	E
Methamidophos	22	22															n.d.		0,01	E
Methidathion	26	26															n.d.		0,02	E
Permethrin	10	10															n.d.		0,05	E
Pirimiphos-methyl	34	34															n.d.		0,05	E
Thiabendazol	16	16															n.d.		0,05	E
Triazophos	9	9															n.d.		0,02	N
Vinclozolin	31	31															n.d.		0,3	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg
 (**) in alphabetical order of the English name
 (***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).
 (##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Head cabbage	
Reporting country:	<u>The Netherlands</u>	Year of sampling:	<u>2000</u>
Total number of samples analysed:		With residues above MRL (EC+national):	
Without detectable residues:		With residues above EC-MRL:	
With detectable residues at or below MRL		With residues above national MRL:	
or without MRL:			

Remark: Only add information regarding head cabbage and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	29	29	0.02																		
Benomyl group(#)	29	28	0.05						1									0.32		3.00	EG
Chlorpyrifos	29	29	0.03																		
Chlorpyrifos-methyl	29	29	0.03																		
Deltamethrin	29	29	0.03																		
Diazinon	29	29	0.02																		
Endosulfan	29	29	0.01																		
Imazalil	29	29	0.02																		
Iprodione	29	28	0.02					1										0.12		5.00	EG
Lambda-cyhalothrin	29	29	0.02																		
Maneb-group(##)	3	0	0.01						3									0.62		2.00	EG
Mecarbam	29	29	0.03																		
Metalaxyl	29	29	0.03																		
Methamidophos	29	29	0.01																		
Methidathion	29	29	0.02																		
Permethrin	29	28	0.03			1												0.05		1.00	EG
Pirimiphos-methyl	29	29	0.03																		
Thiabendazol	29	29	0.05																		
Triazophos	29	29	0.02																		
Vinclozolin	29	29	0.03																		

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg
 (**) in alphabetical order of the English name
 (***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).
 (##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Legume vegetables (fresh)		Food item: Peas (frozen or fresh, analysed without pods)	
Reporting country:	<u>The Netherlands</u>	Year of sampling:	<u>2000</u>
Remark:		Only add information regarding peas and the 20 pesticides and do not change or delete rows or columns	
Total number of samples analysed:	<input type="text" value="16"/>	With residues above MRL (EC+national):	<input type="text" value="0"/>
Without detectable residues:	<input type="text" value="13"/>	With residues above EC-MRL:	<input type="text" value="0"/>
With detectable residues at or below MRL or without MRL:	<input type="text" value="3"/>	With residues above national MRL:	<input type="text" value="0"/>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	16	16	0.02																		
Benomyl group(#)	15	13	0.05				1	1										0.11		0.10	E
Chlorpyrifos	16	16	0.03																		
Chlorpyrifos-methyl	16	16	0.03																		
Deltamethrin	16	16	0.03																		
Diazinon	16	16	0.02																		
Endosulfan	16	16	0.01																		
Imazalil	16	16	0.02																		
Iprodione	16	16	0.02																		
Lambda-cyhalothrin	16	16	0.02																		
Maneb-group(##)	1	0	0.01					1										0.2		2.00	E
Mecarbam	16	16	0.03																		
Metalaxyl	16	16	0.03																		
Methamidophos	16	16	0.01																		
Methidathion	16	16	0.02																		
Permethrin	16	16	0.03																		
Pirimiphos-methyl	16	16	0.03																		
Thiabendazol	15	15	0.05																		
Triazophos	16	16	0.02																		
Vinclozolin	16	16	0.03																		

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg
 (**) in alphabetical order of the English name
 (***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).
 (##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Rice (husked or polished)	
Reporting country:	<u>Austria</u>	Year of sampling:	<u>2000</u>
Total number of samples analysed:		With residues above MRL (EC+national):	
Without detectable residues:		With residues above EC-MRL:	
With detectable residues at or below MRL or without MRL:		With residues above national MRL:	
	12		0
	12		0
	0		0
Remark: Only add information regarding rice and the 20 pesticides and do not change or delete rows or columns			

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	12	12	0,02;0,05																	0.02	E
Benomyl group(#)	12	12	0.10																	0.10	E
Chlorpyrifos	12	12	0,01;0,02																	0.05	E
Chlorpyrifos-methyl	12	12	0,01;0,02																	3.00	E
Deltamethrin	12	12	0.05																	1.00	E
Diazinon	12	12	0,01;0,02																	0.02	E
Endosulfan	12	12	0.01																	0.05	E
Imazalil	12	12	0,03;0,1																	0.02	E
Iprodione	12	12	0,02;0,04																	3.00	E
Lambda-cyhalothrin	12	12	0.05																	0.02	E
Maneb-group(##)	12	12	0,02;0,2																	0.05	E
Mecarbam	12	12	0,01;0,02																	0.05	E
Metalaxyl	12	12	0,03;0,05																	0.05	E
Methamidophos	12	12	0,01;0,03																	0.01	E
Methidathion	12	12	0,01;0,02																	0.02	E
Permethrin	12	12	0.05																	0.05	E
Pirimiphos-methyl	12	12	0,01;0,02																	5.00	E
Thiabendazol	12	12	0,05;0,1																	0.10	E
Triazophos	12	12	0,01;0,02																	0.02	E
Vinclozolin	12	12	0.01																	0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Cucumber	
Reporting country:	<u>Austria</u>	Year of sampling:	<u>2000</u>
Remark:		Only add information regarding cucumbers and the 20 pesticides and do not change or delete rows or columns	
Total number of samples analysed:	13	With residues above MRL (EC+national):	0
Without detectable residues:	9	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	4	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	13	13	0,02;0,05															0.02	13	E
Benomyl group(#)	13	12	0.10				1											0.14	12	E
Chlorpyrifos	13	13	0,01;0,02															0.05	13	E
Chlorpyrifos-methyl	13	13	0,01;0,02															0.05	13	E
Deltamethrin	13	13	0.05															0.10	13	E
Diazinon	13	13	0,01;0,02															0.50	13	E
Endosulfan	13	11	0.01			2												0.31	11	E
Imazalil	13	13	0,03;0,1															0.20	13	E
Iprodione	13	13	0,02;0,04															2.00	13	E
Lambda-cyhalothrin	13	13	0.05															0.02	13	E
Maneb-group(##)	13	13	0,02;0,2															0.50	13	E
Mecarbam	13	13	0,01;0,02															0.02	13	E
Metalaxyl	13	13	0,03;0,05															0.05	13	E
Methamidophos	13	12	0,01;0,03		1													0.012	12	E
Methodathion	13	13	0,01;0,02															0.02	13	E
Permethrin	13	13	0.05															0.10	13	E
Pirimiphos-methyl	13	13	0,01;0,02															0.20	13	E
Thiabendazol	13	13	0,05;0,1															1.50	13	N
Triazophos	13	13	0,01;0,02															0.02	13	E
Vinclozolin	13	13	0.01															1.00	13	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Head cabbage	
Reporting country: <u>Austria</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed:	12	With residues above MRL (EC+national):	0
Without detectable residues:	10	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	2	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	12	12	0,02;0,05																0.02	E
Benomyl group(#)	12	12	0.10																0.50	E
Chlorpyrifos	12	12	0,01;0,02																1.00	E
Chlorpyrifos-methyl	12	12	0,01;0,02																0.05	E
Deltamethrin	12	12	0.05																0.10	E
Diazinon	12	12	0,01;0,02																0.50	E
Endosulfan	12	12	0.01																1.00	E
Imazalil	12	12	0,03;0,1																0.02	E
Iprodione	12	10	0,02;0,04			1				1							0.57		5.00	E
Lambda-cyhalothrin	12	12	0.05																0.20	E
Maneb-group(##)	12	12	0,02;0,2																1.00	E
Mecarbam	12	12	0,01;0,02																0.02	E
Metalaxyl	12	12	0,03;0,05																1.00	E
Methamidophos	12	12	0,01;0,03																0.01	E
Methidathion	12	12	0,01;0,02																0.02	E
Permethrin	12	12	0.05																1.00	E
Pirimiphos-methyl	12	12	0,01;0,02																1.00	E
Thiabendazol	12	12	0,05;0,1																0.05	E
Triazophos	12	12	0,01;0,02																0.02	E
Vinclozolin	12	12	0.01																0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Legume vegetables (fresh)		Food item: Peas (frozen or fresh, analysed without pods)	
Reporting country:	<u>Austria</u>	Year of sampling:	<u>2000</u>
Remark:		Only add information regarding peas and the 20 pesticides and do not change or delete rows or columns	
Total number of samples analysed:	12	With residues above MRL (EC+national):	0
Without detectable residues:	11	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	1	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	12	12	0,02;0,05																3.00	E
Benomyl group(#)	12	12	0.10																0.10	E
Chlorpyrifos	12	12	0,01;0,02																0.05	E
Chlorpyrifos-methyl	12	12	0,01;0,02																0.05	E
Deltamethrin	12	12	0.05																0.10	E
Diazinon	12	12	0,01;0,02																0.50	E
Endosulfan	12	12	0.01																1.00	E
Imazalil	12	12	0,03;0,1																0.02	E
Iprodione	12	12	0,02;0,04																0.20	E
Lambda-cyhalothrin	12	12	0.05																0.20	E
Maneb-group(##)	12	12	0,02;0,2																1.00	E
Mecarbam	12	12	0,01;0,02																0.02	E
Metalaxyl	12	12	0,03;0,05																0.05	E
Methamidophos	12	12	0,01;0,03																0.50	E
Methidathion	12	12	0,01;0,02																0.02	E
Permethrin	12	12	0.05																0.10	E
Pirimiphos-methyl	12	12	0,01;0,02																0.50	E
Thiabendazol	12	12	0,05;0,1																0.05	E
Triazophos	12	12	0,01;0,02																0.02	E
Vinclozolin	12	11	0.01		1												0.016		2.00	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Head cabbage	
Reporting country: <u>Portugal</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed:	69	With residues above MRL (EC+national):	0
Without detectable residues:	64	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	5	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	29	27	0,02			1				1								0,21	0	2,00	E
Benomyl group(#)	29	29	0,1																		
Chlorpyrifos	39	39	0,02																		
Chlorpyrifos-methyl	39	39	0,02																		
Deltamethrin	39	39	0,05																		
Diazinon	39	39	0,02																		
Endosulfan	39	38	0,02		1												0,02	0	1,00	E	
Imazalil	29	29	0,1																		
Iprodione	39	39	0,02																		
Lambda-cyhalothrin	39	39	0,02																		
Maneb-group(##)	30	29	0,1					1									0,12	0	1,00	E	
Mecarbam	29	29	0,05																		
Metalaxyl	39	39	0,1																		
Methamidophos	29	28	0,01				1										0,07	0	0,5	E	
Methidathion	39	39	0,02																		
Permethrin	39	39	0,05																		
Pirimiphos-methyl	39	39	0,02																		
Thiabendazol	39	39	0,2																		
Triazophos	29	29	0,02																		
Vinclozolin	39	39	0,02																		

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Rice (husked or polished)	
Reporting country: <u>Finland</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed: <input type="text" value="22"/>	With residues above MRL (EC+national): <input type="text" value="0"/>	Only add information regarding rice and the 20 pesticides and do not change or delete rows or columns	
Without detectable residues: <input type="text" value="18"/>	With residues above EC-MRL: <input type="text" value="0"/>		
With detectable residues at or below MRL or without MRL: <input type="text" value="4"/>	With residues above national MRL: <input type="text" value="0"/>		

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50				
Acephate	22	22	0.05															0.02	E
Benomyl group(#)	14	14	0.10															0.10	E
Chlorpyrifos	22	22	0.01															0.05	E
Chlorpyrifos-methyl	22	22	0.01															3.00	E
Deltamethrin	22	22	0.20															1.00	E
Diazinon	22	22	0.01																W
Endosulfan	22	22	0.01															0.05	E
Imazalil	22	22	0.15															0.02	E
Iprodione	22	22	0.02															3.00	E
Lambda-cyhalothrin	22	22	0.01															0.02	E
Maneb-group(##)	14	14	0.20															0.05	E
Mecarbam	22	22	0.05															0.05	E
Metalaxyl	22	22	0.05															0.05	E
Methamidophos	22	22	0.02															0.01	E
Methidathion	22	22	0.01															0.02	E
Permethrin	22	22	0.10															2.00	E
Pirimiphos-methyl	22	18	0.01			1		3								0.19		5.00	E
Thiabendazol	22	22	0.03																W
Triazophos	22	22	0.02															0.02	E
Vinclozolin	22	22	0.01															0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Cucumber	
Reporting country: <u>Finland</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed:	115	With residues above MRL (EC+national):	0
Without detectable residues:	92	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	23	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	115	115	0.05															0.02	E	
Benomyl group(#)	65	65	0.10															0.50	E	
Chlorpyrifos	115	115	0.01															0.05	E	
Chlorpyrifos-methyl	115	115	0.01															0.05	E	
Deltamethrin	115	115	0.20															0.10	E	
Diazinon	115	115	0.01															0.10	N	
Endosulfan	115	106	0.01	4	2	3											0.05	1.00	E	
Imazalil	115	115	0.15															0.20	E	
Iprodione	115	110	0.02		1		1	2	1								0.22	2.00	E	
Lambda-cyhalothrin	115	115	0.01															0.10	E	
Maneb-group(##)	5	4	0.20					1										0.2	0.50	E
Mecarbam	115	115	0.05															0.05	E	
Metalaxyl	115	110	0.05			1	4											0.09	0.50	E
Methamidophos	115	112	0.02			1	1		1									0.23	1.00	E
Methidathion	115	115	0.01															0.02	E	
Permethrin	115	115	0.10															0.10	E	
Pirimiphos-methyl	115	115	0.01															0.10	E	
Thiabendazol	115	115	0.03																W	
Triazophos	115	115	0.02																W	
Vinclozolin	115	115	0.01															1.00	E	

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Head cabbage	
Reporting country: <u>Finland</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed: <input type="text" value="20"/>	With residues above MRL (EC+national): <input type="text" value="0"/>	Only add information regarding head cabbage and the 20 pesticides and do not change or delete rows or columns	
Without detectable residues: <input type="text" value="19"/>	With residues above EC-MRL: <input type="text" value="0"/>		
With detectable residues at or below MRL or without MRL: <input type="text" value="1"/>	With residues above national MRL: <input type="text" value="0"/>		

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	20	20	0.05																2.00	E
Benomyl group(#)	11	11	0.10																3.00	E
Chlorpyrifos	20	20	0.01																1.00	E
Chlorpyrifos-methyl	20	20	0.01																0.05	E
Deltamethrin	20	20	0.20																0.10	E
Diazinon	20	19	0.01		1												0.02		2.00	N
Endosulfan	20	20	0.01																1.00	E
Imazalil	20	20	0.15																0.02	E
Iprodione	20	20	0.02																5.00	E
Lambda-cyhalothrin	20	20	0.01																0.20	E
Maneb-group(##)	11	11	0.20																1.00	E
Mecarbam	20	20	0.05																0.05	E
Metalaxyl	20	20	0.05																1.00	E
Methamidophos	20	20	0.02																0.50	E
Methidathion	20	20	0.01																0.02	E
Permethrin	20	20	0.10																1.00	E
Pirimiphos-methyl	20	20	0.01																0.05	E
Thiabendazol	20	20	0.03																	W
Triazophos	20	20	0.02																0.10	N
Vinclozolin	20	20	0.01																0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Legume vegetables (fresh)		Food item: Peas (frozen or fresh, analysed without pods)	
Reporting country: <u>Finland</u>	Year of sampling: <u>2000</u>	Remark: Only add information regarding peas and the 20 pesticides and do not change or delete rows or columns	
Total number of samples analysed:	<input type="text" value="21"/>	With residues above MRL (EC+national):	<input type="text" value="0"/>
Without detectable residues:	<input type="text" value="21"/>	With residues above EC-MRL:	<input type="text" value="0"/>
With detectable residues at or below MRL or without MRL:	<input type="text" value="0"/>	With residues above national MRL:	<input type="text" value="0"/>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50				
Acephate	21	21	0.05															0.02	E
Benomyl group(#)	5	5	0.10															0.10	E
Chlorpyrifos	21	21	0.01															0.05	E
Chlorpyrifos-methyl	21	21	0.01															0.05	E
Deltamethrin	21	21	0.20															0.05	E
Diazinon	21	21	0.01															0.20	N
Endosulfan	21	21	0.01															0.50	N
Imazalil	21	21	0.15															0.02	E
Iprodione	21	21	0.02															0.20	E
Lambda-cyhalothrin	21	21	0.01															0.20	E
Maneb-group(##)	5	5	0.20															0.10	E
Mecarbam	21	21	0.05															0.05	E
Metalaxyl	21	21	0.05															0.05	E
Methamidophos	21	21	0.02															0.01	E
Methodathion	21	21	0.01															0.02	E
Permethrin	21	21	0.10															0.05	E
Pirimiphos-methyl	21	21	0.01															0.05	E
Thiabendazol	21	21	0.03															0.05	E
Triazophos	21	21	0.02															0.10	N
Vinclozolin	21	21	0.01															0.30	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereal grains		Food item: Rice	
Reporting country: <u>Sweden</u>		Year of sampling: <u>2000</u>	Remark: Only add information regarding rice and the 20 pesticides and do not change or delete rows or columns
Total number of samples analysed:	41	With residues above MRL (EC+national):	0
Without detectable residues:	39	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	2	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50				
Acephate	0	0																0.02	E
Benomyl group (#)	12	12	0.10															0.10	E
Chlorpyrifos	41	39	0.01		1	1											0.025	0.05	E
Chlorpyrifos-methyl	41	41	0.02															3.00	E
Deltamethrin	41	41	0.20															1.00	E
Diazinon	41	41	0.02															0.05	E
Endosulfan (sum)	41	41	0.02															0.05	E
Imazalil	41	41	0.30															0.02	E
Iprodione	41	41	0.40															3.00	E
Lambda-cyhalothrin	0	0																0.02	E
Maneb group (##)	0	0																0.05	E
Mecarbam	0	0																0.05	E
Metalaxyl	41	41	0.40															0.05	E
Methamidophos	0	0																0.01	E
Methodathion	0	0																0.02	E
Permethrin	41	41	0.90															2.00	E
Pirimiphos-methyl	41	41	0.05															5.00	E
Thiabendazole	41	41	0.20															No MRL	
Triazophos	0	0																0.02	E
Vinclozolin (sum)	41	41	0.06															0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Cucumbers	
Reporting country:	<u>Sweden</u>	Year of sampling:	<u>2000</u>
Remark: Only add information regarding Cucumbers and the 20 pesticides and do not change or delete rows or columns			
Total number of samples analysed:	141	With residues above MRL (EC+national):	0
Without detectable residues:	129	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	12	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)		
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50	
Acephate	141	141	0.02																	0.02	E
Benomyl group (#)	45	45	0.10																	0.50	E
Chlorpyrifos	141	141	0.05																	0.05	E
Chlorpyrifos-methyl	141	141	0.05																	0.05	E
Deltamethrin	141	141	0.05																	0.10	E
Diazinon	141	141	0.02																	0.50	E
Endosulfan (sum)	141	138	0.01		1	2											0.043		1.00	E	
Imazalil	141	141	0.20																0.20	E	
Iprodione	141	139	0.20						2								0.37		2.00	E	
Lambda-cyhalothrin	141	141	0.10																0.10	E	
Maneb group (##)	35	29	0.10					3	3								0.31		0.50	E	
Mecarbam	141	141	0.05																0.05	E	
Metalaxyl	141	140	0.10					1									0.13		No MRL		
Methamidophos	141	138	0.02			1	1		1								0.325		1.00	E	
Methodathion	141	141	0.02																0.02	E	
Permethrin	141	141	0.05																0.10	E	
Pirimiphos-methyl	141	141	0.05																1.00	N	
Thiabendazol	141	141	0,3/0,2																No MRL		
Triazophos	141	141	0.02																No MRL		
Vinclozolin (sum)	141	141	0.05																1.00	E	

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Head Cabbages	
Reporting country: <u>Sweden</u>	Year of sampling: <u>2000</u>	Remark: Only add information regarding head cabbages and the 20 pesticides and do not change or delete rows or columns	
Total number of samples analysed:	<input type="text" value="53"/>	With residues above MRL (EC+national):	<input type="text" value="0"/>
Without detectable residues:	<input type="text" value="53"/>	With residues above EC-MRL:	<input type="text" value="0"/>
With detectable residues at or below MRL or without MRL:	<input type="text" value="0"/>	With residues above national MRL:	<input type="text" value="0"/>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	53	53	0.02																	2.00	E
Benomyl group (#)	22	22	0.10																	3.00	E
Chlorpyrifos	53	53	0.05																	1.00	E
Chlorpyrifos-methyl	53	53	0.05																	0.05	E
Deltamethrin	53	53	0.05																	0.10	E
Diazinon	53	53	0.02																	0.50	E
Endosulfan	53	53	0.01																	1.00	E
Imazalil	53	53	0.20																	0.02	E
Iprodione	53	53	0.20																	5.00	E
Lambda-cyhalothrin	53	53	0.10																	0.20	E
Maneb group (##)	5	5	0.10																	1.00	E
Mecarbam	53	53	0.05																	0.05	E
Metalaxyl	53	53	0.10																	1.00	E
Methamidophos	53	53	0.02																	0.50	E
Methidathion	53	53	0.02																	0.02	E
Permethrin	53	53	0.05																	1.00	E
Pirimiphos-methyl	53	53	0.05																	1.00	N
Thiabendazol	53	53	0,2/0,3																	6.00	N
Triazophos	53	53	0.02																	0.10	N
Vinclozolin	53	53	0.05																	0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Legume Vegetables		Food item: Peas (without pods)	
Reporting country: <u>Sweden</u>	Year of sampling: <u>2000</u>	Remark:	
Total number of samples analysed:	<input type="text" value="14"/>	With residues above MRL (EC+national):	<input type="text" value="0"/>
Without detectable residues:	<input type="text" value="14"/>	With residues above EC-MRL:	<input type="text" value="0"/>
With detectable residues at or below MRL or without MRL:	<input type="text" value="0"/>	With residues above national MRL:	<input type="text" value="0"/>

Only add information regarding peas and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	14	14	0.02																0.02	E
Benomyl group (#)	5	5	0.10																0.10	E
Chlorpyrifos	14	14	0.05																0.05	E
Chlorpyrifos-methyl	14	14	0.05																0.05	E
Deltamethrin	14	14	0.05																0.05	E
Diazinon	14	14	0.02																0.50	E
Endosulfan (sum)	14	14	0.01																1.00	E
Imazalil	14	14	0.20																0.02	E
Iprodione	14	14	0.20																0.20	E
Lambda-cyhalothrin	14	14	0.10																0.20	E
Maneb group (##)	6	6	0.10																0.10	E
Mecarbam	14	14	0.05																0.05	E
Metalaxyl	14	14	0.10																0.05	E
Methamidophos	14	14	0.02																0.01	E
Methidathion	14	14	0.02																0.02	E
Permethrin	14	14	0.05																0.05	E
Pirimiphos-methyl	14	14	0.05																0.05	E
Thiabendazole	14	14	0,3/0,2																0.05	E
Triazophos	14	14	0.02																0.10	N
Vinclozolin (sum)	14	14	0.05																0.30	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Rice (husked or polished)	
Reporting country:	<u>United Kingdom</u>	Year of sampling:	<u>2000</u>
Remark:		Only add information regarding rice and the 20 pesticides and do not change or delete rows or columns	
Total number of samples analysed:	96	With residues above MRL (EC+national):	1
Without detectable residues:	84	With residues above EC-MRL:	1
With detectable residues at or below MRL or without MRL:	11	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	96	96	0.02															0		
Benomyl group(#)	96	96	0.10															0		
Chlorpyrifos	96	96	0.05															0		
Chlorpyrifos-methyl	96	96	0.05															0		
Deltamethrin	84	84	0.05															0		
Diazinon	96	96	0.05															0		
Endosulfan	96	96	0.05															0		
Imazalil	96	96	0.02															0		
Iprodione	96	96	0.05															0		
Lambda-cyhalothrin																				
Maneb-group(##)	96	96	0.05															0		
Mecarbam	96	96	0.05															0		
Metalaxyl																				
Methamidophos	96	96	0.01															0		
Methodathion	96	96	0.02															0		
Permethrin	96	96	0.05															0		
Pirimiphos-methyl	96	93	0.05				1			1	1						1	0	5.00	E
Thiabendazol	96	96	0.05															0		
Triazophos	96	96	0.02															0		
Vinclozolin	96	96	0.05															0		

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Cucumber	
Reporting country:	<u>United Kingdom</u>	Year of sampling:	<u>2000</u>
Remark: Only add information regarding cucumbers and the 20 pesticides and do not change or delete rows or columns			
Total number of samples analysed:	59	With residues above MRL (EC+national):	0
Without detectable residues:	43	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	16	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	59	59	0.02															0			
Benomyl group(#)	59	57	0.10					2										0.4	0	0.50	E
Chlorpyrifos	59	59	0.05															0			
Chlorpyrifos-methyl	59	59	0.05															0			
Deltamethrin	59	59	0.05															0			
Diazinon	59	59	0.02															0			
Endosulfan	59	59	0.05															0			
Imazalil	59	59	0.02															0			
Iprodione	59	59	0.05															0			
Lambda-cyhalothrin	59	59	0.02															0			
Maneb-group(##)	59	56	0.05			2		1									0.3	0	-	W	
Mecarbam	59	59	0.05															0			
Metalaxyl	59	58	0.05		1												0.05	0	-	W	
Methamidophos	59	59	0.01															0			
Methidathion	59	59	0.02															0			
Permethrin	59	59	0.05															0			
Pirimiphos-methyl	59	59	0.05															0			
Thiabendazol	59	59	0.05															0			
Triazophos	59	59	0.02															0			
Vinclozolin	59	59	0.05															0			

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Head cabbage	
Reporting country:	<u>United Kingdom</u>	Year of sampling:	<u>2000</u>
Total number of samples analysed:		<input type="text" value="72"/>	With residues above MRL (EC+national):
Without detectable residues:		<input type="text" value="61"/>	With residues above EC-MRL:
With detectable residues at or below MRL or without MRL:		<input type="text" value="11"/>	With residues above national MRL:
			<input type="text" value="0"/>
			<input type="text" value="0"/>
			<input type="text" value="0"/>

Remark: Only add information regarding head cabbage and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate																					
Benomyl group(#)	72	70	0.10								1	1						2	0	-	W
Chlorpyrifos	72	72	0.05																		
Chlorpyrifos-methyl	72	72	0.05																		
Deltamethrin	72	71	0.05				1											0.07	0	-	W
Diazinon	72	72	0.02																		
Endosulfan	72	72	0.05																		
Imazalil	72	72	0.02																		
Iprodione	72	68	0.05				1	1		2								0.8	0	-	W
Lambda-cyhalothrin	72	72	0.02																		
Maneb-group(##)																					
Mecarbam	72	72	0.05																		
Metalaxyl	72	72	0.05																		
Methamidophos																					
Methidathion	72	72	0.02																		
Permethrin	72	72	0.05																		
Pirimiphos-methyl	72	72	0.05																		
Thiabendazol	72	72	0.05																		
Triazophos	72	72	0.02																		
Vinclozolin	72	72	0.05																		

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg
 (**) in alphabetical order of the English name
 (***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).
 (##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Legume vegetables (fresh)		Food item: Peas (frozen or fresh, analysed without pods)	
Reporting country:	<u>United Kingdom</u>	Year of sampling:	<u>2000</u>
Remark: Only add information regarding peas and the 20 pesticides and do not change or delete rows or columns			
Total number of samples analysed:	62	With residues above MRL (EC+national):	0
Without detectable residues:	59	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	3	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	62	62	0.02																	
Benomyl group(#)	62	62	0.10																	
Chlorpyrifos	62	62	0.05																	
Chlorpyrifos-methyl	62	62	0.05																	
Deltamethrin	62	62	0.05																	
Diazinon	62	62	0.02																	
Endosulfan	62	62	0.05																	
Imazalil	62	62	0.02																	
Iprodione	62	62	0.05																	
Lambda-cyhalothrin	62	62	0.02																	
Maneb-group(##)	62	62	0.05																	
Mecarbam	62	62	0.05																	
Metalaxyl	62	62	0.05																	
Methamidophos	62	62	0.01																	
Methidathion	62	62	0.02																	
Permethrin	62	62	0.05																	
Pirimiphos-methyl	62	62	0.05																	
Thiabendazol	62	62	0.05																	
Triazophos	62	62	0.02																	
Vinclozolin	62	59	0.05				3										0.1	0	0.30	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg
(**) in alphabetical order of the English name
(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).
(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Cucumber	
Reporting country: <u>Norway</u>		Year of sampling: <u>2000</u>	Remark:
Total number of samples analysed:	84	With residues above MRL (EC+national):	0
Without detectable residues:	76	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	8	With residues above national MRL:	0

Only add information regarding cucumbers and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	82	82	0.05																		
Benomyl group(#)	33	32	0.05				1											0.07		0.50	E
Chlorpyrifos	82	82	0.05																		
Chlorpyrifos-methyl	82	82	0.05																		
Deltamethrin	82	82	0.05																		
Diazinon	82	82	0.02																		
Endosulfan	82	81	0.02				1											0.06		1.00	E
Imazalil	82	82	0.02																		
Iprodione	82	77	0.02			2		1	1	1								0.60		2.00	E
Lambda-cyhalothrin	82	82	0.02																		
Maneb-group(##)	22	21	0.05						1									0.47		0.50	E
Mecarbam	82	82	0.05																		
Metalaxyl	82	82	0.05																		
Methamidophos	82	82	0.05																		
Methidathion	82	82	0.05																		
Permethrin	82	82	0.05																		
Pirimiphos-methyl	82	82	0.05																		
Thiabendazole	82	82	0.10																		
Triazophos	82	82	0.05																		
Vinclozolin	82	82	0.02																		

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

(***) E=EC-MRL, N=National MRL, W=without MRL

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Head cabbage	
Reporting country: <u>Norway</u>		Year of sampling: <u>2000</u>	Remark:
Total number of samples analysed:	44	With residues above MRL (EC+national):	0
Without detectable residues:	44	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	0	With residues above national MRL:	0

Only add information regarding head cabbage and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	44	44	0.05																	
Benomyl group(#)	10	10	0.05																	
Chlorpyrifos	44	44	0.05																	
Chlorpyrifos-methyl	44	44	0.05																	
Deltamethrin	44	44	0.05																	
Diazinon	44	44	0.02																	
Endosulfan	44	44	0.02																	
Imazalil	44	44	0.02																	
Iprodione	44	44	0.02																	
Lambda-cyhalothrin	44	44	0.02																	
Maneb-group(##)	15	15	0.05																	
Mecarbam	44	44	0.05																	
Metalaxyl	44	44	0.05																	
Methamidophos	44	44	0.05																	
Methidathion	44	44	0.05																	
Permethrin	44	44	0.05																	
Pirimiphos-methyl	44	44	0.05																	
Thiabendazole	44	44	0.10																	
Triazophos	44	44	0.05																	
Vinclozolin	44	44	0.02																	

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

(***) E=EC-MRL, N=National MRL, W=without MRL

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables

Food item: Cucumber

Reporting country:	Iceland	Year of sampling:	2000	Remark:
Total number of samples analysed:	11	With residues above MRL (EC+national):	0	Only add information regarding cucumbers and the 20 pesticides and do not change or delete rows or columns
Without detectable residues:	11	With residues above EC-MRL:	0	
With detectable residues at or below MRL or without MRL:	0	With residues above national MRL:	0	

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate																					
Benomyl group(#)																					
Chlorpyrifos	11	11	0.05														0	0	0.05	N	
Chlorpyrifos-methyl	11	11	0.05														0	0	0.05	N	
Deltamethrin																					
Diazinon	11	11	0.05														0	0	0.02	N	
Endosulfan	11	11	0.05														0	0	0.05	N	
Imazalil	11	11	0.05														0	0	0.20	N	
Iprodione	11	11	0.05														0	0	2.00	N	
Lambda-cyhalothrin																					
Maneb-group(##)																					
Mecarbam																					
Metalaxyl	11	11	0.05														0	0	0.05	N	
Methamidophos	11	11	0.05														0	0	1.00	N	
Methodathion	11	11	0.05														0	0	0.02	N	
Permethrin	11	11	0.05														0	0	0.10	N	
Pirimiphos-methyl																					
Thiabendazol	11	11	0.05														0	0	0.05	N	
Triazophos																					
Vinclozolin	11	11	0.05														0	0	1.00	N	

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS2

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables

Food item: Head cabbage

Reporting country: <u>Iceland</u>	Year of sampling: <u>2000</u>	Remark:
Total number of samples analysed: <input type="text" value="6"/>	With residues above MRL (EC+national): <input type="text"/>	Only add information regarding rice and the 20 pesticides and do not change or delete rows or columns
Without detectable residues: <input type="text" value="6"/>	With residues above EC-MRL: <input type="text"/>	
With detectable residues at or below MRL or without MRL: <input type="text" value="0"/>	With residues above national MRL: <input type="text"/>	

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate																		0	0	1	N
Benomyl group(#)																		0	0	0.05	N
Chlorpyrifos	6	6	0.05															0	0	0.02	N
Chlorpyrifos-methy	6	6	0.05															0	0	0.02	N
Deltamethrin																		0	0	5.00	N
Diazinon	6	6	0.05															0	0		N
Endosulfan	6	6	0.05															0	0		N
Imazalil	6	6	0.05															0	0		N
Iprodione	6	6	0.05															0	0		N
Lambda-cyhalothrin																					
Maneb-group(##)																					
Mecarbam																					
Metalaxyl	6	6	0.05															0	0	1.00	N
Methamidophos	6	6	0.05															0	0	0.50	N
Methidathion	6	6	0.05															0	0	0.02	N
Permethrin	6	6	0.05															0	0	1.00	N
Pirimiphos-methyl																					
Thiabendazol	6	6	0.05															0	0	0.05	N
Triazophos																					
Vinclozolin	6	6	0.05															0	0	0.05	N

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(**) in alphabetical order of the English name

(##) Sum of dithiocarbamates, expressed as CS2

(***) E=EC-MRL, N=National MRL, W=without MRL

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Rice (husked or polished)	
Reporting country: Liechtenstein		Year of sampling: 2000	
Total number of samples analysed: <input type="text" value="12"/>		With residues above MRL (EC+national): <input type="text" value="1"/>	
Without detectable residues: <input type="text" value="11"/>		With residues above EC-MRL: <input type="text" value="1"/>	
With detectable residues at or below MRL or without MRL: <input type="text" value="0"/>		With residues above national MRL: <input type="text" value="0"/>	
Remark Only add information regarding rice and the 20 pesticides and do not change or delete rows or columns			

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	12	12	0.05																0.02	E
Benomyl group (#)	12	12	0.1																0.1	E
Chlropyriphos	12	12	0.01																0.05	E
Deltamethrin	12	12	0.01																1	E
Diazinon	12	12	0.02																0.02	E
Endosulfan	12	12	0.01																0.05	E
Imazalil	12	12	0.05																0.02	E
Iprodione	12	12	0.05																1	E
Lambda-cyhalothrin	12	12	0.01																0.02	E
Maneb-group (##)	12	12	0.1																0.05	E
Mecarbam	12	11	0.01				1										0.06	1	0.05	E
Metalaxyl	12	12	0.02																0.05	E
Methamidophos	12	12	0.05																0.01	E
Methidathion	12	12	0.02																0.02	E
Permethrin	12	12	0.05																2	E
Pirimiphos-methyl	12	12	0.01																5	E
Thiabendazol	12	12	0.05																0.05	E
Triazophos	12	12	0.02																0.02	E
Vinclozolin	12	12	0.01																0.05	E

(*) i.e. column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E-EC-MRL, N=National MRL, W=Without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS4

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Cucumber	
Reporting country: Liechtenstein		Year of sampling: 2000	
Total number of samples analysed: <input type="text" value="12"/>		With residues above MRL (EC+national): <input type="text" value="0"/>	
Without detectable residues: <input type="text" value="10"/>		With residues above EC-MRL: <input type="text" value="0"/>	
With detectable residues at or below MRL or without MRL: <input type="text" value="2"/>		With residues above national MRL: <input type="text" value="0"/>	
Remark Only add information regarding rice and the 20 pesticides and do not change or delete rows or columns			

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	12	12	0.05															0.02	E	
Benomyl group (#)	12	10	0.05				1	1										0.17	0.5	E
Chlorpyrifos	12	12	0.01																0.05	E
Chlorpyrifos-meth.	12	12	0.01																0.05	E
Deltamethrin	12	12	0.01																0.2	E
Diazinon	12	12	0.02																0.5	N
Endosulfan	12	12	0.01																1	N
Imazalil	12	12	0.05																0.2	E
Iprodione	12	12	0.05																0.2	E
Lambda-cyhalothrin	12	12	0.01																0.5	E
Maneb-group (##)	12	12	0.50																0.5	E
Mecarbam	12	12	0.01																0.2	N
Metalaxyl	12	12	0.02																0.2	N
Methamidophos	12	12	0.05																1	E
Methidathion	12	12	0.02																0.1	N
Permethrin	12	12	0.05																0.5	N
Pirimiphos-methyl	12	12	0.01																0.2	N
Thiabendazol	12	12	0.05																0.2	N
Triazophos	12	12	0.02																0.2	N
Vinclozolin	12	12	0.01																1	N

(*) i.e. column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E-EC-MRL, N=National MRL, W=Without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS4

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Legume vegetables (fresh)		Food item: Peas (frozen or fresh, analysed without pods)	
Reporting country: Liechtenstein		Year of sampling: 2000	
		Remark	
Total number of samples analysed:	12	With residues above MRL (EC+national):	0
Without detectable residues:	12	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	0	With residues above national MRL:	0

Only add information regarding rice and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50				
Acephate	12	12	0.05															0.02	E
Benomyl group (#)	12	12	0.1															0.1	E
Chlorpyrifos	12	12	0.01															0.05	E
Chlorpyrifos-meth.	12	12	0.01															0.05	E
Deltamethrin	12	12	0.01															0.5	E
Diazinon	12	12	0.02															0.02	E
Endosulfan	12	12	0.01															0.05	E
Imazalil	12	12	0.05															0.02	E
Iprodione	12	12	0.05															0.2	E
Lambda-cyhalothrin	12	12	0.01															0.02	E
Maneb-group (##)	12	12	1															0.05	E
Mecarbam	12	12	0.01															0.05	E
Metalaxyl	12	12	0.02															0.05	E
Methamidophos	12	12	0.05															-	E
Methidathion	12	12	0.02															0.02	E
Permethrin	12	12	0.05															0.1	E
Pirimiphos-methyl	12	12	0.01															0.05	E
Thiabendazol	12	12	0.05															0.05	E
Triazophos	12	12	0.02															0.02	E
Vinclozolin	12	12	0.01															-	E

(*) i.e. column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E-EC-MRL, N=National MRL, W=Without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS4