

Notifications of the co-ordinated programme (specific exercise) to the European Commission for all four products											
	Number of samples analysed	Without detectable residues	in %	residues at or below MRL:	in %	With residues above MRL:	in %	residues above EC-MRL:	in %	residues above national	in %
Country											
Belgium	166	148	89.2	9	5.4	9	5.4	3	1.8	6	3.6
Denmark	143	109	76.2	33	23.1	1	0.7	0	0.0	1	0.7
Germany	1079	591	54.8	406	37.6	82	7.6	75	7.0	7	0.6
Greece	171	147	86.0	17	9.9	7	4.1	7	4.1	0	0.0
Spain	177	147	83.1	12	6.8	18	10.2	18	10.2	0	0.0
France	350	208	59.4	125	35.7	17	4.9	17	4.9	0	0.0
Ireland	50	40	80.0	7	14.0	3	6.0	3	6.0	2	4.0
Italy	441	402	91.2	37	8.4	2	0.5	2	0.5	0	0.0
Luxembourg	65	50	76.9	13	20.0	2	3.1	2	3.1	0	0.0
The Netherlands	389	247	63.5	26	6.7	116	29.8	108	27.8	8	2.1
Austria	48	27	56.3	11	22.9	10	20.8	10	20.8	0	0.0
Portugal	300	232	77.3	51	17.0	17	5.7	3	1.0	14	4.7
Finland	345	113	32.8	151	43.8	81	23.5	81	23.5	0	0.0
Sweden	383	270	70.5	77	20.1	36	9.4	36	9.4	0	0.0
United Kingdom	276	227	82.2	41	14.9	8	2.9	8	2.9	0	0.0
Norway	284	255	89.8	27	9.5	2	0.7	0	0.0	2	0.7
Iceland	40	36	90.0	3	7.5	1	2.5	1	2.5	0	0.0
EU + Norway	4707	3249	69.0	1046	22.2	412	8.8	374	7.9	40	0.8

Notifications of the co-ordinated programme (specific exercise) to the European Commission for melon

Pesticide (**)	Total number of samples	Number of samples without residues	Number of samples with detectable residues	%	Number of samples with residues below/equal MRL	%	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)										Maximum (mg/kg)	Number of samples with residues exceeding the MRL	Source of MRL (***)						
								0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10				20	50	>50			
Acephate	758	751	7	99.08	1	0.92	0	1	0	1	3	0	1	0	0	0	0	0	0	0	0	0	0	6		
Benomyl group (****)	541	522	19	96.49	19	3.51	4	0	4	4	7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Chlorpyrifos	844	839	5	99.41	4	0.47	1	1	2	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1		
Chlorpyrifos-methyl	830	830	0	100.00	0	0.00	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Deltamethrin	803	800	3	99.63	3	0.37	0	1	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Diazinon	843	833	10	98.81	10	1.19	2	0	6	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Endosulfan	850	618	232	72.71	232	27.29	27	38	57	60	38	11	1	0	0	0	0	0	0	0	0	0	0	0	0	
Imazalil	789	786	3	99.62	3	0.38	0	1	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	
Iprodione	831	830	1	99.88	1	0.12	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	
Lambda-cyhalothrin	765	762	3	99.61	3	0.39	0	2	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Maneb-group (****)	405	377	28	93.09	26	6.42	0	2	0	14	4	4	2	0	0	0	0	0	0	0	0	0	0	2		
Mecarbam	773	773	0	100.00	0	0.00	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Metaxyl	792	790	2	99.75	2	0.25	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	
Methamidophos	782	749	33	95.78	4	0.51	3	1	13	11	4	1	0	0	0	0	0	0	0	0	0	0	0	0	29	
Methidathion	853	852	1	99.88	1	0.12	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	
Permethrin	820	819	1	99.88	1	0.12	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Pirimiphos-methyl	811	809	2	99.75	2	0.25	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Thiabendazol	597	552	45	92.46	28	4.69	0	1	3	9	5	10	15	2	0	0	0	0	0	0	0	0	0	17		
Triazophos	785	785	0	100.00	0	0.00	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Vinclozolin	838	835	3	99.64	3	0.36	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL

(****) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(*****) Sum of dithiocarbamates, expressed as CS₂.

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission for all four products

Reporting country: EU + Norway	Year of sampling: 1998	in %	in %	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)										Maximum (mg/kg)	Number of samples with residues exceeding the MRL	% samples with residues exceeding the MRL								
				>50	50.0	20.0	10.0	5.0	2.0	1.0	0.5	0.2	0.1				0.05	0.02	0.01					
Pesticide (**)	Total number of samples	Number of samples without residues	%	Number of samples with detectable residues	%	Number of samples with detectable residues	%	Number of samples with residues below/equal MRL	Reporting level (mg/kg)	0.01	0.02	0.05	0.1	0.2	0.5	1.0	2.0	5.0	10.0	20.0	50.0	>50		
Acetophate	3434	3396	98.89	38	1.11	24	0.70	24	0.70	0	5	12	8	3	6	3	1	0	0	0	0	0	14	0.41
Benomyl group (****)	2552	2514	98.51	38	1.49	29	1.14	29	1.14	4	0	6	8	16	2	2	0	0	0	0	0	0	9	0.35
Chlorpyrifos	4211	4178	99.22	33	0.78	31	0.74	31	0.74	6	6	6	7	3	2	1	0	0	0	0	0	0	2	0.05
Chlorpyrifos-methyl	4215	4111	97.53	104	2.47	101	2.40	101	2.40	5	23	29	11	14	16	6	0	0	0	0	0	0	3	0.07
Deltamethrin	3963	3940	99.42	23	0.58	23	0.58	23	0.58	0	6	9	5	3	0	0	0	0	0	0	0	0	0	0.00
Diazinon	4166	4151	99.64	15	0.36	14	0.34	14	0.34	2	1	7	3	2	0	0	0	0	0	0	0	0	1	0.02
Endosulfan	4071	3387	83.20	684	16.80	678	16.65	678	16.65	51	71	123	148	118	130	37	6	0	0	0	0	0	6	0.15
Imazalil	3557	3551	99.83	6	0.17	3	0.08	3	0.08	0	1	0	1	1	2	0	0	1	0	0	0	0	3	0.08
Iprodione	4034	3988	98.86	46	1.14	46	1.14	46	1.14	2	6	16	10	7	3	1	1	0	0	0	0	0	0	0.00
Lambda-cyhalothrin	3614	3606	99.78	8	0.22	8	0.22	8	0.22	1	4	3	0	0	0	0	0	0	0	0	0	0	0	0.00
Maneb-group (****)	2265	2006	88.57	259	11.43	235	10.38	235	10.38	3	14	43	46	46	50	31	21	5	0	0	0	0	24	1.06
Mecarbam	3500	3499	99.97	1	0.03	1	0.03	1	0.03	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0.00
Metaxyl	3663	3659	99.90	4	0.10	1	0.03	1	0.03	1	0	0	2	1	0	0	0	0	0	0	0	0	3	0.08
Methamidophos	3731	3365	90.19	366	9.81	38	1.02	38	1.02	5	17	63	67	64	92	42	11	5	0	0	0	0	328	8.79
Methidathion	3805	3795	99.74	10	0.26	3	0.08	3	0.08	2	1	4	3	0	0	0	0	0	0	0	0	0	7	0.18
Permethrin	3668	3659	99.77	9	0.23	9	0.23	9	0.23	0	0	0	2	4	3	0	0	0	0	0	0	0	0	0.00
Pyrimiphos-methyl	4216	3954	93.79	262	6.21	260	6.17	260	6.17	18	33	63	57	28	26	16	15	6	0	0	0	0	2	0.05
Thiabendazol	3168	3116	98.36	52	1.64	34	1.07	34	1.07	1	1	6	12	5	10	15	2	0	0	0	0	0	18	0.57
Triazophos	3372	3370	99.94	2	0.06	0	0.00	0	0.00	0	0	1	0	0	1	0	0	0	0	0	0	0	2	0.06
Vinclozolin	4046	4024	99.46	22	0.54	22	0.54	22	0.54	3	3	7	6	1	2	0	0	0	0	0	0	0	0	0.00
SUM				1982																				

(*) i.e column 0.02 includes the range from 0.011 ... mg/kg upto 0.020 ... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL

(****) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(*****) Sum of dithiocarbamates, expressed as CS₂.

Overview on the pesticide* sample combinations for the four products																						
Commodities	Total number of pesticide combinations analysed	Number of pesticide combinations without residues	Number of pesticide combinations with detectable residues	Number of pesticide combinations with residues below/equal MRL	Reporting level (mg/kg)	Number of pesticide*sample combinations, in which a residue was found with quantifiable residues in classes up to and including (In mg/kg) (*)										Check sum						
						0	0.02	0.05	0.1	0.2	0.5	1.0	2.0	5.0	10.0		20.0	50.0	>50	Maximum (mg/kg)	Number of pesticide combinations, in which residues exceeded the MRL	%
Cauliflower	16511	16312	199	171	1.0	5	12	37	28	29	35	30	20	3	0	0	0	0	0	0	28	0.17
Peppers	27147	26004	1143	810	3.0	47	89	207	236	202	250	86	20	6	0	0	0	0	0	0	333	1.23
Wheat grains	14683	14441	242	238	1.6	11	43	67	28	24	29	20	14	6	0	0	0	0	0	0	4	0.03
Melon	15310	14912	398	341	2.2	41	49	89	104	61	31	18	3	2	0	0	0	0	0	0	57	0.37
SUM	73651	71669	1982	1560	2.1																422	0.6

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Peppers	
Reporting country:	<u>Belgium</u>	Year of sampling:	<u>1999</u>
Remark: Only add information regarding peppers and the 20 pesticides and do not change or delete rows or columns			
Total number of samples analysed:	58	With residues above MRL (EC+national):	5
Without detectable residues:	49	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	4	With residues above national MRL:	5

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)		
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50	
Acephate	58	58	0,02-0,2																		
Benomyl group(#)	58	57	0,1-0,2				1											0.1	0	0.50	N-E
Chlorpyrifos	58	58	0.05																		
Chlorpyrifos-methyl	58	58	0.05																		
Deltamethrin	58	58	0,05-0,1																		
Diazinon	58	58	0,02-0,05																		
Endosulfan	58	57	0.05					1										0.13	0	2.00	E
Imazalil	58	58	0.1																		
Iprodione	58	58	0,1-0,2																		
Lambda-cyhalothrin	58	58	0.05																		
Maneb-group(##)	58	53	0,5-0,6							4	1							1.8	0	2.00	E
Mecarbam	58	58	0.05																		
Metalaxyl	58	58	0,05-0,1																		
Methamidophos	58	53	0,01-0,02		1		2		2									0.36	5	0.01	N-E
Methidathion	58	58	0,02-0,05																		
Permethrin	58	58	0,05-0,1																		
Pirimiphos-methyl	58	58	0.05																		
Thiabendazol	58	58	0,05-0,1																		
Triazophos	39	39	0.05																		
Vinclozolin	58	58	0.05																		

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B:		Notifications of the co-ordinated programme (specific exercise) to the European Commission																			
Product group: Fruiting vegetables				Food item: Peppers																	
Reporting country: Denmark				Year of sampling: 1999				Remark:													
Total number of samples analysed:				32		With residues above MRL (EC+national):				0		Only add information regarding peppers and the 20 pesticides and do not change or delete rows or columns									
Without detectable residues:				21		With residues above EC-MRL:				0											
With detectable residues at or below MRL or without MRL:				11		With residues above national MRL:				0											
Samples with quantifiable residues in classes up to and including (in mg/kg) (*)																					
Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50	Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
Acephate	32	32	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Benomyl group(##)	32	30	0.05	0	0	0	0	2	0	0	0	0	0	0	0	0	0.15	0	(none)	W	
Chlorpyrifos	32	32	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Chlorpyrifos-methyl	32	32	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Deltamethrin	32	31	0.10	0	0	1	0	0	0	0	0	0	0	0	0	0	0.044	0	0.20	E	
Diazinon	32	32	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Endosulfan	32	22	0.02	0	0	3	2	2	1	2	0	0	0	0	0	0	0.57	0	1.00	E	
Imazail	32	32	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Iprodione	32	32	0.20	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Lambda-cyhalothrin	0																	0			
Maneb-group(##)	31	30	0.10	0	0	0	0	1	0	0	0	0	0	0	0	0	0.12	0	2.00	E	
Mecarbam	32	32	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Metalaxyl	32	32	0.10	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Methamidophos	32	32	0.20	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Methidathion	32	32	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Permethrin	32	32	0.20	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Pirimiphos-methyl	32	31	0.05	0	1	0	0	0	0	0	0	0	0	0	0	0	0.02	0	(none)	W	
Thiabendazol	32	32	0.08	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Triazophos	32	32	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Vinclozolin	32	32	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg								(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).													
(**) in alphabetical order of the English name								(##) Sum of dithiocarbamates, expressed as CS ₂													
(***) E=EC-MRL, N=National MRL, W=without MRL																					

Table B:		Notifications of the co-ordinated programme (specific exercise) to the European Commission																			
Product group: Cereals				Food item: Wheat grains																	
Reporting country:		Denmark		Year of sampling:				1999				Remark:									
Total number of samples analysed:		58		With residues above MRL (EC+national):				0				Only add information regarding wheat grains and the 20 pesticides and do not change or delete rows or columns									
Without detectable residues:		55		With residues above EC-MRL:				0													
With detectable residues at or below MRL or without MRL:		3		With residues above national MRL:				0													
Samples with quantifiable residues in classes up to and including (in mg/kg) (*)																					
Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50	Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
Acephate	0																				
Benomyl group(#)	0																				
Chlorpyrifos	48	48	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0		0			
Chlorpyrifos-methyl	48	45	0.01	0	2	0	0	0	1	0	0	0	0	0	0	0	0.24	0	5.00	E	
Deltamethrin	48	48	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0		0			
Diazinon	48	48	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0		0			
Endosulfan	0																				
Imazalil	0																				
Iprodione	0																				
Lambda-cyhalothrin	48	48	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0		0			
Maneb-group(##)	0																				
Mecarbam	0																				
Metalaxyl	0																				
Methamidophos	0																				
Methidathion	0																				
Permethrin	0																				
Pirimiphos-methyl	48	48	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0		0			
Thiabendazol	0																				
Triazophos	0																				
Vinclozolin	48	48	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0		0			
(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg										(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).											
(**) in alphabetical order of the English name										(##) Sum of dithiocarbamates, expressed as CS ₂											
(***) E=EC-MRL, N=National MRL, W=without MRL																					

Table B:		Notifications of the co-ordinated programme (specific exercise) to the European Commission																		
Product group: Fruiting vegetables			Food item: Melon, not squash or watermelon																	
Reporting country:		Denmark			Year of sampling:					1999					Remark:					
Total number of samples analysed:		30			With residues above MRL (EC+national):					1					Only add information regarding melon and the 20 pesticides and do not change or delete rows or columns					
Without detectable residues:		11			With residues above EC-MRL:					0										
With detectable residues at or below MRL or without MRL:		18			With residues above national MRL:					1										
Samples with quantifiable residues in classes up to and including (in mg/kg) (*)																				
Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50	Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
Acephate	30	30	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.50	E
Benomyl group(##)	30	28	0.05	0	0	0	1	1	0	0	0	0	0	0	0	0	0.12	0	0.50	E
Chlorpyrifos	30	30	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Chlorpyrifos-methyl	30	30	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Deltamethrin	30	30	0.10	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Diazinon	30	28	0.05	1	0	1	0	0	0	0	0	0	0	0	0	0	0.04	0	0.50	E
Endosulfan	30	16	0.02	0	3	3	5	1	2	0	0	0	0	0	0	0	0.28	0	1.00	E
Imazalil	30	29	0.05	0	1	0	0	0	0	0	0	0	0	0	0	0	0.02	0	2.00	E
Iprodione	30	30	0.20	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Lambda-cyhalothrin	0																			
Maneb-group(##)	30	28	0.10	0	0	0	0	1	0	0	0	1	0	0	0	0	2.9	1	1.00	N
Mecarbam	30	30	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Metalaxyl	30	30	0.10	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Methamidophos	30	30	0.20	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Methidathion	30	30	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Permethrin	30	30	0.20	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Pirimiphos-methyl	30	30	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Thiabendazol	30	25	0.08	0	0	0	0	0	2	2	1	0	0	0	0	0	1.4	0	(none)	W
Triazophos	30	30	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Vinclozolin	30	30	0.05	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(##) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(**) in alphabetical order of the English name

(###) Sum of dithiocarbamates, expressed as CS₂

(***) E=EC-MRL, N=National MRL, W=without MRL

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Cauliflower (fresh or frozen)	
Reporting country: <u>Germany</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed:	296	With residues above MRL (EC+national):	27
Without detectable residues:	136	With residues above EC-MRL:	27
With detectable residues at or below MRL or without MRL:	133	With residues above national MRL:	0

Only add information regarding Cauliflower and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	261	261	0.02														0.02		E	
Benomyl group(#)	246	240				1	1	1	2	1							0.6	4	0.10	E
Chlorpyrifos	277	276	0.01	1													0.01		0.05	E
Chlorpyrifos-methyl	278	278	0.01																0.05	E
Deltamethrin	278	278	0.05																0.10	E
Diazinon	278	275	0.02			1	1	1									0.18		0.50	E
Endosulfan	278	278																	1.00	E
Imazalil	261	260	0.10				1										0.1	1	0.02	E
Iprodione	278	277	0.02			1											0.03		0.05	E
Lambda-cyhalothrin	256	256	0.02																0.02	N
Maneb-group(##)	263	111	0.02	2	9	26	20	22	31	20	19	3					2.6	22	1.00	E
Mecarbam	278	278	0.02																0.05	E
Metalaxyl	272	272	0.05																0.05	N
Methamidophos	261	261	0.05																0.50	E
Methodathion	278	278	0.02																0.02	E
Permethrin	278	278	0.05																0.10	E
Pirimiphos-methyl	272	272	0.01																1.00	E
Thiabendazol	269	265	0.10	1		3											0.05		0.05	N
Triazophos	278	278	0.02																0.20	N
Vinclozolin	277	275	0.01	1		1											0.031		0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Peppers	
Reporting country:	<u>Germany</u>	Year of sampling:	<u>1999</u>
Remark:			
Only add information regarding peppers and the 20 pesticides and do not change or delete rows or columns			
Total number of samples analysed:	345	With residues above MRL (EC+national):	44
Without detectable residues:	182	With residues above EC-MRL:	44
With detectable residues at or below MRL or without MRL:	119	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)		
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50	
Acephate	301	299	0.02			1	1											0.062	2	0.02	E
Benomyl group(#)	37	37																		0.10	E
Chlorpyrifos	338	333	0.01	2		2			1									0.29		0.50	E
Chlorpyrifos-methyl	339	334	0.01		4			1										0.15		0.50	E
Deltamethrin	320	315	0.05		3		2											0.061		0.20	E
Diazinon	339	339	0.02																	0.50	E
Endosulfan	339	214		6	7	27	24	19	34	6	2							1.3	2	1.00	E
Imazalil	311	311	0.10																	0.02	E
Iprodione	322	302	0.02	2	4	5	6	3										0.19		5.00	E
Lambda-cyhalothrin	316	316	0.02																	0.02	N
Maneb-group(##)	273	226	0.02	1	2	15	9	11	8	1								0.67		2.00	E
Mecarbam	338	338	0.02																	0.05	E
Metalaxyl	309	309	0.05																	0.05	N
Methamidophos	305	265	0.05		1	6	6	13	12	1	1							1.6	40	0.01	E
Methidathion	339	338	0.02			1												0.035	1	0.02	E
Permethrin	327	327	0.05																	0.50	E
Pirimiphos-methyl	334	308	0.01	2	2	5	8	3	6									0.37		0.50	N
Thiabendazol	39	39	0.05																	0.05	N
Triazophos	327	326	0.02			1												0.03	1	0.02	E
Vinclozolin	333	328	0.01	1	2		2											0.099		3.00	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Wheat grains	
Reporting country:	<u>Germany</u>	Year of sampling:	<u>1999</u>
Remark:			
Only add information regarding wheat grains and the 20 pesticides and do not change or delete rows or columns			
Total number of samples analysed:	181	With residues above MRL (EC+national):	1
Without detectable residues:	159	With residues above EC-MRL:	1
With detectable residues at or below MRL or without MRL:	21	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	56	56	0.02																0.02	E
Benomyl group(#)	114	114																	0.10	E
Chlorpyrifos	89	89	0.01																0.05	E
Chlorpyrifos-methyl	175	175	0.01																3.00	E
Deltamethrin	174	173	0.05					1									0.12		1.00	E
Diazinon	89	89	0.02																0.05	E
Endosulfan	175	174		1													0.002		0.10	E
Imazalil	45	44	0.05						1								0.24	1	0.02	E
Iprodione	174	173	0.04					1									0.16		0.50	E
Lambda-cyhalothrin	173	173	0.04																0.02	E
Maneb-group(##)	1	1	0.02																1.00	E
Mecarbam	50	50	0.03																0.05	E
Metalaxyl	69	69	0.09																0.05	E
Methamidophos	42	42	0.05																0.01	E
Methidathion	58	57	0.02	1													0.005		0.02	E
Permethrin	82	82	0.002																2.00	E
Pirimiphos-methyl	169	150	0.02	2	6	7	1		2		1						1.1		5.00	E
Thiabendazol	115	115	0.05																0.20	N
Triazophos	82	82	0.011																0.02	N
Vinclozolin	89	89	0.01																	

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(**) in alphabetical order of the English name

(##) Sum of dithiocarbamates, expressed as CS₂

(***) E=EC-MRL, N=National MRL, W=without MRL

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Melon, not squash or watermelon	
Reporting country:	<u>Germany</u>	Year of sampling:	<u>1999</u>
Remark:			
Only add information regarding melon and the 20 pesticides and do not change or delete rows or columns			
Total number of samples analysed:	257	With residues above MRL (EC+national):	10
Without detectable residues:	114	With residues above EC-MRL:	3
With detectable residues at or below MRL or without MRL:	133	With residues above national MRL:	7

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	216	215	0.02		1													0.02		0.02	E
Benomyl group(#)	53	45		4		2	1	1										0.15		0.50	E
Chlorpyrifos	256	253	0.01	1	1	1												0.05		0.05	E
Chlorpyrifos-methyl	255	255	0.01																	0.05	E
Deltamethrin	254	251	0.05		1	2												0.05		0.05	E
Diazinon	256	254	0.02	1		1												0.023		0.50	E
Endosulfan	256	122		26	28	34	26	16	3	1								0.57		1.00	E
Imazalil	238	236	0.10					1	1									0.29		2.00	E
Iprodione	254	254	0.02																	0.30	E
Lambda-cyhalothrin	254	252	0.02		2													0.014		0.02	N
Maneb-group(##)	21	20	0.02		1													0.015		0.50	E
Mecarbam	255	255	0.02																	0.05	N
Metalaxyl	238	237	0.05	1														0.002		0.05	N
Methamidophos	221	218	0.05		1	1		1										0.15	3	0.01	E
Methidathion	256	256	0.02																	0.02	E
Permethrin	254	254	0.05																	0.10	E
Pirimiphos-methyl	247	245	0.01	2														0.001		0.50	N
Thiabendazol	58	47	0.05		1	3	2	3	2									0.47	7	0.05	N
Triazophos	254	254	0.02																	0.02	N
Vinclozolin	256	256	0.01																	1.00	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Wheat grains	
Reporting country:	<u>GREECE</u>	Year of sampling:	<u>1999</u>
Remark:		Only add information regarding wheat grains and the 20 pesticides and do not change or delete rows or columns	
Total number of samples analysed:	<input type="text" value="30"/>	With residues above MRL (EC+national):	<input type="text" value="0"/>
Without detectable residues:	<input type="text" value="30"/>	With residues above EC-MRL:	<input type="text" value="0"/>
With detectable residues at or below MRL or without MRL:	<input type="text"/>	With residues above national MRL:	<input type="text" value="0"/>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50				
Acephate	30	30																	
Benomyl group(#)	0	0																	
Chlorpyrifos	30	30																	
Chlorpyrifos-methyl	30	30																	
Deltamethrin	30	30																	
Diazinon	30	30																	
Endosulfan	30	30																	
Imazalil	30	30																	
Iprodione	30	30																	
Lambda-cyhalothrin	30	30																	
Maneb-group(##)	0	0																	
Mecarbam	30	30																	
Metalaxyl	30	30																	
Methamidophos	30	30																	
Methodathion	30	30																	
Permethrin	30	30																	
Pirimiphos-methyl	30	30																	
Thiabendazol	30	30																	
Triazophos	30	30																	
Vinclozolin	30	30																	

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables **Food item:** Cauliflower (fresh or frozen)

Reporting country: SPAIN **Year of sampling:** 1999 **Remark:** Only add information regarding Cauliflower and the 20 pesticides and do not change or delete rows or columns

Total number of samples analysed:

42

1

Without detectable residues:

36

1

With detectable residues at or below MRL:

5

0

or without MRL:

5

0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)										Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)			
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10					20	50	>50
Acceptate	42	40	0.02													0.08		2.00	E	
Benomyl group(#)	42	42	0.10																	
Chlorpyrifos	42	41	0.05										1			0.06	1	0.05	E	
Chlorpyrifos-methyl	42	42	0.05																	
Deltamethrin	42	42	0.05																	
Diazinon	42	41	0.02										1			0.02		0.50	E	
Endosulfan	42	41	0.05													0.72		1.00	E	
Imazalil	42	42	0.02																	
Iprodione	42	42	0.02																	
Lambda-cyhalothrin	42	42	0.02																	
Maneb-group(##)	42	41	0.05											1				1.00	E	
Mecarbam	42	42	0.05																	
Metaxyl	42	42	0.05																	
Methamidophos	42	42	0.01																	
Methidathion	42	42	0.02																	
Permethrin	42	42	0.05																	
Pirimiphos-methyl	42	42	0.05																	
Thiabendazol	42	42	0.05																	
Triazophos	42	42	0.02																	
Vinclozolin	42	42	0.05																	

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS2

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Peppers	
Reporting country:	SPAIN	Year of sampling:	1999
Total number of samples analysed:	45	With residues above MRL (EC+national):	11
Without detectable residues:	30	With residues above EC-MRL:	11
With detectable residues at or below MRL or without MRL:	4	With residues above national MRL:	0

Remark:
Only add information regarding peppers and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)										Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)		
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10					20	50
Acephate	45	45	0.02																
Benomyl group(#)	45	45	0.10																
Chlorpyrifos	45	45	0.05																
Chlorpyrifos-methyl	45	45	0.05																
Deltamethrin	45	45	0.05																
Diazinon	45	45	0.02																
Endosulfan	45	42	0.05							1	1	1							E
Imazalil	45	45	0.02																
Iprodione	45	45	0.02																
Lambda-cyhalothrin	45	45	0.02																
Maneb-group(##)	45	43	0.05							1	1								E
Mecarbam	45	45	0.05																
Metalaxyl	45	45	0.05																
Methamidophos	45	35	0.01																
Metidathion	45	45	0.02																
Permethrin	45	45	0.05																
Pirimiphos-methyl	45	45	0.05																
Thiabendazol	45	45	0.05																
Triazophos	45	45	0.02																
Vinclozolin	45	45	0.05																

(*) i.e column 0.02 includes the range from 0.011 ... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS2

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Wheat grains																
Reporting country:	SPAIN	Year of sampling: 1999																
Total number of samples analysed:	45	With residues above MRL (EC+national):																
Without detectable residues:	43	With residues above EC-MRL:																
With detectable residues at or below MRL or without MRL:	2	With residues above national MRL:																
Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)										Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10					20
Acephate	45	45	0.02															
Benomyl group(#)	45	45	0.10															
Chlorpyrifos	45	45	0.05															
Chlorpyrifos-methyl	45	45	0.05															
Deltamethrin	45	45	0.05															
Diazinon	45	45	0.02															
Endosulfan	45	45	0.05															
Imazalil	45	45	0.02															
Iprodione	45	45	0.02															
Lambda-cyhalothrin	45	45	0.02															
Maneb-group(##)	45	45	0.05															
Mecarbam	45	45	0.05															
Metaxyl	45	45	0.05															
Methamidophos	45	45	0.01															
Methidathion	45	45	0.02															
Permethrin	45	45	0.05															
Pirimiphos-methyl	45	43	0.05							1							0.37	5.00
Thiabendazol	45	45	0.05															E
Triazophos	45	45	0.02															
Vinclozolin	45	45	0.05															

(*) i.e column 0.02 includes the range from 0.011 ... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim)

(##) Sum of dithiocarbamates, expressed as CSZ

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables **Food item:** Melon, not squash or watermelon

Reporting country: SPAIN **Year of sampling:** 1999 **Remark:** Only add information regarding melon and the 20 pesticides and do not change or delete rows or columns

Total number of samples analysed: 45
 Without detectable residues: 38
 With detectable residues at or below MRL or without MRL: 1

With residues above MRL (EC+national): 6
 With residues above EC-MRL: 6
 With residues above national MRL: 0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)										Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10					20
Acephate	45	45	0.02															
Benomyl group(#)	45	45	0.10															
Chlorpyrifos	45	45	0.05															
Chlorpyrifos-methyl	45	45	0.05															
Deltamethrin	45	45	0.05															
Diazinon	45	45	0.02															
Endosulfan	45	45	0.05															
Imazalil	45	45	0.02															
Iprodione	45	45	0.02															
Lambda-cyhalothrin	45	45	0.02															
Maneb-group(##)	45	44	0.05				1											
Mecarbam	45	45	0.05															
Metaxyl	45	45	0.05															
Methamidophos	45	39	0.01		1	3	2											
Methidathion	45	45	0.02															
Permethrin	45	45	0.05															
Pirimiphos-methyl	45	45	0.05															
Thiabendazol	45	45	0.05															
Triazophos	45	45	0.02															
Vinclozolin	45	45	0.05															

(*) i.e column 0.02 includes the range from 0.011 ... mg/kg upto 0.020... mg/kg
 (**) in alphabetical order of the English name
 (***) E=EC-MRL, N=National MRL, W=without MRL
 (#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).
 (##) Sum of dithiocarbamates, expressed as CSZ

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Cauliflower (fresh or frozen)	
Reporting country:	<u>France</u>	Year of sampling:	<u>1999</u>
Total number of samples analysed:	66	With residues above MRL (EC+national):	0
Without detectable residues:	60	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	6	With residues above national MRL:	0

Remark:
Only add information regarding Cauliflower and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	66	66	0.01														0.01	0	2	E
Benomyl group(#)	66	66	0.02														0.02	0	0.1	E
Chlorpyrifos	66	66	0.01														0.01	0	0.05	E
Chlorpyrifos-methyl	66	66	0.01														0.01	0	0.05	E
Deltamethrin	66	66	0.01														0.01	0	0.1	E
Diazinon	66	66	0.01														0.01	0	0.5	E
Endosulfan	66	66	0.02														0.02	0	1	E
Imazalil	44	44	0.05														0.05	0	0.05	E
Iprodione	66	66	0.01														0.01	0	0.05	E
Lambda-cyhalothrin	66	66	0.01														0.01	0	0.02	E
Maneb-group(##)	66	62	0.05				1	2		1							0.75	0	0,5/1	E
Mecarbam	66	66	0.01														0.01	0	0.05	E
Metalaxyl	66	66	0.01														0.01	0	0.05	E
Methamidophos	66	66	0.01														0.01	0	0.5	E
Methidathion	66	66	0.01														0.01	0	0.02	E
Permethrin	66	66	0.02														0.02	0	0.1	E
Pirimiphos-methyl	66	66	0.01														0.01	0	1	E
Thiabendazol	66	64	0.02				2										0.08	0	0,05/5	E
Triazophos	66	66	0.02														0.02	0	0.1	E
Vinclozolin	66	66	0.01														0.01	0	0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Peppers	
Reporting country: <u>France</u>		Year of sampling: <u>1999</u>	Remark:
Total number of samples analysed:	54	With residues above MRL (EC+national):	16
Without detectable residues:	22	With residues above EC-MRL:	16
With detectable residues at or below MRL or without MRL:	16	With residues above national MRL:	0

Only add information regarding peppers and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Accephate	54	54	0.01														0.01	0	0.02	E
Benomyl group(#)	54	53	0.02				1										0.09	0	0.1	E
Chlorpyrifos	54	50	0.01		1	2	1										0.06	0	0.5	E
Chlorpyrifos-methyl	54	54	0.01														0.01	0	0.5	E
Deltamethrin	54	54	0.01														0.01	0	0.2	E
Diazinon	54	54	0.01														0.01	0	0.5	E
Endosulfan	54	46	0.02		1	1	4		2								0.24	0	0,05/1	E
Imazalil	54	54	0.05														0.05	0	0.02	E
Iprodione	54	54	0.01														0.01	0	5	E
Lambda-cyhalothrin	54	54	0.01														0.01	0	0,02/0,1	E
Maneb-group(##)	54	54	0.05														0.05	0	2	E
Mecarbam	54	54	0.01														0.01	0	0.05	E
Metalaxyl	54	54	0.01														0.01	0	0.05	E
Methamidophos	54	37	0.01	1	2	5	2	3	3		1						2.2	15	0.01	E
Methidathion	54	54	0.01														0.01	0	0.02	E
Permethrin	54	54	0.02														0.02	0	0.5	E
Pirimiphos-methyl	54	47	0.01	1	1	3	1	1									0.17	1	0.05	E
Thiabendazol	54	54	0.02														0.02	0	0.05	E
Triazophos	54	54	0.02														0.02	0	0.02	E
Vinclozolin	54	53	0.01						1								0.24	0	3	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(**) in alphabetical order of the English name

(##) Sum of dithiocarbamates, expressed as CS₂

(***) E=EC-MRL, N=National MRL, W=without MRL

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Wheat grains	
Reporting country: <u>France</u>		Year of sampling: <u>1999</u>	Remark:
Total number of samples analysed:	173	With residues above MRL (EC+national):	0
Without detectable residues:	91	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	82	With residues above national MRL:	0

Only add information regarding wheat grains and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	66	66	0.01														0.01	0	0.02	E
Benomyl group(#)	66	66	0.02														0.02	0	0.1	E
Chlorpyrifos	173	173	0.01														0.01	0	0.05	E
Chlorpyrifos-methyl	173	125	0.01	2	10	18	1	5	11	1							0.93	0	3	E
Deltamethrin	173	164	0.01		2	4	1	2									0.17	0	1	E
Diazinon	173	173	0.01														0.01	0	0.05	E
Endosulfan	173	173	0.02														0.02	0	0.1	E
Imazalil	66	66	0.05														0.05	0	0.02	E
Iprodione	173	173	0.01														0.01	0	0.5	E
Lambda-cyhalothrin	173	170	0.01	1	2												0.015	0	0.02	E
Maneb-group(##)	66	66	0.05														0.05	0	1	E
Mecarbam	173	173	0.01														0.01	0	0.05	E
Metalaxyl	173	173	0.01														0.01	0	0.05	E
Methamidophos	173	173	0.01														0.01	0	0.01	E
Methodathion	173	173	0.01														0.01	0	0.02	E
Permethrin	173	173	0.02														0.02	0	2	E
Pirimiphos-methyl	173	103	0.01	4	14	26	3	3	4	7	6	3					2.13	0	5	E
Thiabendazol	66	66	0.02														0.02	0	0.05	E
Triazophos	66	66	0.02														0.02	0	0.02	E
Vinclozolin	173	173	0.01														0.01	0	0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Melon, not squash or watermelon	
Reporting country: <u>France</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed:	57	With residues above MRL (EC+national):	1
Without detectable residues:	35	With residues above EC-MRL:	1
With detectable residues at or below MRL or without MRL:	21	With residues above national MRL:	0

Only add information regarding melon and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)		
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50	
Acephate	57	56	0.01														1	1.39	1	0.02	E
Benomyl group(#)	57	55	0.02					2										0.095	0	0.5	E
Chlorpyrifos	57	57	0.01															0.01	0	0.05	E
Chlorpyrifos-methyl	57	57	0.01															0.01	0	0.05	E
Deltamethrin	57	57	0.01															0.01	0	0.05	E
Diazinon	57	57	0.01															0.01	0	0.5	E
Endosulfan	57	48	0.02		4	4	1											0.06	0	1	E
Imazalil	57	57	0.05															0.05	0	2	E
Iprodione	57	57	0.01															0.01	0	0.3	E
Lambda-cyhalothrin	57	56	0.01			1												0.02	0	0.05	E
Maneb-group(##)	57	50	0.05				4	2	1									0.22	0	0.5	E
Mecarbam	57	57	0.01															0.01	0	0.05	E
Metalaxyl	57	57	0.01															0.01	0	0.05	E
Methamidophos	57	56	0.01				1											0.05	1	0.01	E
Methodathion	57	57	0.01															0.01	0	0.02	E
Permethrin	57	57	0.02															0.02	0	0.1	E
Pirimiphos-methyl	57	57	0.01															0.01	0		E
Thiabendazol	57	57	0.02															0.02	0	0.05	E
Triazophos	57	57	0.02															0.02	0		E
Vinclozolin	57	55	0.01	1	1													0.015	0	1	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg
 (**) in alphabetical order of the English name
 (***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).
 (##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Wheat grains	
Reporting country:	<u>Ireland</u>	Year of sampling:	<u>1999</u>
Remark:		Only add information regarding wheat grains and the 20 pesticides and do not change or delete rows or columns	
Total number of samples analysed:	<u>13</u>	With residues above MRL (EC+national):	<u>0</u>
Without detectable residues:	<u>13</u>	With residues above EC-MRL:	<u>0</u>
With detectable residues at or below MRL or without MRL:	<u>0</u>	With residues above national MRL:	<u>0</u>

Pesticide (**)	Total number of samples	Number of samples without residues	Lowest Calibrated Level (lcl) (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)											Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20					50
Acephate	13	13	0.05																
Benomyl group(#)	13	13	0.10																
Chlorpyrifos	13	13	0.05																
Chlorpyrifos-methyl	13	13	0.05																
Deltamethrin	13	13	0.05																
Diazinon	13	13	0.02																
Endosulfan	13	13	0.02 for each component of the residue.																
Imazalil																			
Iprodione	13	13	0.10																
Lambda-Cyhalothrin	13	13	0.02																
Maneb-group(##)																			
Mecarbam																			
Metalaxyl																			
Methamidophos	13	13	0.05																
Methodathion	13	13	0.02																
Permethrin	13	13	0.20																
Pirimiphos-methyl	13	13	0.05																
Thiabendazol	13	13	0.05																
Triazophos	13	13	0.05																
Vinclozolin	13	13	0.02																

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg
 (**) in alphabetical order of the English name
 (***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).
 (##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Melon, not squash or watermelon	
Reporting country: <u>Ireland</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed:	12	With residues above MRL (EC+national):	1
Without detectable residues:	5	With residues above EC-MRL:	1
With detectable residues at or below MRL or without MRL:	6	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Lowest Calibrated Level (lcl) (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	12	12	0.05														0.14	0	0.50	E
Benomyl group(#)	12	11	0.10					1												
Chlorpyrifos	12	12	0.05																	
Chlorpyrifos-methyl	12	12	0.05																	
Deltamethrin	12	12	0.05																	
Diazinon	12	12	0.02																	
Endosulfan	12	7	0.02 for each compon	1	2	1	1										0.2	0	0.10	E
Imazalil																				
Iprodione	12	12	0.10																	
Lambda-cyhalothrin	12	12	0.02																	
Maneb-group(##)																				
Mecarbam																				
Metalaxyl																				
Methamidophos	12	11	0.05			1											0.04	1	0.01	E
Methodathion	12	12	0.02																	
Permethrin	12	12	0.20																	
Pirimiphos-methyl	12	12	0.05																	
Thiabendazol	12	11	0.05				1										0.08	0		
Triazophos	12	12	0.05																	
Vinclzolin	12	12	0.02																	

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Cauliflower (fresh or frozen)	
Reporting country: <u>ITALY</u>		Year of sampling: <u>1999</u>	Remark:
Total number of samples analysed:	74	With residues above MRL (EC-national):	0
Without detectable residues:	70	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	4	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50				
Acephate	48	48															n.d.	2	E
Benomyl group(#)	37	37															n.d.	0,1	E
Chlorpyrifos	67	67															n.d.	0,05	E
Chlorpyrifos-methyl	61	61															n.d.	0,05	E
Deltamethrin	45	45															n.d.	0,1	E
Diazinon	66	66															n.d.	0,5	E
Endosulfan	56	56															n.d.	1	E
Imazalil	51	51															n.d.	0,02	E
Iprodione	55	55															n.d.	0,05	E
Lambda-cyhalothrin	24	24															n.d.	0,05	N
Maneb-group(##)	57	53						1	2	1							1,9	2	E
Mecarbam	32	32															n.d.	0,05	E
Metalaxyl	42	42															n.d.	0,5	N
Methamidophos	49	49															n.d.	0,5	E
Methodathion	65	65															n.d.	0,02	E
Permethrin	45	45															n.d.	0,1	E
Pirimiphos-methyl	63	63															n.d.	1	E
Thiabendazol	38	38															n.d.	0,05	E
Triazophos	41	41															n.d.	0,1	N
Vinclozolin	65	65															n.d.	0,05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Peppers	
Reporting country:	<u>ITALY</u>	Year of sampling:	<u>1999</u>
Remark:		Only add information regarding peppers and the 20 pesticides and do not change or delete rows or columns	
Total number of samples analysed:	<u>144</u>	With residues above MRL (EC+national):	<u>2</u>
Without detectable residues:	<u>127</u>	With residues above EC-MRL:	<u>2</u>
With detectable residues at or below MRL or without MRL:	<u>15</u>	With residues above national MRL:	<u>0</u>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	80	80															n.d.		0,02	E
Benomyl group(#)	72	71						1									0,13	1	0,1	E
Chlorpyrifos	120	119					1										0,06		0,5	E
Chlorpyrifos-methyl	105	104				1											0,04		0,5	E
Deltamethrin	88	88															n.d.		0,2	E
Diazinon	112	112															n.d.		0,5	E
Endosulfan	102	95			1		1	1	3	1							0,69		1	E
Imazalil	89	89									1						n.d.		0,02	E
Iprodione	107	106									1						0,84		5	E
Lambda-cyhalothrin	28	28															n.d.		0,02	N
Maneb-group(##)	104	101					1		2								0,5		2	E
Mecarbam	57	57															n.d.		0,05	E
Metalaxyl	92	92															n.d.		1	N
Methamidophos	91	90							1								0,5	1	0,01	E
Methidathion	112	112															n.d.		0,02	E
Permethrin	58	58															n.d.		0,5	E
Pirimiphos-methyl	108	106				2											0,05		0,5	N
Thiabendazol	85	85															n.d.		0,05	N
Triazophos	49	49															n.d.		0,02	E
Vinclozolin	120	120															n.d.		1	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Wheat grains	
Reporting country:	<u>ITALY</u>	Year of sampling:	<u>1999</u>
Remark:			
Only add information regarding wheat grains and the 20 pesticides and do not change or delete rows or columns			
Total number of samples analysed:	166	With residues above MRL (EC+national):	0
Without detectable residues:	151	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	15	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	37	37															n.d.		0,02	E
Benomyl group(#)	20	20															n.d.		0,1	E
Chlorpyrifos	116	116															n.d.		0,05	E
Chlorpyrifos-methyl	67	64			1		1		1								0,28		2,5	E
Deltamethrin	35	35															n.d.		1	E
Diazinon	79	79															n.d.		0,05	E
Endosulfan	59	59															n.d.		0,1	E
Imazalil	18	18															n.d.		0,02	E
Iprodione	43	43															n.d.		0,5	E
Lambda-cyhalothrin	2	2															n.d.		0,02	E
Maneb-group(##)	14	14															n.d.		1	E
Mecarbam	14	14															n.d.		0,05	E
Metalaxyl	41	41															n.d.		0,05	E
Methamidophos	29	29															n.d.		0,01	E
Methodathion	72	72															n.d.		0,02	E
Permethrin	34	33						1									0,15		2	E
Pirimiphos-methyl	126	115			2	2	1		4								3,2		5	E
Thiabendazol	25	25															n.d.		0,2	N
Triazophos	8	8															n.d.		0,02	N
Vinclozolin	52	52															n.d.		0,05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Melon, not squash or watermelon	
Reporting country:	<u>ITALY</u>	Year of sampling:	<u>1999</u>
		Remark:	Only add information regarding melon and the 20 pesticides and do not change or delete rows or columns
Total number of samples analysed:	<input type="text" value="57"/>	With residues above MRL (EC+national):	<input type="text" value="0"/>
Without detectable residues:	<input type="text" value="54"/>	With residues above EC-MRL:	<input type="text" value="0"/>
With detectable residues at or below MRL or without MRL:	<input type="text" value="3"/>	With residues above national MRL:	<input type="text" value="0"/>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	33	33															n.d.		0,02	E
Benomyl group(#)	32	32															n.d.		0,5	E
Chlorpyrifos	55	55															n.d.		0,05	E
Chlorpyrifos-methyl	42	42															n.d.		0,05	E
Deltamethrin	29	29															n.d.		0,05	E
Diazinon	54	54															n.d.		0,5	E
Endosulfan	48	46				2											0,1		1	E
Imazalil	41	41															n.d.		2	E
Iprodione	44	44															n.d.		0,3	E
Lambda-cyhalothrin	21	21															n.d.		0,02	N
Maneb-group(##)	30	30															n.d.		0,05	E
Mecarbam	21	21															n.d.		0,05	E
Metalaxyl	33	33															n.d.		0,5	N
Methamidophos	39	38		1													0,004		0,01	E
Methodathion	51	51															n.d.		0,02	E
Permethrin	33	33															n.d.		0,1	E
Pirimiphos-methyl	44	44															n.d.		0,05	N
Thiabendazol	41	41															n.d.		0,05	N
Triazophos	22	22															n.d.		0,02	N
Vinclozolin	49	49															n.d.		1	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetable:		Food item: Cauliflower (fresh or frozen)	
Reporting country:	<u>The Netherlands</u>	Year of sampling:	<u>1999</u>
Total number of samples analysed:	37	With residues above MRL (EC+national):	0
Without detectable residues:	37	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	0	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50				
Acephate	36	36	0.02															0.00	
Benomyl group(#)	32	32	0.10															0.10	Ec
Chlorpyrifos	34	34	0.05															0.05	Ec
Chlorpyrifos-methyl	34	34	0.05															0.05	Ec
Deltamethrin	34	34	0.05															0.10	Ec
Diazinon	34	34	0.02															0.50	E
Endosulfan	34	34	0.05															1.00	E
Imazalil	34	34	0.02															0.00	
Iprodione	34	34	0.02															0.05	Ec
Lambda-cyhalothrin	34	34	0.05															0.00	
Maneb-group(##)	0	0	0.05															2.00	Ec
Mecarbam	36	36	0.05															0.00	
Metalaxyl	34	34	0.05															0.05	E-O
Methamidophos	36	36	0.01															0.50	Ec
Methodathion	34	34	0.02															0.02	E
Permethrin	34	34	0.05															0.10	Ec
Pirimiphos-methyl	34	34	0.05															1.00	E
Thiabendazol	32	32	0.05															0.05	E
Triazophos	34	34	0.02															0.00	
Vinclozolin	34	34	0.05															0.05	Ec

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(**) in alphabetical order of the English name

(##) Sum of dithiocarbamates, expressed as CS₂

(***) E=EC-MRL, N=National MRL, E-O national MRL in open position, W=without MRL, c=change in 1999

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Peppers	
Reporting country:	<u>The Netherlands</u>	Year of sampling:	<u>1999</u>
Total number of samples analysed:	283	With residues above MRL (EC+national):	109
Without detectable residues:	163	With residues above EC-MRL:	108
With detectable residues at or below MRL or without MRL:	11	With residues above national MRL:	1

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	151	151	0.02																0.02	E-O
Benomyl group(#)	117	115	0.10					2									0.18	1	0.10	Ec
Chlorpyrifos	121	120	0.05					1									0.14		0.50	Ec
Chlorpyrifos-methyl	121	121	0.05																0.50	Ec
Deltamethrin	124	124	0.05																0.20	Ec
Diazinon	121	121	0.02																0.50	E
Endosulfan	124	114	0.05				2	5	3								0.34		1.00	E
Imazalil	121	121	0.02																0.02	Ec
Iprodione	121	116	0.02				1	3									1.6		5.00	Ec
Lambda-cyhalothrin	121	121	0.05																	
Maneb-group(##)	0	0	0.05																2.00	Ec
Mecarbam	151	151	0.05																0.05	E
Metalaxyl	121	119	0.05				1	1									0.13	2	0.05	E
Methamidophos	283	178	0.01		8	9	13	12	33	18	8	4					2.8	105	0.01	Ec
Methidathion	121	121	0.02																0.02	E
Permethrin	121	121	0.05																0.50	Ec
Pirimiphos-methyl	121	118	0.05				2	1									0.17		0.20	E-O
Thiabendazol	117	116	0.05				1										0.07	1	0.05	E-O
Triazophos	121	120	0.02						1								0.24	1	0.02	E
Vinclozolin	121	121	0.05																3.00	Ec

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(**) in alphabetical order of the English name

(##) Sum of dithiocarbamates, expressed as CS₂

(***) E=EC-MRL, N=National MRL, E-O national MRL in open position, W=without MRL, c=change in 1999

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Cauliflower (fresh or frozen)	
Reporting country: <u>Austria</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed: <input type="text" value="12"/>	With residues above MRL (EC+national): <input type="text" value="0"/>	Only add information regarding Cauliflower and the 20 pesticides and do not change or delete rows or columns Maneb-group: Analytical problems with matrix	
Without detectable residues: <input type="text" value="11"/>	With residues above EC-MRL: <input type="text" value="0"/>		
With detectable residues at or below MRL or without MRL: <input type="text" value="1"/>	With residues above national MRL: <input type="text" value="0"/>		

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)		
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50	
Acephate	12	12	0,02;0,05																	2.00	E
Benomyl group(#)	12	12	0.10																	0.50	E
Chlorpyrifos	12	12	.01;0,02;0,05																	0.05	E
Chlorpyrifos-methyl	12	12	.01;0,02;0,05																	0.05	E
Deltamethrin	12	12	0.05																	0.00	E
Diazinon	12	12	.01;0,02;0,05																	0.50	E
Endosulfan	12	12	0.01																	1.00	E
Imazalil	12	12	0,03;0,1																	0.02	E
Iprodione	12	12	0,02;0,04																	0.05	E
Lambda-cyhalothrin	12	12	0.05																	0.05	E
Maneb-group(##)			0,01;0,02;0,05;0,1																	1.00	E
Mecarbam	12	12	.01;0,02;0,05																	0.05	E
Metalaxyl	12	12	0,03;0,05																	0.05	E
Methamidophos	12	11	0,01;0,03				1											0.04		0.50	E
Methodathion	12	12	0,01;0,02																	0.02	E
Permethrin	12	12	0.05																	0.05	E
Pirimiphos-methyl	12	12	.01;0,02;0,05																	2.00	E
Thiabendazol	12	12	0,05;0,1																	0.05	E
Triazophos	12	12	0,01;0,02																	0.20	N
Vinclozolin	12	12	0.01																	0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Peppers	
Reporting country: <u>Austria</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed:	12	With residues above MRL (EC+national):	6
Without detectable residues:	4	With residues above EC-MRL:	6
With detectable residues at or below MRL or without MRL:	2	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	12	11	0,02;0,05						1								0.362	1	0.02	E
Benomyl group(#)	12	10	0.10				1	1									0.14	1	0.10	E
Chlorpyrifos	12	12	,01;0,02;0,05																0.50	E
Chlorpyrifos-methyl	12	12	,01;0,02;0,05																0.50	E
Deltamethrin	12	12	0.05																0.20	E
Diazinon	12	12	,01;0,02;0,05																0.50	E
Endosulfan	12	8	0.01			2	2										0.096		1.00	E
Imazalil	12	12	0,03;0,1																0.02	E
Iprodione	12	12	0,02;0,04																5.00	E
Lambda-cyhalothrin	12	12	0.05																0.02	N
Maneb-group(##)	12	10	1;0,02;0,05;0,1			2											0.04		2.00	E
Mecarbam	12	12	,01;0,02;0,05																0.05	E
Metalaxyl	12	12	0,03;0,05																0.10	N
Methamidophos	12	6	0,01;0,03				2	1	3								0.395	6	0.05	E
Methidathion	12	12	0,01;0,02																0.02	E
Permethrin	12	12	0.05																0.50	E
Pirimiphos-methyl	12	12	,01;0,02;0,05																0.50	N
Thiabendazol	12	12	0,05;0,1																1.50	N
Triazophos	12	12	0,01;0,02																0.02	E
Vinclozolin	12	12	0.01																3.00	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Wheat grains	
Reporting country: <u>Austria</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed:	<u>12</u>	With residues above MRL (EC+national):	<u>0</u>
Without detectable residues:	<u>11</u>	With residues above EC-MRL:	<u>0</u>
With detectable residues at or below MRL or without MRL:	<u>1</u>	With residues above national MRL:	<u>0</u>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50				
Acephate	12	12	0,02;0,05															0.02	E
Benomyl group(#)	12	12	0.10															0.10	E
Chlorpyrifos	12	12	.01;0,02;0,05															0.05	E
Chlorpyrifos-methyl	12	12	.01;0,02;0,05															3.00	E
Deltamethrin	12	12	0.05															1.00	E
Diazinon	12	12	.01;0,02;0,05															0.05	E
Endosulfan	12	12	0.01															0.10	E
Imazalil	12	12	0,03;0,1															0.02	E
Iprodione	12	11	0,02;0,04			1											0.05	0.50	E
Lambda-cyhalothrin	12	12	0.05															0.02	E
Maneb-group(##)	12	12	1;0,02;0,05;0,1															2.00	E
Mecarbam	12	12	.01;0,02;0,05															0.06	E
Metalaxyl	12	12	0,03;0,05															0.05	E
Methamidophos	12	12	0,01;0,03															0.01	E
Methodathion	12	12	0,01;0,02															0.02	E
Permethrin	12	12	0.05															2.00	E
Pirimiphos-methyl	12	12	.01;0,02;0,05															5.00	E
Thiabendazol	12	12	0,05;0,1															0.50	N
Triazophos	12	12	0,01;0,02															0.02	N
Vinclazolin	12	12	0.01															0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Melon, not squash or watermelon	
Reporting country: <u>Austria</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed:	12	With residues above MRL (EC+national):	4
Without detectable residues:	1	With residues above EC-MRL:	4
With detectable residues at or below MRL or without MRL:	7	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	12	10	0,02;0,05					1	1								0.321	2	0.02	E
Benomyl group(#)	12	12	0.10																0.50	E
Chlorpyrifos	12	12	.01;0,02;0,05																0.05	E
Chlorpyrifos-methyl	12	12	.01;0,02;0,05																0.05	E
Deltamethrin	12	12	0.05																0.05	E
Diazinon	12	12	.01;0,02;0,05																0.50	E
Endosulfan	12	4	0.01		1	3	1	2	1								0.285		1.00	E
Imazalil	12	12	0,03;0,1							1									2.00	E
Iprodione	12	11	0,02;0,04							1									0.30	E
Lambda-cyhalothrin	12	12	0.05																0.02	E
Maneb-group(##)	12	11	1;0,02;0,05;0,1		1												0.014		0.50	E
Mecarbam	12	12	.01;0,02;0,05																0.05	E
Metalaxyl	12	12	0,03;0,05																0.10	E
Methamidophos	12	9	0,01;0,03			2	1										0.06	3	0.01	E
Methidathion	12	12	0,01;0,02																0.02	E
Permethrin	12	12	0.05																0.10	E
Pirimiphos-methyl	12	12	.01;0,02;0,05																0.50	E
Thiabendazol	12	9	0,05;0,1							2	1						1.1		1.50	E
Triazophos	12	12	0,01;0,02																0.02	E
Vinclzolin	12	12	0.01																1.00	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Cauliflower (fresh or frozen)	
Reporting country: <u>Portugal</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed:	65	With residues above MRL (EC+national):	0
Without detectable residues:	61	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	4	With residues above national MRL:	0

Only add information regarding Cauliflower and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)		
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50	
Acephate	24	23	0.02			1												0.05	0	1.00	N
Benomyl group(#)	24	24	0.10																0	3.00	N
Chlorpyrifos	34	34	0.02																0	0.50	N
Chlorpyrifos-methyl	34	34	0.02																0	0.05	E
Deltamethrin	34	34	0.05																0	0.10	E
Diazinon	34	34	0.02																0	0.50	E
Endosulfan	34	34	0.05																0	1.00	E
Imazalil	24	24	0.10																0	0.02	E
Iprodione	34	34	0.02																0	0.02	N
Lambda-cyhalothrin	34	34	0.02																0	0.20	N
Maneb-group(##)	31	28	0.05				1	1	1									0.3	0	1.00	N
Mecarbam	24	24	0.05																0	0.05	E
Metalaxyl	34	34	0.10																0	0.20	N
Methamidophos	24	24	0.01																0	0.20	N
Methidathion	34	34	0.02																0	0.02	E
Permethrin	34	34	0.05																0	0.10	E
Pirimiphos-methyl	34	34	0.02																0	1.00	E
Thiabendazol	34	34	0.20																0	0.05	E
Triazophos	24	24	0.02																0	0.02	N
Vinclozolin	34	34	0.02																0	0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Peppers	
Reporting country: <u>Portugal</u>	Year of sampling: <u>1999</u>	Remark: By the end of 1999 directive 98/82/CE was still not transposed into the national legislation.	
Total number of samples analysed:	<u>97</u>	With residues above MRL (EC+national):	<u>13</u>
Without detectable residues:	<u>62</u>	With residues above EC-MRL:	<u>0</u>
With detectable residues at or below MRL or without MRL:	<u>22</u>	With residues above national MRL:	<u>13</u>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)		
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50	
Acephate	82	65	0.02		3	9	3	1	1									0.29	0	0.50	N
Benomyl group(#)	18	18	0.10																0	2.00	N
Chlorpyrifos	82	82	0.02																0	0.50	E
Chlorpyrifos-methyl	82	82	0.02																0	0.50	E
Deltamethrin	18	18	0.05																0	0.20	E
Diazinon	82	82	0.02																0	0.50	E
Endosulfan	18	17	0.05				1											0.08	0	1.00	E
Imazalil	65	65	0.10																0	0.02	N
Iprodione	18	18	0.02																0	5.00	E
Lambda-cyhalothrin	18	18	0.02																0	0.20	N
Maneb-group(##)	31	29	0.05					1		1								0.6	0	2.00	E
Mecarbam	65	65	0.05																0	0.05	E
Metalaxyl	65	65	0.10																0	0.20	N
Methamidophos	82	49	0.01	1	1	13	5	5	6	2								0.93	13	0.10	N
Methidathion	65	65	0.02																0	0.02	E
Permethrin	18	18	0.05																0	0.50	E
Pirimiphos-methyl	82	82	0.02																0	0.50	N
Thiabendazol	65	65	0.20																0	0.05	N
Triazophos	65	65	0.02																0	0.02	E
Vinclozolin	18	18	0.02																0	3.00	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(**) in alphabetical order of the English name

(##) Sum of dithiocarbamates, expressed as CS₂

(***) E=EC-MRL, N=National MRL, W=without MRL

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Wheat grains	
Reporting country:	<u>Portugal</u>	Year of sampling:	<u>1999</u>
Total number of samples analysed:		With residues above MRL (EC+national):	
Without detectable residues:		With residues above EC-MRL:	
With detectable residues at or below MRL or without MRL:		With residues above national MRL:	
	70		0
	58		0
	12		0

Remark: Only add information regarding wheat grains and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	16	16	0.02															0	0.02	E
Benomyl group(#)	16	16	0.10															0	0.10	E
Chlorpyrifos	65	65	0.02															0	0.05	E
Chlorpyrifos-methyl	65	61	0.02			1		2		1							0.68	0	5.00	N
Deltamethrin	41	41	0.05															0	1.00	N
Diazinon	65	65	0.02															0	0.05	E
Endosulfan	41	41	0.05															0	0.10	E
Imazalil	16	16	0.10															0	0.02	E
Iprodione	41	41	0.02															0	0.50	E
Lambda-cyhalothrin	41	41	0.02															0	0.02	E
Maneb-group(##)	24	24	0.05															0	0.05	N
Mecarbam	16	16	0.05															0	0.05	E
Metalaxyl	41	41	0.10															0	0.05	E
Methamidophos	16	16	0.01															0	0.01	E
Methodathion	56	56	0.02															0	0.02	E
Permethrin	41	41	0.05															0	2.00	E
Pirimiphos-methyl	65	55	0.02			1	1	4		4							0.98	0	5.00	E
Thiabendazol	41	41	0.20															0	0.20	N
Triazophos	16	16	0.02															0	0.02	N
Vinclozolin	41	41	0.02															0	0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Melon, not squash or watermelon	
Reporting country:	<u>Portugal</u>	Year of sampling:	<u>1999</u>
		Remark:	Only add information regarding melon and the 20 pesticides and do not change or delete rows or columns
Total number of samples analysed:	68	With residues above MRL (EC+national):	4
Without detectable residues:	51	With residues above EC-MRL:	3
With detectable residues at or below MRL or without MRL:	13	With residues above national MRL:	1

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)		
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50	
Acephate	28	26	0.02						2									0.22	2	0.02	E
Benomyl group(#)	28	27	0.10				1											0.1	0	0.50	E
Chlorpyrifos	39	39	0.02																0	0.05	E
Chlorpyrifos-methyl	39	39	0.02																0	0.05	E
Deltamethrin	39	39	0.05																0	0.05	E
Diazinon	39	39	0.02																0	0.50	E
Endosulfan	39	35	0.05				2	2										0.17	0	1.00	E
Imazalil	28	28	0.10																0	0.02	N
Iprodione	39	39	0.02																0	0.02	N
Lambda-cyhalothrin	39	39	0.02																0	0.05	N
Maneb-group(##)	29	21	0.05				2	1	3	1		1						3.8	1	1.00	N
Mecarbam	28	28	0.05																0	0.05	E
Metalaxyl	39	39	0.10																0	0.05	N
Methamidophos	28	24	0.01			2	2											0.06	3	0.01	E
Methidathion	39	39	0.02																0	0.02	E
Permethrin	39	39	0.05																0	0.10	E
Pirimiphos-methyl	39	39	0.02																0	0.20	N
Thiabendazol	39	39	0.20																0	0.05	N
Triazophos	28	28	0.02																0	0.02	N
Vinclozolin	39	39	0.02																0	1.00	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Cauliflower (fresh or frozen)	
Reporting country:	<u>Finland</u>	Year of sampling:	<u>1999</u>
Remark: Only add information regarding Cauliflower and the 20 pesticides and do not change or delete rows or columns			
Total number of samples analysed:	44	With residues above MRL (EC+national):	0
Without detectable residues:	44	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	0	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	44	44	0.05															0	2	E
Benomyl group(#)	6	6	0.1															0	0.1	E
Chlorpyrifos	44	44	0.01															0	0.05	E
Chlorpyrifos-methyl	44	44	0.01															0	0.05	E
Deltamethrin	44	44	0.2															0	0.1	E
Diazinon	44	44	0.01															0	0.5	E
Endosulfan	44	44	0.01															0	1	E
Imazalil	44	44	0.15															0	0.02	E
Iprodione	44	44	0.02															0	0.05	E
Lambda-cyhalothrin	44	44	0.01															0	0.02	E
Maneb-group(##)	6	6	0.2															0	1	E
Mecarbam	44	44	0.05															0	0.05	E
Metalaxyl	44	44	0.05															0	0.05	E
Methamidophos	44	44	0.02															0	0.5	E
Methidathion	44	44	0.01															0	0.02	E
Permethrin	44	44	0.1															0	0.1	E
Pirimiphos-methyl	44	44	0.01															0	1	E
Thiabendazol	44	44	0.03															0	0.05	E
Triazophos	35	35	0.02															0	0.1	N
Vinclozolin	44	44	0.01															0	0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS2

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Peppers	
Reporting country:	<u>Finland</u>	Year of sampling:	<u>1999</u>
Remark: Only add information regarding peppers and the 20 pesticides and do not change or delete rows or columns			
Total number of samples analysed:	279	With residues above MRL (EC+national):	80
Without detectable residues:	52	With residues above EC-MRL:	80
With detectable residues at or below MRL or without MRL:	147	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	279	275	0.05			1	1	1	1									0.31	4	0.02	E
Benomyl group(#)	260	260	0.1																0	0.1	E
Chlorpyrifos	279	268	0.01	2	3	2	3	1										0.11	0	0.5	E
Chlorpyrifos-methyl	279	260	0.01	3	3	5	5	2		1								0.53	1	0.5	E
Deltamethrin	279	279	0.2																0	0.2	E
Diazinon	279	279	0.01																0	0.5	E
Endosulfan	279	75	0.01	17	20	28	26	38	57	15	3							1.5	3	1	E
Imazalil	279	279	0.15																0	0.02	E
Iprodione	279	267	0.02		2	6		2	2									0.42	0	5	E
Lambda-cyhalothrin	279	279	0.01																		W
Maneb-group(##)	260	260	0.2																0	2	E
Mecarbam	279	279	0.05																0	0.05	E
Metalaxyl	279	279	0.05																0	0.05	E
Methamidophos	279	209	0.02			7	15	16	18	14								0.97	70	0.01	E
Methidathion	279	277	0.01			1	1											0.06	2	0.02	E
Permethrin	279	273	0.1				1	3	2									0.47	0	0.5	E
Pirimiphos-methyl	279	205	0.01	7	6	15	25	13	7	1								0.93	0	1	N
Thiabendazol	279	279	0.03																0	0.05	E
Triazophos	265	265	0.02																0	0.02	E
Vinclozolin	279	274	0.01			2	2	1										0.11	0	3	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS2

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Wheat grains	
Reporting country:	<u>Finland</u>	Year of sampling:	<u>1999</u>
Remark: Only add information regarding wheat grains and the 20 pesticides and do not change or delete rows or columns			
Total number of samples analysed:	<input type="text" value="13"/>	With residues above MRL (EC+national):	<input type="text" value="0"/>
Without detectable residues:	<input type="text" value="13"/>	With residues above EC-MRL:	<input type="text" value="0"/>
With detectable residues at or below MRL or without MRL:	<input type="text" value="0"/>	With residues above national MRL:	<input type="text" value="0"/>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	13	13	0.05															0	0.02	E
Benomyl group(#)	13	13	0.1															0	0.1	E
Chlorpyrifos	13	13	0.01															0	0.05	E
Chlorpyrifos-methyl	13	13	0.01															0	3	E
Deltamethrin	13	13	0.2															0	1	E
Diazinon	13	13	0.01															0	0.05	E
Endosulfan	13	13	0.01															0	0.1	E
Imazalil	13	13	0.15															0	0.02	E
Iprodione	13	13	0.02															0	0.5	E
Lambda-cyhalothrin	13	13	0.01															0	0.02	E
Maneb-group(##)	13	13	0.2															0	1	E
Mecarbam	13	13	0.05															0	0.05	E
Metalaxyl	13	13	0.05															0	0.05	E
Methamidophos	13	13	0.02															0	0.01	E
Methodathion	13	13	0.01															0	0.02	E
Permethrin	13	13	0.1															0	2	E
Pirimiphos-methyl	13	13	0.01															0	5	E
Thiabendazol	13	13	0.03															0	0.05	E
Triazophos	13	13	0.02															0	0.02	E
Vinclozolin	13	13	0.01															0	0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS2

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Melon, not squash or watermelon	
Reporting country:	<u>Finland</u>	Year of sampling:	<u>1999</u>
Total number of samples analysed: <input type="text" value="9"/>		With residues above MRL (EC+national): <input type="text" value="1"/>	
Without detectable residues: <input type="text" value="4"/>		With residues above EC-MRL: <input type="text" value="1"/>	
With detectable residues at or below MRL or without MRL: <input type="text" value="4"/>		With residues above national MRL: <input type="text" value="0"/>	
Remark: Only add information regarding melon and the 20 pesticides and do not change or delete rows or columns			

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50				
Acephate	9	9	0.05															0	0.02	E
Benomyl group(#)	9	9	0.1															0	0.5	E
Chlorpyrifos	9	9	0.01															0	0.05	E
Chlorpyrifos-methyl	9	9	0.01															0	0.05	E
Deltamethrin	9	9	0.2															0	0.05	E
Diazinon	9	9	0.01															0	0.5	E
Endosulfan	9	4	0.01	1			2	2									0.13	0	1	E
Imazalil	9	9	0.15															0	2	E
Iprodione	9	9	0.02															0	0.3	E
Lambda-cyhalothrin	9	9	0.01																	W
Maneb-group(##)	9	9	0.2															0	0.5	E
Mecarbam	9	9	0.05															0	0.05	E
Metalaxyl	9	9	0.05															0	0.05	E
Methamidophos	9	9	0.02															0	0.01	E
Methodathion	9	9	0.01															0	0.02	E
Permethrin	9	9	0.1															0	0.1	E
Pirimiphos-methyl	9	9	0.01																	W
Thiabendazol	9	8	0.03				1										0.08	1	0.05	E
Triazophos	9	9	0.02																	W
Vinclozolin	9	9	0.01															0	1	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS2

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Cauliflower (fresh or frozen)	
Reporting country: <u>Sweden</u>	Year of sampling: <u>1999</u>	Remark: Only add information regarding Cauliflower and the 20 pesticides and do not change or delete rows or columns	
Total number of samples analysed:	<input type="text" value="48"/>	With residues above MRL (EC+national):	<input type="text" value="0"/>
Without detectable residues:	<input type="text" value="47"/>	With residues above EC-MRL:	<input type="text" value="0"/>
With detectable residues at or below MRL or without MRL:	<input type="text" value="1"/>	With residues above national MRL:	<input type="text" value="0"/>

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)		
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50	
Acephate	48	47	0.02								1							0.72		2.00	E
Benomyl group(#)	31	31	0.10																	0.10	E
Chlorpyrifos	48	48	0.05																	0.5/0.05	N/E
Chlorpyrifos-methyl	48	48	0.05																	0.05	E
Deltamethrin	48	48	0.05																	0.10	E
Diazinon	48	48	0.02																	0.50	E
Endosulfan	48	48	0.01																	1.00	E
Imazalil	48	48	0.20																	0.02	E
Iprodione	48	48	0.20																	10/0.05	N/E
Lambda-cyhalothrin	48	48	0.10																	No MRL	
Maneb-group(##)	12	12	0.10																	1.00	E
Mecarbam	48	48	0.05																	0.05	E
Metalaxyl	48	48	0.10																	No MRL	
Methamidophos	48	47	0.02					1										0.15		0.50	E
Methidathion	48	48	0.02																	0.02	E
Permethrin	48	48	0.05																	0.10	E
Pirimiphos-methyl	48	48	0.05																	1.00	E
Thiabendazol	48	48	0.3/0.2																	0.05	E
Triazophos	48	48	0.02																	0.10	N
Vinclozolin	48	48	0.05																	0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Peppers	
Reporting country: <u>Sweden</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed:	126	With residues above MRL (EC+national):	30
Without detectable residues:	62	With residues above EC-MRL:	30
With detectable residues at or below MRL or without MRL:	34	With residues above national MRL:	0

Only add information regarding peppers and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)											Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)			
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20					50	>50	
Acephate	126	123	0.02		1		2											0.098	1	1/0.02	N/E
Benomyl group(#)	52	52	0.10																	0.10	E
Chlorpyrifos	126	125	0.05				1											0.066		0.50	E
Chlorpyrifos-methyl	126	123	0.05					2	1									0.3		0.50	E
Deltamethrin	126	126	0.05																	0.20	E
Diazinon	126	126	0.02																	0.50	E
Endosulfan	126	75	0.01		3	1	17	8	16	6								0.834		1.00	E
Imazalil	126	126	0.20																	0.02	E
Iprodione	126	126	0.20																	5.00	E
Lambda-cyhalothrin	126	126	0.10																	No MRL	
Maneb-group(##)	31	30	0.10					1										0.12		2.00	E
Mecarbam	126	126	0.05																	0.05	E
Metalaxyl	126	126	0.10																	No MRL	
Methamidophos	126	95	0.02		1	4	4	5	11	5	1							1.6	27	1/0.01	N/E
Methidathion	126	122	0.02		1	2	1											0.062	3	0.02	E
Permethrin	126	126	0.05																	0.50	E
Pirimiphos-methyl	126	123	0.05				2	1										0.11		1.00	N
Thiabendazol	126	126	0.3/0.2																	6.00	N
Triazophos	126	126	0.02																	0.02	E
Vinclozolin	126	123	0.05				2		1									0.31		3.00	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Wheat grains	
Reporting country: <u>Sweden</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed:	140	With residues above MRL (EC+national):	0
Without detectable residues:	128	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	12	With residues above national MRL:	0

Only add information regarding wheat grains and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate																			0.02	E
Benomyl group(##)	15	15	0.10																0.10	E
Chlorpyrifos	140	140	0.01																0.05	E
Chlorpyrifos-methyl	140	131	0.01		3	3	1	1		1							0.73		3.00	E
Deltamethrin	140	140	0.20																1.00	E
Diazinon	140	140	0.02																0.05	E
Endosulfan	140	140	0.02																0.10	E
Imazalil	140	140	0.30																0.02	E
Iprodione	140	140	0.40																0.50	E
Lambda-cyhalothrin																			0.02	E
Maneb-group(##)																			1.00	E
Mecarbam																			0.05	E
Metalaxyl	140	140	0.40																0.05	E
Methamidophos																			0.01	E
Methodathion																			0.02	E
Permethrin	140	140	0.90																2.00	E
Pirimiphos-methyl	140	136	0.05				2	1		1							1.15		5.00	E
Thiabendazol	140	140	0.20																No MRL	E
Triazophos																			No MRL	E
Vinclozolin	140	140	0.06																0.05	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Melon, not squash or watermelon	
Reporting country: <u>Sweden</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed:	69	With residues above MRL (EC+national):	6
Without detectable residues:	33	With residues above EC-MRL:	6
With detectable residues at or below MRL or without MRL:	30	With residues above national MRL:	0

Only add information regarding melon and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)		
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50	
Acephate	69	68	0.02				1											0.07	1	0.02	E
Benomyl group(#)	68	68	0.10																	0.50	E
Chlorpyrifos	69	68	0.05					1										0.108	1	0.05	E
Chlorpyrifos-methyl	69	69	0.05																	0.05	E
Deltamethrin	69	69	0.05																	0.05	E
Diazinon	69	67	0.02			1	1											0.054		0.50	E
Endosulfan	69	39	0.01		1	10	15	2	2									0.218		1.00	E
Imazalil	69	69	0.20																	2.00	E
Iprodione	69	69	0.20																	10/0.3	N/E
Lambda-cyhalothrin	69	69	0.10																	No MRL	
Maneb-group(##)	17	16	0.10				1											0.1		1/0.5	N/E
Mecarbam	69	69	0.05																	0.05	E
Metalaxyl	69	69	0.10																	No MRL	
Methamidophos	69	65	0.02			2	2											0.096	4	0.01	E
Methidathion	69	69	0.02																	0.02	E
Permethrin	69	69	0.05																	0.10	E
Pirimiphos-methyl	69	69	0.05																	No MRL	
Thiabendazol	69	68	0.3/0.2						1									0.26		6.00	N
Triazophos	69	69	0.02																	No MRL	
Vinclozolin	69	69	0.05																	1.00	E

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Cauliflower (fresh or frozen)	
Reporting country: <u>UK</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed:	71	With residues above MRL (EC-national):	0
Without detectable residues:	71	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	0	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	71	71	0.02															0		
Benomyl group(#)	71	71	0.10															0		
Chlorpyrifos	71	71	0.05															0		
Chlorpyrifos-methyl	71	71	0.05															0		
Deltamethrin	71	71	0.05															0		
Diazinon	71	71	0.02															0		
Endosulfan	71	71	0.05															0		
Imazalil	71	71	0.02															0		
Iprodione	71	71	0.02															0		
Lambda-cyhalothrin	71	71	0.02															0		
Maneb-group(##)****	0	0																0		
Mecarbam	71	71	0.05															0		
Metalaxyl	71	71	0.05															0		
Methamidophos	71	71	0.03															0		
Methodathion	71	71	0.02															0		
Permethrin	71	71	0.05															0		
Pirimiphos-methyl	71	71	0.05															0		
Thiabendazol	71	71	0.05															0		
Triazophos	71	71	0.02															0		
Vinclozolin	71	71	0.05															0		

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

**** analytical difficulties with maneb group

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Cereals		Food item: Wheat grains	
Reporting country: <u>UK</u>		Year of sampling: <u>1999</u>	Remark:
Total number of samples analysed:	62	With residues above MRL (EC+national):	0
Without detectable residues:	51	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	11	With residues above national MRL:	0

Only add information regarding wheat grains and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	62	62	0.02															0		
Benomyl group(#)	62	62	0.10															0		
Chlorpyrifos	62	62	0.05															0		
Chlorpyrifos-methyl	62	62	0.05															0		
Deltamethrin	62	58	0.05			2	2										0.06	0	1.00	E
Diazinon	62	62	0.05															0		
Endosulfan	62	62	0.05															0		
Imazalil	62	62	0.02															0		
Iprodione	62	62	0.02															0		
Lambda-cyhalothrin	62	62	0.02															0		
Maneb-group(##)	62	62	0.05															0		
Mecarbam	62	62	0.05															0		
Metalaxyl	62	62	0.05															0		
Methamidophos	62	62	0.01															0		
Methidathion	62	62	0.02															0		
Permethrin	62	62	0.05															0		
Pirimiphos-methyl	62	54	0.05			1	7										0.1	0	5.00	E
Thiabendazol	62	62	0.05															0		
Triazophos	62	62	0.02															0		
Vinclzolin	62	62	0.05															0		

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Melon, not squash or watermelon	
Reporting country: <u>UK</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed:	72	With residues above MRL (EC+national):	6
Without detectable residues:	42	With residues above EC-MRL:	6
With detectable residues at or below MRL or without MRL:	24	With residues above national MRL:	0

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)		
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50	
Acephate	72	72	0.02															0.2	0	0.5	E
Benomyl group(#)	72	70	0.50					2													
Chlorpyrifos	72	72	0.05																		
Chlorpyrifos-methyl	72	72	0.05																		
Deltamethrin	72	72	0.05																		
Diazinon	72	70	0.02			1	1											0.07	0	0.5	E
Endosulfan	72	63	0.05				3	5	1									0.4	0	1	E
Imazalil	72	72	0.10																		
Iprodione	72	72	0.05																		
Lambda-cyhalothrin	72	72	0.02																		
Maneb-group(##)	72	67	0.05				4			1								0.6	0	-	W
Mecarbam	72	72	0.05																		
Metalaxyl	72	72	0.05																		
Methamidophos	72	64	0.01	2		2	2	1	1									0.3	6	0.01	E
Methodathion	72	72	0.02																		
Permethrin	72	71	0.05				1											0.06	0	0.1	E
Pirimiphos-methyl	72	72	0.05																		
Thiabendazol	72	65	0.05				3		1	3								0.8	0	-	W
Triazophos	72	72	0.02																		
Vinclzolin	72	72	0.05																		

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Peppers	
Reporting country: <u>Iceland</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed:	13	With residues above MRL (EC+national):	1
Without detectable residues:	11	With residues above EC-MRL:	1
With detectable residues at or below MRL or without MRL:	1	With residues above national MRL:	0

Only add information regarding peppers and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50				
Acephate	0	0																	
Benomyl group(#)	0	0																	
Chlorpyrifos	13	13	0.05															0.50	N
Chlorpyrifos-methyl	13	13	0.05															0.50	N
Deltamethrin	0	0																	
Diazinon	13	13	0.05															0.50	N
Endosulfan	13	13	0.05															1.00	N
Imazalil	13	13	0.05															0.02	N
Iprodione	13	13	0.05															5.00	N
Lambda-cyhalothrin	0	0																	
Maneb-group(##)	0	0																	
Mecarbam	0	0																	
Metalaxyl	13	13	0.05															0.05	N
Methamidophos	13	12	0.05					1								0.18	1	0.01	E
Methidathion	13	13	0.05															0.02	N
Permethrin	13	13	0.05															0.50	N
Pirimiphos-methyl	0	0																	
Thiabendazol	13	13	0.05															0.05	N
Triazophos	0	0																	
Vinclozolin	13	13	0.05															3.00	N

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Fruiting vegetables		Food item: Melon, not squash or watermelon	
Reporting country: <u>Iceland</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed:	13	With residues above MRL (EC+national):	0
Without detectable residues:	11	With residues above EC-MRL:	0
With detectable residues at or below MRL or without MRL:	2	With residues above national MRL:	0

Only add information regarding melon and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)													Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50	>50					
Acephate	0	0																			
Benomyl group(#)	0	0																			
Chlorpyrifos	13	12	0.05			1											0.05	0	0.05	N	
Chlorpyrifos-methyl	13	13	0.05																0.05	N	
Deltamethrin																					
Diazinon	13	13	0.05																0.50	N	
Endosulfan	13	13	0.05																1.00	N	
Imazalil	13	13	0.05																2.00	N	
Iprodione	13	13	0.05																0.30	N	
Lambda-cyhalothrin																					
Maneb-group(##)																					
Mecarbam																					
Metalaxyl	13	13	0.05																0.05	N	
Methamidophos	13	13	0.05																0.01	N	
Methidathion	13	13	0.05																0.02	N	
Permethrin	13	13	0.05																0.10	N	
Pirimiphos-methyl																					
Thiabendazol	13	12	0.05					1									0.11	0	0.05	N	
Triazophos																					
Vinclozolin	13	13	0.05																1.00	N	

(*) i.e column 0.02 includes the range from 0.011... mg/kg upto 0.020... mg/kg

(**) in alphabetical order of the English name

(***) E=EC-MRL, N=National MRL, W=without MRL

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

Table B: Notifications of the co-ordinated programme (specific exercise) to the European Commission

Product group: Brassica vegetables		Food item: Cauliflower (fresh or frozen)	
Reporting country: <u>Norway</u>	Year of sampling: <u>1999</u>	Remark:	
Total number of samples analysed:	<u>79</u>	With residues above MRL (EC+national):	<u>0</u>
Without detectable residues:	<u>78</u>	With residues above EC-MRL:	<u>0</u>
With detectable residues at or below MRL or without MRL:	<u>1</u>	With residues above national MRL:	<u>0</u>

Only add information regarding Cauliflower and the 20 pesticides and do not change or delete rows or columns

Pesticide (**)	Total number of samples	Number of samples without residues	Reporting level (mg/kg)	Samples with quantifiable residues in classes up to and including (in mg/kg) (*)												Maximum residue level found (mg/kg)	Number of samples with residues exceeding the MRL	MRL (mg/kg)	Source of MRL (***)	
				0.01	0.02	0.05	0.1	0.2	0.5	1	2	5	10	20	50					>50
Acephate	79	79	0.05																	
Benomyl group(#)	27	26	0.05					1										0.12		W
Chlorpyrifos	79	79	0.05																	
Chlorpyrifos-methyl	79	79	0.05																	
Deltamethrin	79	79	0.05																	
Diazinon	79	79	0.02																	
Endosulfan	79	79	0.02																	
Imazalil	79	79	0.02																	
Iprodione	79	79	0.02																	
Lambda-cyhalothrin	79	79	0.02																	
Maneb-group(##)	12	12	0.05																	
Mecarbam	79	79	0.05																	
Metalaxyl	79	79	0.05																	
Methamidophos	79	79	0.05																	
Methidathion	79	79	0.05																	
Permethrin	79	79	0.05																	
Pirimiphos-methyl	79	79	0.05																	
Thiabendazol	79	79	0.05																	
Triazophos	79	79	0.05																	
Vinclozolin	79	79	0.02																	

(#) Benomyl, carbendazim, thiophanate-methyl (sum of residues expressed as carbendazim).

(##) Sum of dithiocarbamates, expressed as CS₂

(***) E=EC-MRL, N=National MRL, W=without MRL

