

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

## 1. NAME OF THE MEDICINAL PRODUCT

Jardiance 10 mg film-coated tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 10 mg empagliflozin.

Excipient with known effect:

Each tablet contains lactose monohydrate equivalent to 154.3 mg lactose anhydrous.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Film-coated tablet (tablet).

Round, pale yellow, biconvex, bevel-edged film-coated tablet debossed with “S10” on one side and the Boehringer Ingelheim logo on the other (tablet diameter: 9.1 mm).

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Jardiance is indicated in the treatment of type 2 diabetes mellitus to improve glycaemic control in adults as:

#### Monotherapy

When diet and exercise alone do not provide adequate glycaemic control in patients for whom use of metformin is considered inappropriate due to intolerance.

#### Add-on combination therapy

In combination with other glucose-lowering medicinal products including insulin, when these, together with diet and exercise, do not provide adequate glycaemic control (see sections 4.4, 4.5 and 5.1 for available data on different combinations).

### 4.2 Posology and method of administration

#### Posology

##### Monotherapy and add-on combination

The recommended starting dose is 10 mg empagliflozin once daily for monotherapy and add-on combination therapy with other glucose-lowering medicinal products including insulin. In patients tolerating empagliflozin 10 mg once daily who have an eGFR  $\geq 60$  ml/min/1.73 m<sup>2</sup> and need tighter glycaemic control, the dose can be increased to 25 mg once daily. The maximum daily dose is 25 mg (see below and section 4.4).

When empagliflozin is used in combination with a sulphonylurea or with insulin, a lower dose of the sulphonylurea or insulin may be considered to reduce the risk of hypoglycaemia (see sections 4.5 and 4.8).

### Special populations

#### *Patients with renal impairment*

Due to the mechanism of action, the efficacy of empagliflozin is dependent on renal function. No dose adjustment is required for patients with an eGFR  $\geq 60$  ml/min/1.73 m<sup>2</sup> or CrCl  $\geq 60$  ml/min.

Empagliflozin should not be initiated in patients with an eGFR  $<60$  ml/min/1.73 m<sup>2</sup> or CrCl  $<60$  ml/min. In patients tolerating empagliflozin whose eGFR falls persistently below 60 ml/min/1.73 m<sup>2</sup> or CrCl below 60 ml/min, the dose of empagliflozin should be adjusted to or maintained at 10 mg once daily. Empagliflozin should be discontinued when eGFR is persistently below 45 ml/min/1.73 m<sup>2</sup> or CrCl persistently below 45 ml/min (see sections 4.4, 4.8, 5.1, and 5.2).

Empagliflozin should not be used in patients with end stage renal disease (ESRD) or in patients on dialysis as it is not expected to be effective in these patients (see sections 4.4 and 5.2).

#### *Patients with hepatic impairment*

No dose adjustment is required for patients with hepatic impairment. Empagliflozin exposure is increased in patients with severe hepatic impairment. Therapeutic experience in patients with severe hepatic impairment is limited and therefore not recommended for use in this population (see section 5.2).

#### *Elderly patients*

No dose adjustment is recommended based on age. In patients 75 years and older, an increased risk for volume depletion should be taken into account (see sections 4.4 and 4.8). In patients aged 85 years and older, initiation of empagliflozin therapy is not recommended due to the limited therapeutic experience (see section 4.4).

#### *Paediatric population*

The safety and efficacy of empagliflozin in children and adolescents has not yet been established. No data are available.

### Method of administration

The tablets can be taken with or without food, swallowed whole with water. If a dose is missed, it should be taken as soon as the patient remembers. A double dose should not be taken on the same day.

## **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

## **4.4 Special warnings and precautions for use**

### General

Jardiance should not be used in patients with type 1 diabetes or for the treatment of diabetic ketoacidosis.

### Use in patients with renal impairment

Jardiance should not be initiated in patients with an eGFR below 60 ml/min/1.73 m<sup>2</sup> or CrCl  $<60$  ml/min. In patients tolerating empagliflozin whose eGFR is persistently below 60 ml/min/1.73 m<sup>2</sup> or CrCl  $<60$  ml/min, the dose of empagliflozin should be adjusted to or maintained at 10 mg once daily. Empagliflozin should be discontinued when eGFR is persistently below 45 ml/min/1.73 m<sup>2</sup> or CrCl persistently below 45 ml/min. Empagliflozin should not be used in patients

with ESRD or in patients on dialysis as it is not expected to be effective in these patients (see section 4.2 and 5.2).

#### Monitoring of renal function

Due to the mechanism of action, the efficacy of empagliflozin is dependent on renal function.

Therefore assessment of renal function is recommended as follows:

- Prior to empagliflozin initiation and periodically during treatment, i.e. at least yearly (see sections 4.2, 5.1 and 5.2).
- Prior to initiation of any concomitant medicinal product that may have a negative impact on renal function.

#### Hepatic injury

Cases of hepatic injury have been reported with empagliflozin in clinical trials. A causal relationship between empagliflozin and hepatic injury has not been established.

#### Elderly patients

The effect of empagliflozin on urinary glucose excretion is associated with osmotic diuresis, which could affect the hydration status. Patients aged 75 years and older may be at an increased risk of volume depletion. A higher number of these patients treated with empagliflozin had adverse reactions related to volume depletion as compared to placebo (see section 4.8).

Therapeutic experience in patients aged 85 years and older is limited. Initiation of empagliflozin therapy in this population is not recommended (see section 4.2).

#### Use in patients at risk for volume depletion

Based on the mode of action of SGLT-2 inhibitors, osmotic diuresis accompanying therapeutic glucosuria may lead to a modest decrease in blood pressure (see section 5.1). Therefore, caution should be exercised in patients for whom an empagliflozin-induced drop in blood pressure could pose a risk, such as patients with known cardiovascular disease, patients on anti-hypertensive therapy with a history of hypotension or patients aged 75 years and older.

In case of conditions that may lead to fluid loss (e.g. gastrointestinal illness), careful monitoring of volume status (e.g. physical examination, blood pressure measurements, laboratory tests including haematocrit) and electrolytes is recommended for patients receiving empagliflozin. Temporary interruption of treatment with empagliflozin should be considered until the fluid loss is corrected.

#### Urinary tract infections

The overall frequency of urinary tract infection reported as adverse event was similar in patients treated with empagliflozin 25 mg and placebo and higher in patients treated with empagliflozin 10 mg (see section 4.8). Complicated urinary tract infection (e.g. pyelonephritis or urosepsis) occurred at a similar frequency in patients treated with empagliflozin compared to placebo. However, temporary interruption of empagliflozin should be considered in patients with complicated urinary tract infections.

#### Cardiac failure

Experience in New York Heart Association (NYHA) class I-II is limited, and there is no experience in clinical studies with empagliflozin in NYHA class III-IV.

#### Urine laboratory assessments

Due to its mechanism of action, patients taking Jardiance will test positive for glucose in their urine.

## Lactose

The tablets contain lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency, or glucose-galactose malabsorption should not take this medicinal product.

## **4.5 Interaction with other medicinal products and other forms of interaction**

### Pharmacodynamic interactions

#### Diuretics

Empagliflozin may add to the diuretic effect of thiazide and loop diuretics and may increase the risk of dehydration and hypotension (see section 4.4).

#### Insulin and insulin secretagogues

Insulin and insulin secretagogues, such as sulphonylureas, may increase the risk of hypoglycaemia. Therefore, a lower dose of insulin or an insulin secretagogue may be required to reduce the risk of hypoglycaemia when used in combination with empagliflozin (see sections 4.2 and 4.8).

### Pharmacokinetic interactions

#### Effects of other medicinal products on empagliflozin

*In vitro* data suggest that the primary route of metabolism of empagliflozin in humans is glucuronidation by uridine 5'-diphosphoglucuronosyltransferases UGT1A3, UGT1A8, UGT1A9, and UGT2B7. Empagliflozin is a substrate of the human uptake transporters OAT3, OATP1B1, and OATP1B3, but not OAT1 and OCT2. Empagliflozin is a substrate of P-glycoprotein (P-gp) and breast cancer resistance protein (BCRP).

Co-administration of empagliflozin with probenecid, an inhibitor of UGT enzymes and OAT3, resulted in a 26% increase in peak empagliflozin plasma concentrations ( $C_{max}$ ) and a 53% increase in area under the concentration-time curve (AUC). These changes were not considered to be clinically meaningful.

The effect of UGT induction on empagliflozin has not been studied. Co-medication with known inducers of UGT enzymes should be avoided due to a potential risk of decreased efficacy.

An interaction study with gemfibrozil, an *in vitro* inhibitor of OAT3 and OATP1B1/1B3 transporters, showed that empagliflozin  $C_{max}$  increased by 15% and AUC increased by 59% following co-administration. These changes were not considered to be clinically meaningful.

Inhibition of OATP1B1/1B3 transporters by coadministration with rifampicin resulted in a 75% increase in  $C_{max}$  and a 35% increase in AUC of empagliflozin. These changes were not considered to be clinically meaningful.

Empagliflozin exposure was similar with and without co-administration with verapamil, a P-gp inhibitor, indicating that inhibition of P-gp does not have any clinically relevant effect on empagliflozin.

Interaction studies conducted in healthy volunteers suggest that the pharmacokinetics of empagliflozin were not influenced by coadministration with metformin, glimepiride, pioglitazone, sitagliptin, linagliptin, warfarin, verapamil, ramipril, simvastatin, torasemide and hydrochlorothiazide.

#### Effects of empagliflozin on other medicinal products

Based on *in vitro* studies, empagliflozin does not inhibit, inactivate, or induce CYP450 isoforms. Empagliflozin does not inhibit UGT1A1. Drug-drug interactions involving the major CYP450 isoforms or UGT1A1 with empagliflozin and concomitantly administered substrates of these enzymes are therefore considered unlikely. The potential for empagliflozin to inhibit UGT2B7 has not been studied.

Empagliflozin does not inhibit P-gp at therapeutic doses. Based on *in vitro* studies, empagliflozin is considered unlikely to cause interactions with drugs that are P-gp substrates. Co-administration of digoxin, a P-gp substrate, with empagliflozin resulted in a 6% increase in AUC and 14% increase in  $C_{max}$  of digoxin. These changes were not considered to be clinically meaningful.

Empagliflozin does not inhibit human uptake transporters such as OAT3, OATP1B1, and OATP1B3 *in vitro* at clinically relevant plasma concentrations and, as such, drug-drug interactions with substrates of these uptake transporters are considered unlikely.

Interaction studies conducted in healthy volunteers suggest that empagliflozin had no clinically relevant effect on the pharmacokinetics of metformin, glimepiride, pioglitazone, sitagliptin, linagliptin, simvastatin, warfarin, ramipiril, digoxin, diuretics and oral contraceptives.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy

There are no data from the use of empagliflozin in pregnant women. Animal studies show that empagliflozin crosses the placenta during late gestation to a very limited extent but do not indicate direct or indirect harmful effects with respect to early embryonic development. However, animal studies have shown adverse effects on postnatal development (see section 5.3). As a precautionary measure, it is preferable to avoid the use of Jardiance during early pregnancy. Jardiance is not recommended during the second and third trimester of pregnancy.

##### Breast-feeding

No data in humans are available on excretion of empagliflozin into milk. Available toxicological data in animals have shown excretion of empagliflozin in milk. A risk to the newborns/infants cannot be excluded. Jardiance should not be used during breast-feeding.

##### Fertility

No studies on the effect on human fertility have been conducted for Jardiance. Animal studies do not indicate direct or indirect harmful effects with respect to fertility (see section 5.3).

#### **4.7 Effects on ability to drive and use machines**

Jardiance has minor influence on the ability to drive and use machines. Patients should be advised to take precautions to avoid hypoglycaemia while driving and using machines, in particular when Jardiance is used in combination with a sulphonylurea and/or insulin.

#### **4.8 Undesirable effects**

##### Summary of the safety profile

A total of 13,076 patients with type 2 diabetes were included in clinical studies to evaluate the safety of empagliflozin. 2856 patients received empagliflozin 10 mg and 3,738 patients received empagliflozin 25 mg for at least 24 weeks and 601 or 881 patients for at least 76 weeks, either alone or in combination with metformin, a sulphonylurea, pioglitazone, DPP-4 inhibitors, or insulin.

In 5 placebo-controlled trials of 18 to 24 weeks duration, 2,971 patients were included of which 995 were treated with placebo and 1,976 with empagliflozin. The overall incidence of adverse events in patients treated with empagliflozin was similar to placebo. The most frequently reported adverse reaction was hypoglycaemia when used with sulphonylurea or insulin (see description of selected adverse reactions).

## Tabulated list of adverse reactions

Adverse reactions classified by system organ class and MedDRA preferred terms reported in patients who received empagliflozin in placebo-controlled studies are presented in the table below (Table 1).

The adverse reactions are listed by absolute frequency. Frequencies are defined as very common ( 1/10), common ( 1/100 to <1/10), uncommon ( 1/1,000 to <1/100), rare ( 1/10,000 to <1/1,000), or very rare (<1/10,000), and not known (cannot be estimated from the available data).

Table 1: Adverse reactions reported in placebo-controlled studies

<b>System organ class</b>	<b>Very common</b>	<b>Common</b>	<b>Uncommon</b>
<i>Infections and infestations</i>		Vaginal moniliasis, vulvovaginitis, balanitis and other genital infection <sup>a</sup> Urinary tract infection <sup>a</sup>	
<i>Metabolism and nutrition disorders</i>	Hypoglycaemia (when used with sulphonylurea or insulin) <sup>a</sup>		
<i>Skin and subcutaneous disorders</i>		Pruritus (generalised)	
<i>Vascular disorders</i>			Volume depletion <sup>a</sup>
<i>Renal and urinary disorders</i>		Increased urination <sup>a</sup>	Dysuria

<sup>a</sup>see subsections below for additional information

## Description of selected adverse reactions

### Hypoglycaemia

The frequency of hypoglycaemia depended on the background therapy in the respective studies.

#### *Minor hypoglycaemia*

The frequency of patients with minor hypoglycaemia was similar for empagliflozin and placebo as monotherapy, add-on to metformin, and add-on to pioglitazone with or without metformin. An increased frequency was noted when given as add-on to metformin and a sulphonylurea (empagliflozin 10 mg: 16.1%, empagliflozin 25 mg: 11.5%, placebo: 8.4%), and as add-on to insulin with or without metformin and with or without a sulphonylurea (empagliflozin 10 mg: 19.5%, empagliflozin 25 mg: 27.1%, placebo: 20.6% during initial 18 weeks treatment when insulin could not be adjusted; empagliflozin 10 mg: 36.1%, empagliflozin 25 mg: 34.8%, placebo 35.3% over the 78-week trial).

#### *Major hypoglycaemia (hypoglycaemia requiring assistance)*

No increase in major hypoglycaemia was observed with empagliflozin compared to placebo as monotherapy, add-on to metformin, add-on to metformin and a sulphonylurea, and add-on to pioglitazone with or without metformin. An increased frequency was noted when given as add-on to insulin with or without metformin and with or without a sulphonylurea (empagliflozin 10 mg: 0%, empagliflozin 25 mg: 1.3%, placebo: 0% during initial 18 weeks treatment when insulin could not be adjusted; empagliflozin 10 mg: 0%, empagliflozin 25 mg: 1.3%, placebo 0% over the 78-week trial).

### Vaginal moniliasis, vulvovaginitis, balanitis and other genital infection

Vaginal moniliasis, vulvovaginitis, balanitis and other genital infections were reported more frequently in patients treated with empagliflozin (empagliflozin 10 mg: 4.1%, empagliflozin 25 mg: 3.7%) compared to placebo (0.9%). These infections were reported more frequently in females treated with empagliflozin compared to placebo, and the difference in frequency was less pronounced in males. The genital tract infections were mild or moderate in intensity.

### Increased urination

Increased urination (including the predefined terms pollakiuria, polyuria, and nocturia) was observed at higher frequencies in patients treated with empagliflozin (empagliflozin 10 mg: 3.4%, empagliflozin 25 mg: 3.2%) compared to placebo (1.0%). Increased urination was mostly mild or moderate in intensity. The frequency of reported nocturia was similar for placebo and empagliflozin (<1%).

### Urinary tract infection

The overall frequency of urinary tract infection reported as adverse event was similar in patients treated with empagliflozin 25 mg and placebo (7.6%) and higher in empagliflozin 10 mg (9.3%). Similar to placebo, urinary tract infection was reported more frequently for empagliflozin in patients with a history of chronic or recurrent urinary tract infections. The intensity (mild, moderate, severe) of urinary tract infection was similar in patients treated with empagliflozin and placebo. Urinary tract infection was reported more frequently in females treated with empagliflozin compared to placebo; there was no difference in males.

### Volume depletion

The overall frequency of volume depletion (including the predefined terms blood pressure (ambulatory) decreased, blood pressure systolic decreased, dehydration, hypotension, hypovolaemia, orthostatic hypotension, and syncope) was similar in patients treated with empagliflozin (empagliflozin 10 mg: 0.5%, empagliflozin 25 mg: 0.3%) and placebo (0.3%). The frequency of volume depletion events was increased in patients 75 years and older treated with empagliflozin 10 mg (2.3%) or empagliflozin 25 mg (4.4%) compared to placebo (2.1%).

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via **the national reporting system** listed in [Appendix V](#).

## **4.9 Overdose**

### Symptoms

In controlled clinical studies single doses of up to 800 mg empagliflozin (equivalent to 32 times the highest recommended daily dose) in healthy volunteers and multiple daily doses of up to 100 mg empagliflozin (equivalent to 4 times the highest recommended daily dose) in patients with type 2 diabetes did not show any toxicity. Empagliflozin increased urine glucose excretion leading to an increase in urine volume. The observed increase in urine volume was not dose-dependent and is not clinically meaningful. There is no experience with doses above 800 mg in humans.

### Therapy

In the event of an overdose, treatment should be initiated as appropriate to the patient's clinical status. The removal of empagliflozin by haemodialysis has not been studied.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Drugs used in diabetes, Other blood glucose lowering drugs, excl. insulins, ATC code: A10BX12



## Mechanism of action

Empagliflozin is a reversible, highly potent ( $IC_{50}$  of 1.3 nmol) and selective competitive inhibitor of sodium-glucose co-transporter 2 (SGLT2). Empagliflozin does not inhibit other glucose transporters important for glucose transport into peripheral tissues and is 5000 times more selective for SGLT2 versus SGLT1, the major transporter responsible for glucose absorption in the gut. SGLT2 is highly expressed in the kidney, whereas expression in other tissues is absent or very low. It is responsible, as the predominant transporter, for the reabsorption of glucose from the glomerular filtrate back into the circulation. In patients with type 2 diabetes and hyperglycaemia a higher amount of glucose is filtered and reabsorbed.

Empagliflozin improves glycaemic control in patients with type 2 diabetes by reducing renal glucose reabsorption. The amount of glucose removed by the kidney through this glucuretic mechanism is dependent on blood glucose concentration and GFR. Inhibition of SGLT2 in patients with type 2 diabetes and hyperglycaemia leads to excess glucose excretion in the urine.

In patients with type 2 diabetes, urinary glucose excretion increased immediately following the first dose of empagliflozin and is continuous over the 24 hour dosing interval. Increased urinary glucose excretion was maintained at the end of the 4-week treatment period, averaging approximately 78 g/day. Increased urinary glucose excretion resulted in an immediate reduction in plasma glucose levels in patients with type 2 diabetes.

Empagliflozin improves both fasting and post-prandial plasma glucose levels. The mechanism of action of empagliflozin is independent of beta cell function and insulin pathway and this contributes to a low risk of hypoglycaemia. Improvement of surrogate markers of beta cell function including Homeostasis Model Assessment- (HOMA- ) was noted. In addition, urinary glucose excretion triggers calorie loss, associated with body fat loss and body weight reduction. The glucosuria observed with empagliflozin is accompanied by mild diuresis which may contribute to sustained and moderate reduction of blood pressure.

## Clinical efficacy and safety

A total of 11,250 patients with type 2 diabetes were treated in 10 double-blind, placebo- and active-controlled clinical studies, of which 6,015 received empagliflozin (empagliflozin 10 mg: 3021 patients; empagliflozin 25 mg: 3994 patients). Four studies had treatment durations of 24 weeks; extensions of those and other studies had patients exposed to empagliflozin for up to 102 weeks.

Treatment with empagliflozin as monotherapy and in combination with metformin, pioglitazone, a sulphonylurea, DPP-4 inhibitors, and insulin lead to clinically relevant improvements in HbA1c, fasting plasma glucose (FPG), body weight, and systolic and diastolic blood pressure. Administration of empagliflozin 25 mg resulted in a higher proportion of patients achieving HbA1c goal of less than 7% and fewer patients needing glycaemic rescue compared to empagliflozin 10 mg and placebo. Higher baseline HbA1c was associated with a greater reduction in HbA1c.

### *Monotherapy*

The efficacy and safety of empagliflozin as monotherapy was evaluated in a double-blind, placebo- and active-controlled study of 24 weeks duration in treatment-naïve patients. Treatment with empagliflozin resulted in a statistically significant ( $p < 0.0001$ ) reduction in HbA1c compared to placebo (Table 2) and a clinically meaningful decrease in FPG.

In a prespecified analysis of patients ( $N=201$ ) with a baseline HbA1c  $\geq 8.5\%$ , treatment resulted in a reduction in HbA1c from baseline of  $-1.44\%$  for empagliflozin 10 mg,  $-1.43\%$  for empagliflozin 25 mg,  $-1.04\%$  for sitagliptin, and an increase of  $0.01\%$  for placebo.

In the double-blind placebo-controlled extension of this study, reductions of HbA1c, body weight and blood pressure were sustained up to Week 52.

Table 2: Efficacy results of a 24 week placebo-controlled study of empagliflozin as monotherapy<sup>a</sup>

	Placebo	Jardiance		Sitagliptin
		10 mg	25 mg	100 mg
N	228	224	224	223
<b>HbA1c (%)</b>				
Baseline (mean)	7.91	7.87	7.86	7.85
Change from baseline <sup>1</sup>	0.08	-0.66	-0.78	-0.66
Difference from placebo <sup>1</sup> (97.5% CI)		-0.74* (-0.90, -0.57)	-0.85* (-1.01, -0.69)	-0.73 (-0.88, -0.59) <sup>3</sup>
N	208	204	202	200
<b>Patients (%) achieving HbA1c &lt;7% with baseline HbA1c ≥7%<sup>2</sup></b>	12.0	35.3	43.6	37.5
N	228	224	224	223
<b>Body Weight (kg)</b>				
Baseline (mean)	78.23	78.35	77.80	79.31
Change from baseline <sup>1</sup>	-0.33	-2.26	-2.48	0.18
Difference from placebo <sup>1</sup> (97.5% CI)		-1.93* (-2.48, -1.38)	-2.15* (-2.70, -1.60)	0.52 (-0.04, 1.00) <sup>3</sup>
N	228	224	224	223
<b>SBP (mmHg)<sup>4</sup></b>				
Baseline (mean)	130.4	133.0	129.9	132.5
Change from baseline <sup>1</sup>	-0.3	-2.9	-3.7	0.5
Difference from placebo <sup>1</sup> (95% CI)		-2.6* (-5.2, -0.0)	-3.4* (-6.0, -0.9)	0.8 (-1.4, 3.1) <sup>3</sup>

<sup>a</sup> Full analysis set (FAS) using last observation carried forward (LOCF) prior to glycaemic rescue therapy

<sup>1</sup> Mean adjusted for baseline value

<sup>2</sup> Not evaluated for statistical significance as a result of the sequential confirmatory testing procedure

<sup>3</sup> 95% CI

<sup>4</sup> LOCF, values after antihypertensive rescue censored

\*p-value <0.0001

#### Combination therapy

##### *Empagliflozin as add on to metformin, sulphonylurea, pioglitazone*

Empagliflozin as add-on to metformin, metformin and a sulphonylurea, or pioglitazone with or without metformin resulted in statistically significant (p<0.0001) reductions in HbA1c and body weight compared to placebo (Table 3). In addition it resulted in a clinically meaningful reduction in FPG, systolic and diastolic blood pressure compared to placebo.

In the double-blind placebo-controlled extension of these studies, reduction of HbA1c, body weight and blood pressure were sustained up to Week 52.

Table 3: Efficacy results of 24 week placebo-controlled studies<sup>a</sup>

<b>Add-on to metformin therapy</b>			
	<b>Placebo</b>	<b>Jardiance</b>	
		<b>10 mg</b>	<b>25 mg</b>
N	207	217	213
<b>HbA1c (%)</b>			
Baseline (mean)	7.90	7.94	7.86
Change from baseline <sup>1</sup>	-0.13	-0.70	-0.77
Difference from placebo <sup>1</sup> (97.5% CI)		-0.57* (-0.72, -0.42)	-0.64* (-0.79, -0.48)
N	184	199	191
<b>Patients (%) achieving HbA1c &lt;7% with baseline HbA1c 7%<sup>2</sup></b>	12.5	37.7	38.7
N	207	217	213
<b>Body Weight (kg)</b>			
Baseline (mean)	79.73	81.59	82.21
Change from baseline <sup>1</sup>	-0.45	-2.08	-2.46
Difference from placebo <sup>1</sup> (97.5% CI)		-1.63* (-2.17, -1.08)	-2.01* (-2.56, -1.46)
N	207	217	213
<b>SBP (mmHg)<sup>2</sup></b>			
Baseline (mean)	128.6	129.6	130.0
Change from baseline <sup>1</sup>	-0.4	-4.5	-5.2
Difference from placebo <sup>1</sup> (95% CI)		-4.1* (-6.2, -2.1)	-4.8* (-6.9, -2.7)
<b>Add-on to metformin and a sulphonylurea therapy</b>			
	<b>Placebo</b>	<b>Jardiance</b>	
		<b>10 mg</b>	<b>25 mg</b>
N	225	225	216
<b>HbA1c (%)</b>			
Baseline (mean)	8.15	8.07	8.10
Change from baseline <sup>1</sup>	-0.17	-0.82	-0.77
Difference from placebo <sup>1</sup> (97.5% CI)		-0.64* (-0.79, -0.49)	-0.59* (-0.74, -0.44)
N	216	209	202
<b>Patients (%) achieving HbA1c &lt;7% with baseline HbA1c 7%<sup>2</sup></b>	9.3	26.3	32.2
N	225	225	216
<b>Body Weight (kg)</b>			
Baseline (mean)	76.23	77.08	77.50
Change from baseline <sup>1</sup>	-0.39	-2.16	-2.39
Difference from placebo <sup>1</sup> (97.5% CI)		-1.76* (-2.25, -1.28)	-1.99* (-2.48, -1.50)
N	225	225	216
<b>SBP (mmHg)<sup>2</sup></b>			
Baseline (mean)	128.8	128.7	129.3
Change from baseline <sup>1</sup>	-1.4	-4.1	-3.5
Difference from placebo <sup>1</sup> (95% CI)		-2.7 (-4.6, -0.8)	-2.1 (-4.0, -0.2)

<b>Add-on to pioglitazone +/- metformin therapy</b>			
	<b>Placebo</b>	<b>Jardiance</b>	
		<b>10 mg</b>	<b>25 mg</b>
N	165	165	168
<b>HbA1c (%)</b>			
Baseline (mean)	8.16	8.07	8.06
Change from baseline <sup>1</sup>	-0.11	-0.59	-0.72
Difference from placebo <sup>1</sup> (97.5% CI)		-0.48* (-0.69, -0.27)	-0.61* (-0.82, -0.40)
N	155	151	160
<b>Patients (%) achieving HbA1c &lt;7% with baseline HbA1c 7%<sup>2</sup></b>	7.7	24	30
N	165	165	168
<b>Body Weight (kg)</b>			
Baseline (mean)	78.1	77.97	78.93
Change from baseline <sup>1</sup>	0.34	-1.62	-1.47
Difference from placebo <sup>1</sup> (97.5% CI)		-1.95* (-2.64, -1.27)	-1.81* (-2.49, -1.13)
N	165	165	168
<b>SBP (mmHg)<sup>3</sup></b>			
Baseline (mean)	125.7	126.5	126
Change from baseline <sup>1</sup>	0.7	-3.1	-4.0
Difference from placebo <sup>1</sup> (95% CI)		-3.9 (-6.23, -1.50)	-4.7 (-7.08, -2.37)

<sup>a</sup> Full analysis set (FAS) using last observation carried forward (LOCF) prior to glycaemic rescue therapy

<sup>1</sup> Mean adjusted for baseline value

<sup>2</sup> Not evaluated for statistical significance as a result of the sequential confirmatory testing procedure

<sup>3</sup> LOCF, values after antihypertensive rescue censored

\* p-value <0.0001

*Empagliflozin 24 months data, as add on to metformin in comparison to glimepiride*

In a study comparing the efficacy and safety of empagliflozin 25 mg versus glimepiride (up to 4 mg per day) in patients with inadequate glycaemic control on metformin alone, treatment with empagliflozin daily resulted in superior reduction in HbA1c (Table 4), and a clinically meaningful reduction in FPG, compared to glimepiride. Empagliflozin daily resulted in a statistically significant reduction in body weight, systolic and diastolic blood pressure and a statistically significantly lower proportion of patients with hypoglycaemic events compared to glimepiride (1.6% for empagliflozin, 20.4% for glimepiride, p<0.0001).

Table 4: Efficacy results at 104 week in an active controlled study comparing empagliflozin to glimepiride as add on to metformin<sup>a</sup>

	<b>Empagliflozin 25 mg</b>	<b>Glimepiride<sup>b</sup></b>
N	765	780
<b>HbA1c (%)</b>		
Baseline (mean)	7.92	7.92
Change from baseline <sup>1</sup>	-0.66	-0.55
Difference from glimepiride <sup>1</sup> (97.5% CI)	-0.11* (-0.20, -0.01)	
N	690	715
<b>Patients (%) achieving HbA1c &lt;7% with baseline HbA1c 7%<sup>2</sup></b>	33.6	30.9
N	765	780
<b>Body Weight (kg)</b>		
Baseline (mean)	82.52	83.03
Change from baseline <sup>1</sup>	-3.12	1.34
Difference from glimepiride <sup>1</sup> (97.5% CI)	-4.46** (-4.87, -4.05)	
N	765	780
<b>SBP (mmHg)<sup>2</sup></b>		
Baseline (mean)	133.4	133.5
Change from baseline <sup>1</sup>	-3.1	2.5
Difference from glimepiride <sup>1</sup> (97.5% CI)	-5.6** (-7.0, -4.2)	

<sup>a</sup> Full analysis set (FAS) using last observation carried forward (LOCF) prior to glycaemic rescue therapy

<sup>b</sup> Up to 4 mg glimepiride

<sup>1</sup> Mean adjusted for baseline value

<sup>2</sup> LOCF, values after antihypertensive rescue censored

\* p-value <0.0001 for non-inferiority, and p-value = 0.0153 for superiority

\*\* p-value <0.0001

#### *Add-on to insulin therapy*

##### *Empagliflozin as add on to multiple daily insulin*

The efficacy and safety of empagliflozin as add-on to multiple daily insulin with or without concomitant metformin therapy was evaluated in a double-blind, placebo-controlled trial of 52 weeks duration. During the initial 18 weeks and the last 12 weeks, the insulin dose was kept stable, but was adjusted to achieve pre-prandial glucose levels <100 mg/dl [5.5 mmol/l], and post-prandial glucose levels <140 mg/dl [7.8 mmol/l] between Weeks 19 and 40.

At Week 18, empagliflozin provided statistically significant improvement in HbA1c compared with placebo (Table 5).

At Week 52, treatment with empagliflozin resulted in a statistically significant decrease in HbA1c and insulin sparing compared with placebo and a reduction in FPG and body weight.

Table 5: Efficacy results at 18 and 52 weeks in a placebo-controlled study of empagliflozin as add on to multiple daily doses of insulin with or without metformin

	<b>Placebo</b>	<b>Jardiance</b>	
		<b>10 mg</b>	<b>25 mg</b>
N	188	186	189
<b>HbA1c (%) at week 18</b>			
Baseline (mean)	8.33	8.39	8.29
Change from baseline <sup>1</sup>	-0.50	-0.94	-1.02
Difference from placebo <sup>1</sup> (97.5% CI)		-0.44* (-0.61, -0.27)	-0.52* (-0.69, -0.35)
N	115	119	118
<b>HbA1c (%) at week 52<sup>2</sup></b>			
Baseline (mean)	8.25	8.40	8.37
Change from baseline <sup>1</sup>	-0.81	-1.18	-1.27
Difference from placebo <sup>1</sup>		-0.38*** (-0.62, -0.13)	-0.46* (-0.70, -0.22)

(97.5% CI)			
N	113	118	118
<b>Patients (%) achieving HbA1c &lt;7% with baseline HbA1c 7% at week 52</b>	26.5	39.8	45.8
N	115	118	117
<b>Insulin dose (IU/day) at week 52<sup>2</sup></b>			
Baseline (mean)	89.94	88.57	90.38
Change from baseline <sup>1</sup>	10.16	1.33	-1.06
Difference from placebo <sup>1</sup> (97.5% CI)		-8.83 <sup>#</sup> (-15.69, -1.97)	-11.22** (-18.09, -4.36)
N	115	119	118
<b>Body Weight (kg) at week 52<sup>2</sup></b>			
Baseline (mean)	96.34	96.47	95.37
Change from baseline <sup>1</sup>	0.44	-1.95	-2.04
Difference from placebo <sup>1</sup> (97.5% CI)		-2.39* (-3.54, -1.24)	-2.48* (-3.63, -1.33)

<sup>1</sup> Mean adjusted for baseline value

<sup>2</sup> Week 19-40: treat-to-target regimen for insulin dose adjustment to achieve pre-defined glucose target levels (pre-prandial <100 mg/dl (5.5 mmol/l), post-prandial <140 mg/dl (7.8 mmol/l))

\* p-value <0.0001

\*\* p-value = 0.0003

\*\*\* p-value = 0.0005

# p-value = 0.0040

#### *Empagliflozin as add on to basal insulin*

The efficacy and safety of empagliflozin as add on to basal insulin with or without metformin and/or a sulphonylurea was evaluated in a double-blind, placebo-controlled trial of 78 weeks duration. During the initial 18 weeks the insulin dose was kept stable, but was adjusted to achieve a FPG <110 mg/dl in the following 60 weeks.

At week 18, empagliflozin provided statistically significant improvement in HbA1c (Table 6).

At 78 weeks, empagliflozin resulted in a statistically significant decrease in HbA1c and insulin sparing compared to placebo. Furthermore, empagliflozin resulted in a reduction in FPG, body weight, and blood pressure.

Table 6 Efficacy results at 18 and 78 weeks in a placebo-controlled study of empagliflozin as add on to basal insulin with or without metformin or a sulphonylurea<sup>a</sup>

	Placebo	Empagliflozin 10 mg	Empagliflozin 25 mg
N	125	132	117
<b>HbA1c (%) at week 18</b>			
Baseline (mean)	8.10	8.26	8.34
Change from baseline <sup>1</sup>	-0.01	-0.57	-0.71
Difference from placebo <sup>1</sup> (97.5% CI)		-0.56* (-0.78, -0.33)	-0.70* (-0.93, -0.47)
N	112	127	110
<b>HbA1c (%) at week 78</b>			
Baseline (mean)	8.09	8.27	8.29
Change from baseline <sup>1</sup>	-0.02	-0.48	-0.64
Difference from placebo <sup>1</sup> (97.5% CI)		-0.46* (-0.73, -0.19)	-0.62* (-0.90, -0.34)
N	112	127	110
<b>Basal insulin dose (IU/day) at</b>			

<b>week 78</b>			
Baseline (mean)	47.84	45.13	48.43
Change from baseline <sup>1</sup>	5.45	-1.21	-0.47
Difference from placebo <sup>1</sup> (97.5% CI)		-6.66** (-11.56, -1.77)	-5.92** (-11.00, -0.85)

<sup>a</sup> Full analysis set (FAS) - Completers using last observation carried forward (LOCF) prior to glycaemic rescue therapy

<sup>1</sup> mean adjusted for baseline value

\* p-value <0.0001

\*\* p-value <0.025

*Patients with renal impairment, 52 week placebo controlled data*

The efficacy and safety of empagliflozin as add on to antidiabetic therapy was evaluated in patients with renal impairment in a double-blind, placebo-controlled study for 52 weeks. Treatment with empagliflozin led to a statistically significant reduction of HbA1c (Table 7) and clinically meaningful improvement in FPG compared to placebo at Week 24. The improvement in HbA1c, body weight, and blood pressure was sustained up to 52 weeks.

Table 7 Results at 24 week in a placebo-controlled study of empagliflozin in renally impaired type 2 diabetes patients<sup>a</sup>

	Placebo	Empagliflozin 10 mg	Empagliflozin 25 mg	Placebo	Empagliflozin 25 mg
	eGFR 60 to <90 ml/min/1.73 m <sup>2</sup>			eGFR 45 to <60 ml/min/1.73 m <sup>2</sup>	
N	95	98	97	89	91
<b>HbA1c (%)</b>					
Baseline (mean)	8.09	8.02	7.96	8.08	8.12
Change from baseline <sup>1</sup>	0.06	-0.46	-0.63	-0.08	-0.54
Difference from placebo <sup>1</sup> (95% CI)		-0.52* (-0.72, -0.32)	-0.68* (-0.88, -0.49)		-0.46 (-0.66, -0.27)
N	89	94	91	84	86
<b>Patients (%) achieving HbA1c &lt;7% with baseline HbA1c 7%<sup>2</sup></b>					
	6.7	17.0	24.2	10.7	15.1
N	95	98	97	89	91
<b>Body Weight (kg)<sup>2</sup></b>					
Baseline (mean)	86.00	92.05	88.06	83.20	84.90
Change from baseline <sup>1</sup>	-0.33	-1.76	-2.33	-0.11	-1.39
Difference from placebo <sup>1</sup> (95% CI)		-1.43 (-2.09, -0.77)	-2.00 (-2.66, -1.34)		-1.28 (-2.08, -0.48)
N	95	98	97	89	91
<b>SBP (mmHg)<sup>2</sup></b>					
Baseline (mean)	134.69	137.37	133.68	137.29	135.04
Change from baseline <sup>1</sup>	0.65	-2.92	-4.47	0.37	-5.69
Difference from placebo <sup>1</sup> (95% CI)		-3.57 (-6.86, -0.29)	-5.12 (-8.41, -1.82)		-6.07 (-9.79, -2.34)

<sup>a</sup> Full analysis set (FAS) using last observation carried forward (LOCF) prior to glycaemic rescue therapy

<sup>1</sup> Mean adjusted for baseline value

<sup>2</sup> Not evaluated for statistical significance as a result of the sequential confirmatory testing procedure

\* p<0.0001

### Cardiovascular safety

In a prospective, pre-specified meta-analysis of independently adjudicated cardiovascular events from 12 phase 2 and 3 clinical studies involving 10,036 patients with type 2 diabetes, empagliflozin did not increase cardiovascular risk.

### Fasting plasma glucose

In four placebo-controlled studies, treatment with empagliflozin as monotherapy or add-on therapy to metformin, pioglitazone, or metformin plus a sulfonylurea resulted in mean changes from baseline in FPG of -20.5 mg/dl [-1.14 mmol/l] for empagliflozin 10 mg and -23.2 mg/dl [-1.29 mmol/l] for empagliflozin 25 mg compared to placebo (7.4 mg/dl [0.41 mmol/l]). This effect was observed after 24 weeks and maintained for 76 weeks.

### 2-hour post-prandial glucose

Treatment with empagliflozin as add-on to metformin or metformin and a sulphonylurea resulted in a clinically meaningful reduction of 2-hour post-prandial glucose (meal tolerance test) at 24 weeks (add-on to metformin: placebo +5.9 mg/dl, empagliflozin 10 mg: -46.0 mg/dl, empagliflozin 25 mg: -44.6 mg/dl, add-on to metformin and a sulphonylurea: placebo -2.3 mg/dl, empagliflozin 10 mg: -35.7 mg/dl, empagliflozin 25 mg: -36.6 mg/dl).

### Patients with high baseline HbA1c >10%

In a pre-specified pooled analysis of three phase 3 studies, treatment with open-label empagliflozin 25 mg in patients with severe hyperglycaemia (N=257, mean baseline HbA1c 11.26%) resulted in a clinically meaningful reduction in HbA1c from baseline of 3.27%; no placebo or empagliflozin 10 mg arms were included in these studies.

### Body weight

In a pre-specified pooled analysis of 4 placebo controlled studies, treatment with empagliflozin resulted in body weight reduction (-0.24 kg for placebo, -2.04 kg for empagliflozin 10 mg and -2.26 kg for empagliflozin 25 mg) at week 24 that was maintained up to week 52 (-0.16 kg for placebo, -1.96 kg for empagliflozin 10 mg and -2.25 kg for empagliflozin 25 mg).

### Blood pressure

The efficacy and safety of empagliflozin was evaluated in a double-blind, placebo controlled study of 12 weeks duration in patients with type 2 diabetes and high blood pressure on different antidiabetic and up to 2 antihypertensive therapies. Treatment with empagliflozin once daily resulted in statistically significant improvement in HbA1c, and 24 hour mean systolic and diastolic blood pressure as determined by ambulatory blood pressure monitoring (Table 8). Treatment with empagliflozin provided reductions in seated SBP and DBP.

Table 8 Efficacy results at 12 week in a placebo-controlled study of empagliflozin in patients with type 2 diabetes and uncontrolled blood pressure<sup>a</sup>

	Placebo	Jardiance	
		10 mg	25 mg
N	271	276	276
<b>HbA1c (%) at week 12<sup>1</sup></b>			
Baseline (mean)	7.90	7.87	7.92
Change from baseline <sup>2</sup>	0.03	-0.59	-0.62
Difference from placebo <sup>2</sup> (95% CI)		-0.62* (-0.72, -0.52)	-0.65* (-0.75, -0.55)
<b>24 hour SBP at week 12<sup>3</sup></b>			
Baseline (mean)	131.72	131.34	131.18



Change from baseline <sup>4</sup>	0.48	-2.95	-3.68
Difference from placebo <sup>4</sup> (95% CI)		-3.44* (-4.78, -2.09)	-4.16* (-5.50, -2.83)
<b>24 hour DBP at week 12<sup>3</sup></b>			
Baseline (mean)	75.16	75.13	74.64
Change from baseline <sup>5</sup>	0.32	-1.04	-1.40
Difference from placebo <sup>5</sup> (95% CI)		-1.36** (-2.15, -0.56)	-1.72* (-2.51, -0.93)

<sup>a</sup> Full analysis set (FAS)

<sup>1</sup> LOCF, values after taking antidiabetic rescue therapy censored

<sup>2</sup> Mean adjusted for baseline HbA1c, baseline eGFR, geographical region and number of antihypertensive medicinal products

<sup>3</sup> LOCF, values after taking antidiabetic rescue therapy or changing antihypertensive rescue therapy censored

<sup>4</sup> Mean adjusted for baseline SBP, baseline HbA1c, baseline eGFR, geographical region and number of antihypertensive medicinal products

<sup>5</sup> Mean adjusted for baseline DBP, baseline HbA1c, baseline eGFR, geographical region and number of antihypertensive medicinal products

\* p-value <0.0001

\*\* p-value <0.001

In a pre-specified pooled analysis of 4 placebo-controlled studies, treatment with empagliflozin resulted in a reduction in systolic blood pressure (empagliflozin 10 mg: -3.9 mmHg; empagliflozin 25 mg: -4.3 mmHg) compared with placebo (-0.5 mmHg) and in diastolic blood pressure (empagliflozin 10 mg: -1.8 mmHg; empagliflozin 25 mg: -2.0 mmHg) compared with placebo (-0.5 mmHg) at week 24 that were maintained up to week 52.

### Paediatric population

The European Medicines Agency has deferred the obligation to submit the results of studies with Jardiance in one or more subsets of the paediatric population in type 2 diabetes mellitus (see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

### Absorption

The pharmacokinetics of empagliflozin have been extensively characterised in healthy volunteers and patients with type 2 diabetes. After oral administration, empagliflozin was rapidly absorbed with peak plasma concentrations occurring at a median  $t_{max}$  of 1.5 hours post-dose. Thereafter, plasma concentrations declined in a biphasic manner with a rapid distribution phase and a relatively slow terminal phase. The steady state mean plasma AUC and  $C_{max}$  were 1870 nmol.h and 259 nmol/l with empagliflozin 10 mg and 4740 nmol.h and 687 nmol/l with empagliflozin 25 mg once daily. Systemic exposure of empagliflozin increased in a dose-proportional manner. The single-dose and steady-state pharmacokinetic parameters of empagliflozin were similar suggesting linear pharmacokinetics with respect to time. There were no clinically relevant differences in empagliflozin pharmacokinetics between healthy volunteers and patients with type 2 diabetes.

Administration of empagliflozin 25 mg after intake of a high-fat and high calorie meal resulted in slightly lower exposure; AUC decreased by approximately 16% and  $C_{max}$  by approximately 37% compared to fasted condition. The observed effect of food on empagliflozin pharmacokinetics was not considered clinically relevant and empagliflozin may be administered with or without food.

### Distribution

The apparent steady-state volume of distribution was estimated to be 73.8 l based on the population pharmacokinetic analysis. Following administration of an oral [<sup>14</sup>C]-empagliflozin solution to healthy

volunteers, the red blood cell partitioning was approximately 37% and plasma protein binding was 86%.

### Biotransformation

No major metabolites of empagliflozin were detected in human plasma and the most abundant metabolites were three glucuronide conjugates (2-, 3-, and 6-O glucuronide). Systemic exposure of each metabolite was less than 10% of total drug-related material. *In vitro* studies suggested that the primary route of metabolism of empagliflozin in humans is glucuronidation by the uridine 5'-diphospho-glucuronosyltransferases UGT2B7, UGT1A3, UGT1A8, and UGT1A9.

### Elimination

Based on the population pharmacokinetic analysis, the apparent terminal elimination half-life of empagliflozin was estimated to be 12.4 hours and apparent oral clearance was 10.6 l/hour. The inter-subject and residual variabilities for empagliflozin oral clearance were 39.1% and 35.8%, respectively. With once-daily dosing, steady-state plasma concentrations of empagliflozin were reached by the fifth dose. Consistent with the half-life, up to 22% accumulation, with respect to plasma AUC, was observed at steady-state. Following administration of an oral [<sup>14</sup>C]-empagliflozin solution to healthy volunteers, approximately 96% of the drug-related radioactivity was eliminated in faeces (41%) or urine (54%). The majority of drug-related radioactivity recovered in faeces was unchanged parent drug and approximately half of drug related radioactivity excreted in urine was unchanged parent drug.

### Special populations

#### *Renal impairment*

In patients with mild, moderate or severe renal impairment (eGFR <30 - <90 ml/min/1.73 m<sup>2</sup>) and patients with kidney failure/end stage renal disease (ESRD), AUC of empagliflozin increased by approximately 18%, 20%, 66%, and 48%, respectively compared to subjects with normal renal function. Peak plasma levels of empagliflozin were similar in subjects with moderate renal impairment and kidney failure/ESRD compared to patients with normal renal function. Peak plasma levels of empagliflozin were roughly 20% higher in subjects with mild and severe renal impairment as compared to subjects with normal renal function. The population pharmacokinetic analysis showed that the apparent oral clearance of empagliflozin decreased with a decrease in eGFR leading to an increase in drug exposure.

#### *Hepatic impairment*

In subjects with mild, moderate, and severe hepatic impairment according to the Child-Pugh classification, AUC of empagliflozin increased approximately by 23%, 47%, and 75% and C<sub>max</sub> by approximately 4%, 23%, and 48%, respectively, compared to subjects with normal hepatic function.

#### *Body Mass Index*

Body mass index had no clinically relevant effect on the pharmacokinetics of empagliflozin based on the population pharmacokinetic analysis. In this analysis, AUC was estimated to be 5.82%, 10.4%, and 17.3% lower in subjects with BMI of 30, 35, and 45 kg/m<sup>2</sup>, respectively, compared to subjects with a body mass index of 25 kg/m<sup>2</sup>.

#### *Gender*

Gender had no clinically relevant effect on the pharmacokinetics of empagliflozin based on the population pharmacokinetic analysis.

#### *Race*

In the population pharmacokinetic analysis, AUC was estimated to be 13.5% higher in Asians with a body mass index of 25 kg/m<sup>2</sup> compared to non-Asians with a body mass index of 25 kg/m<sup>2</sup>.

### *Elderly patients*

Age did not have a clinically meaningful impact on the pharmacokinetics of empagliflozin based on the population pharmacokinetic analysis.

### *Paediatric patients*

Studies characterising the pharmacokinetics of empagliflozin in paediatric patients have not been performed.

## **5.3 Preclinical safety data**

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, genotoxicity, fertility and early embryonic development.

In long term toxicity studies in rodents and dogs, signs of toxicity were observed at exposures greater than or equal to 10-times the clinical dose of empagliflozin. Most toxicity was consistent with secondary pharmacology related to urinary glucose loss and electrolyte imbalances including decreased body weight and body fat, increased food consumption, diarrhea, dehydration, decreased serum glucose and increases in other serum parameters reflective of increased protein metabolism and gluconeogenesis, urinary changes such as polyuria and glucosuria, and microscopic changes including mineralisation in kidney and some soft and vascular tissues. Microscopic evidence of the effects of exaggerated pharmacology on the kidney observed in some species included tubular dilatation, and tubular and pelvic mineralisation at approximately 4-times the clinical AUC exposure of empagliflozin associated with the 25 mg dose.

Empagliflozin is not genotoxic.

In a 2 year carcinogenicity study, empagliflozin did not increase the incidence of tumors in female rats up to the highest dose of 700 mg/kg/day, which corresponds to approximately 72-times the maximal clinical AUC exposure to empagliflozin. In male rats, treatment-related benign vascular proliferative lesions (haemangiomas) of the mesenteric lymph node were observed at the highest dose, but not at 300 mg/kg/day, which corresponds to approximately 26-times the maximal clinical exposure to empagliflozin. Interstitial cell tumors in the testes were observed with a higher incidence in rats at 300 mg/kg/day and above, but not at 100 mg/kg/day which corresponds to approximately 18-times the maximal clinical exposure to empagliflozin. Both tumors are common in rats and are unlikely to be relevant to humans.

Empagliflozin did not increase the incidence of tumors in female mice at doses up to 1000 mg/kg/day, which corresponds to approximately 62-times the maximal clinical exposure to empagliflozin. Empagliflozin induced renal tumors in male mice at 1000 mg/kg/day, but not at 300 mg/kg/day, which corresponds to approximately 11-times the maximal clinical exposure to empagliflozin. The mode of action for these tumors is dependent on the natural predisposition of the male mouse to renal pathology and a metabolic pathway not reflective of humans. The male mouse renal tumors are considered not relevant to humans.

At exposures sufficiently in excess of exposure in humans after therapeutic doses, empagliflozin had no adverse effects on fertility or early embryonic development. Empagliflozin administered during the period of organogenesis was not teratogenic. Only at maternally toxic doses, empagliflozin also caused bent limb bones in the rat and increased embryofetal loss in the rabbit.

In pre- and postnatal toxicity studies in rats, reduced weight gain of offspring was observed at maternal exposures approximately 4-times the maximal clinical exposure to empagliflozin. No such effect was seen at systemic exposure equal to the maximal clinical exposure to empagliflozin. The relevance of this finding to humans is unclear.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Lactose monohydrate  
Microcrystalline cellulose  
Hydroxypropylcellulose  
Croscarmellose sodium  
Colloidal anhydrous silica  
Magnesium stearate

#### Film coating

Hypromellose  
Titanium dioxide (E171)  
Talc  
Macrogol (400)  
Iron oxide yellow (E172)

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

3 years

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

PVC/aluminium perforated unit dose blisters.  
Pack sizes of 7 x 1, 10 x 1, 14 x 1, 28 x 1, 30 x 1, 60 x 1, 70 x 1, 90 x 1, and 100 x 1 film-coated tablets.

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim International GmbH  
Binger Str. 173  
D-55216 Ingelheim am Rhein  
Germany

## **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/14/930/010  
EU/1/14/930/011

EU/1/14/930/012  
EU/1/14/930/013  
EU/1/14/930/014  
EU/1/14/930/015  
EU/1/14/930/016  
EU/1/14/930/017  
EU/1/14/930/018

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

## 1. NAME OF THE MEDICINAL PRODUCT

Jardiance 25 mg film-coated tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 25 mg empagliflozin.

Excipient with known effect:

Each tablet contains lactose monohydrate equivalent to 107.4 mg lactose anhydrous.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Film-coated tablet (tablet).

Oval, pale yellow, biconvex film-coated tablet debossed with “S25” on one side and the Boehringer Ingelheim logo on the other (tablet length: 11.1 mm, tablet width: 5.6 mm).

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Jardiance is indicated in the treatment of type 2 diabetes mellitus to improve glycaemic control in adults as:

#### Monotherapy

When diet and exercise alone do not provide adequate glycaemic control in patients for whom use of metformin is considered inappropriate due to intolerance.

#### Add-on combination therapy

In combination with other glucose-lowering medicinal products including insulin, when these, together with diet and exercise, do not provide adequate glycaemic control (see sections 4.4, 4.5 and 5.1 for available data on different combinations).

### 4.2 Posology and method of administration

#### Posology

##### Monotherapy and add-on combination

The recommended starting dose is 10 mg empagliflozin once daily for monotherapy and add-on combination therapy with other glucose-lowering medicinal products including insulin. In patients tolerating empagliflozin 10 mg once daily who have an eGFR  $\geq 60$  ml/min/1.73 m<sup>2</sup> and need tighter glycaemic control, the dose can be increased to 25 mg once daily. The maximum daily dose is 25 mg (see below and section 4.4).

When empagliflozin is used in combination with a sulphonylurea or with insulin, a lower dose of the sulphonylurea or insulin may be considered to reduce the risk of hypoglycaemia (see sections 4.5 and 4.8).

### Special populations

#### *Patients with renal impairment*

Due to the mechanism of action, the efficacy of empagliflozin is dependent on renal function. No dose adjustment is required for patients with an eGFR  $\geq 60$  ml/min/1.73 m<sup>2</sup> or CrCl  $\geq 60$  ml/min.

Empagliflozin should not be initiated in patients with an eGFR  $<60$  ml/min/1.73 m<sup>2</sup> or CrCl  $<60$  ml/min. In patients tolerating empagliflozin whose eGFR falls persistently below 60 ml/min/1.73 m<sup>2</sup> or CrCl below 60 ml/min, the dose of empagliflozin should be adjusted to or maintained at 10 mg once daily. Empagliflozin should be discontinued when eGFR is persistently below 45 ml/min/1.73 m<sup>2</sup> or CrCl persistently below 45 ml/min (see sections 4.4, 4.8, 5.1, and 5.2).

Empagliflozin should not be used in patients with end stage renal disease (ESRD) or in patients on dialysis as it is not expected to be effective in these patients (see sections 4.4 and 5.2).

#### *Patients with hepatic impairment*

No dose adjustment is required for patients with hepatic impairment. Empagliflozin exposure is increased in patients with severe hepatic impairment. Therapeutic experience in patients with severe hepatic impairment is limited and therefore not recommended for use in this population (see section 5.2).

#### *Elderly patients*

No dose adjustment is recommended based on age. In patients 75 years and older, an increased risk for volume depletion should be taken into account (see sections 4.4 and 4.8). In patients aged 85 years and older, initiation of empagliflozin therapy is not recommended due to the limited therapeutic experience (see section 4.4).

#### *Paediatric population*

The safety and efficacy of empagliflozin in children and adolescents has not yet been established. No data are available.

### Method of administration

The tablets can be taken with or without food, swallowed whole with water. If a dose is missed, it should be taken as soon as the patient remembers. A double dose should not be taken on the same day.

## **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

## **4.4 Special warnings and precautions for use**

### General

Jardiance should not be used in patients with type 1 diabetes or for the treatment of diabetic ketoacidosis.

### Use in patients with renal impairment

Jardiance should not be initiated in patients with an eGFR below 60 ml/min/1.73 m<sup>2</sup> or CrCl  $<60$  ml/min. In patients tolerating empagliflozin whose eGFR is persistently below 60 ml/min/1.73 m<sup>2</sup> or CrCl  $<60$  ml/min, the dose of empagliflozin should be adjusted to or maintained at 10 mg once daily. Empagliflozin should be discontinued when eGFR is persistently below 45 ml/min/1.73 m<sup>2</sup> or CrCl persistently below 45 ml/min. Empagliflozin should not be used in patients

with ESRD or in patients on dialysis as it is not expected to be effective in these patients (see sections 4.2 and 5.2).

#### Monitoring of renal function

Due to the mechanism of action, the efficacy of empagliflozin is dependent on renal function.

Therefore assessment of renal function is recommended as follows:

- Prior to empagliflozin initiation and periodically during treatment, i.e. at least yearly (see sections 4.2, 5.1 and 5.2)
- Prior to initiation of any concomitant medicinal product that may have a negative impact on renal function.

#### Hepatic injury

Cases of hepatic injury have been reported with empagliflozin in clinical trials. A causal relationship between empagliflozin and hepatic injury has not been established.

#### Elderly patients

The effect of empagliflozin on urinary glucose excretion is associated with osmotic diuresis, which could affect the hydration status. Patients aged 75 years and older may be at an increased risk of volume depletion. A higher number of these patients treated with empagliflozin had adverse reactions related to volume depletion as compared to placebo (see section 4.8).

Therapeutic experience in patients aged 85 years and older is limited. Initiation of empagliflozin therapy in this population is not recommended (see section 4.2).

#### Use in patients at risk for volume depletion

Based on the mode of action of SGLT-2 inhibitors, osmotic diuresis accompanying therapeutic glucosuria may lead to a modest decrease in blood pressure (see section 5.1). Therefore, caution should be exercised in patients for whom an empagliflozin-induced drop in blood pressure could pose a risk, such as patients with known cardiovascular disease, patients on anti-hypertensive therapy with a history of hypotension or patients aged 75 years and older.

In case of conditions that may lead to fluid loss (e.g. gastrointestinal illness), careful monitoring of volume status (e.g. physical examination, blood pressure measurements, laboratory tests including haematocrit) and electrolytes is recommended for patients receiving empagliflozin. Temporary interruption of treatment with empagliflozin should be considered until the fluid loss is corrected.

#### Urinary tract infections

The overall frequency of urinary tract infection reported as adverse event was similar in patients treated with empagliflozin 25 mg and placebo and higher in patients treated with empagliflozin 10 mg (see section 4.8). Complicated urinary tract infection (e.g. pyelonephritis or urosepsis) occurred at a similar frequency in patients treated with empagliflozin compared to placebo. However, temporary interruption of empagliflozin should be considered in patients with complicated urinary tract infections.

#### Cardiac failure

Experience in New York Heart Association (NYHA) class I-II is limited, and there is no experience in clinical studies with empagliflozin in NYHA class III-IV.

#### Urine laboratory assessments

Due to its mechanism of action, patients taking Jardiance will test positive for glucose in their urine.



## Lactose

The tablets contain lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency, or glucose-galactose malabsorption should not take this medicinal product.

## **4.5 Interaction with other medicinal products and other forms of interaction**

### Pharmacodynamic interactions

#### Diuretics

Empagliflozin may add to the diuretic effect of thiazide and loop diuretics and may increase the risk of dehydration and hypotension (see section 4.4).

#### Insulin and insulin secretagogues

Insulin and insulin secretagogues, such as sulphonylureas, may increase the risk of hypoglycaemia. Therefore, a lower dose of insulin or an insulin secretagogue may be required to reduce the risk of hypoglycaemia when used in combination with empagliflozin (see sections 4.2 and 4.8).

### Pharmacokinetic interactions

#### Effects of other medicinal products on empagliflozin

*In vitro* data suggest that the primary route of metabolism of empagliflozin in humans is glucuronidation by uridine 5'-diphosphoglucuronosyltransferases UGT1A3, UGT1A8, UGT1A9, and UGT2B7. Empagliflozin is a substrate of the human uptake transporters OAT3, OATP1B1, and OATP1B3, but not OAT1 and OCT2. Empagliflozin is a substrate of P-glycoprotein (P-gp) and breast cancer resistance protein (BCRP).

Co-administration of empagliflozin with probenecid, an inhibitor of UGT enzymes and OAT3, resulted in a 26% increase in peak empagliflozin plasma concentrations ( $C_{max}$ ) and a 53% increase in area under the concentration-time curve (AUC). These changes were not considered to be clinically meaningful.

The effect of UGT induction on empagliflozin has not been studied. Co-medication with known inducers of UGT enzymes should be avoided due to a potential risk of decreased efficacy.

An interaction study with gemfibrozil, an *in vitro* inhibitor of OAT3 and OATP1B1/1B3 transporters, showed that empagliflozin  $C_{max}$  increased by 15% and AUC increased by 59% following co-administration. These changes were not considered to be clinically meaningful.

Inhibition of OATP1B1/1B3 transporters by coadministration with rifampicin resulted in a 75% increase in  $C_{max}$  and a 35% increase in AUC of empagliflozin. These changes were not considered to be clinically meaningful.

Empagliflozin exposure was similar with and without co-administration with verapamil, a P-gp inhibitor, indicating that inhibition of P-gp does not have any clinically relevant effect on empagliflozin.

Interaction studies conducted in healthy volunteers suggest that the pharmacokinetics of empagliflozin were not influenced by coadministration with metformin, glimepiride, pioglitazone, sitagliptin, linagliptin, warfarin, verapamil, ramipril, simvastatin, torasemide and hydrochlorothiazide.

#### Effects of empagliflozin on other medicinal products

Based on *in vitro* studies, empagliflozin does not inhibit, inactivate, or induce CYP450 isoforms. Empagliflozin does not inhibit UGT1A1. Drug-drug interactions involving the major CYP450 isoforms or UGT1A1 with empagliflozin and concomitantly administered substrates of these enzymes are therefore considered unlikely. The potential for empagliflozin to inhibit UGT2B7 has not been studied.

Empagliflozin does not inhibit P-gp at therapeutic doses. Based on *in vitro* studies, empagliflozin is considered unlikely to cause interactions with drugs that are P-gp substrates. Co-administration of digoxin, a P-gp substrate, with empagliflozin resulted in a 6% increase in AUC and 14% increase in  $C_{max}$  of digoxin. These changes were not considered to be clinically meaningful.

Empagliflozin does not inhibit human uptake transporters such as OAT3, OATP1B1, and OATP1B3 *in vitro* at clinically relevant plasma concentrations and, as such, drug-drug interactions with substrates of these uptake transporters are considered unlikely.

Interaction studies conducted in healthy volunteers suggest that empagliflozin had no clinically relevant effect on the pharmacokinetics of metformin, glimepiride, pioglitazone, sitagliptin, linagliptin, simvastatin, warfarin, ramipiril, digoxin, diuretics and oral contraceptives.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy

There are no data from the use of empagliflozin in pregnant women. Animal studies show that empagliflozin crosses the placenta during late gestation to a very limited extent but do not indicate direct or indirect harmful effects with respect to early embryonic development. However, animal studies have shown adverse effects on postnatal development (see section 5.3). As a precautionary measure, it is preferable to avoid the use of Jardiance during early pregnancy. Jardiance is not recommended during the second and third trimester of pregnancy.

##### Breast-feeding

No data in humans are available on excretion of empagliflozin into milk. Available toxicological data in animals have shown excretion of empagliflozin in milk. A risk to the newborns/infants cannot be excluded. Jardiance should not be used during breast-feeding.

##### Fertility

No studies on the effect on human fertility have been conducted for Jardiance. Animal studies do not indicate direct or indirect harmful effects with respect to fertility (see section 5.3).

#### **4.7 Effects on ability to drive and use machines**

Jardiance has minor influence on the ability to drive and use machines. Patients should be advised to take precautions to avoid hypoglycaemia while driving and using machines, in particular when Jardiance is used in combination with a sulphonylurea and/or insulin.

#### **4.8 Undesirable effects**

##### Summary of the safety profile

A total of 13,076 patients with type 2 diabetes were included in clinical studies to evaluate the safety of empagliflozin. 2856 patients received empagliflozin 10 mg and 3,738 patients received empagliflozin 25 mg for at least 24 weeks and 601 or 881 patients for at least 76 weeks, either alone or in combination with metformin, a sulphonylurea, pioglitazone, DPP-4 inhibitors, or insulin.

In 5 placebo-controlled trials of 18 to 24 weeks duration, 2,971 patients were included of which 995 were treated with placebo and 1,976 with empagliflozin. The overall incidence of adverse events in patients treated with empagliflozin was similar to placebo. The most frequently reported adverse reaction was hypoglycaemia when used with sulphonylurea or insulin (see description of selected adverse reactions).

## Tabulated list of adverse reactions

Adverse reactions classified by system organ class and MedDRA preferred terms reported in patients who received empagliflozin in placebo-controlled studies are presented in the table below (Table 1).

The adverse reactions are listed by absolute frequency. Frequencies are defined as very common ( 1/10), common ( 1/100 to <1/10), uncommon ( 1/1,000 to <1/100), rare ( 1/10,000 to <1/1,000), or very rare (<1/10,000), and not known (cannot be estimated from the available data).

Table 1: Adverse reactions reported in placebo-controlled studies

<b>System organ class</b>	<b>Very common</b>	<b>Common</b>	<b>Uncommon</b>
<i>Infections and infestations</i>		Vaginal moniliasis, vulvovaginitis, balanitis and other genital infection <sup>a</sup> Urinary tract infection <sup>a</sup>	
<i>Metabolism and nutrition disorders</i>	Hypoglycaemia (when used with sulphonylurea or insulin) <sup>a</sup>		
<i>Skin and subcutaneous disorders</i>		Pruritus (generalised)	
<i>Vascular disorders</i>			Volume depletion <sup>a</sup>
<i>Renal and urinary disorders</i>		Increased urination <sup>a</sup>	Dysuria

<sup>a</sup>see subsections below for additional information

## Description of selected adverse reactions

### Hypoglycaemia

The frequency of hypoglycaemia depended on the background therapy in the respective studies.

#### *Minor hypoglycaemia*

The frequency of patients with minor hypoglycaemia was similar for empagliflozin and placebo as monotherapy, add-on to metformin, and add-on to pioglitazone with or without metformin. An increased frequency was noted when given as add-on to metformin and a sulphonylurea (empagliflozin 10 mg: 16.1%, empagliflozin 25 mg: 11.5%, placebo: 8.4%), and as add-on to insulin with or without metformin and with or without a sulphonylurea (empagliflozin 10 mg: 19.5%, empagliflozin 25 mg: 27.1%, placebo: 20.6% during initial 18 weeks treatment when insulin could not be adjusted; empagliflozin 10 mg: 36.1%, empagliflozin 25 mg: 34.8%, placebo 35.3% over the 78-week trial).

#### *Major hypoglycaemia (hypoglycaemia requiring assistance)*

No increase in major hypoglycaemia was observed with empagliflozin compared to placebo as monotherapy, add-on to metformin, add-on to metformin and a sulphonylurea, and add-on to pioglitazone with or without metformin. An increased frequency was noted when given as add-on to insulin with or without metformin and with or without a sulphonylurea (empagliflozin 10 mg: 0%, empagliflozin 25 mg: 1.3%, placebo: 0% during initial 18 weeks treatment when insulin could not be adjusted; empagliflozin 10 mg: 0%, empagliflozin 25 mg: 1.3%, placebo 0% over the 78-week trial).

### Vaginal moniliasis, vulvovaginitis, balanitis and other genital infection

Vaginal moniliasis, vulvovaginitis, balanitis and other genital infections were reported more frequently in patients treated with empagliflozin (empagliflozin 10 mg: 4.1%, empagliflozin 25 mg: 3.7%) compared to placebo (0.9%). These infections were reported more frequently in females treated with empagliflozin compared to placebo, and the difference in frequency was less pronounced in males. The genital tract infections were mild or moderate in intensity.

### Increased urination

Increased urination (including the predefined terms pollakiuria, polyuria, and nocturia) was observed at higher frequencies in patients treated with empagliflozin (empagliflozin 10 mg: 3.4%, empagliflozin 25 mg: 3.2%) compared to placebo (1.0%). Increased urination was mostly mild or moderate in intensity. The frequency of reported nocturia was similar for placebo and empagliflozin (<1%).

### Urinary tract infection

The overall frequency of urinary tract infection reported as adverse event was similar in patients treated with empagliflozin 25 mg and placebo (7.6%) and higher in empagliflozin 10 mg (9.3%). Similar to placebo, urinary tract infection was reported more frequently for empagliflozin in patients with a history of chronic or recurrent urinary tract infections. The intensity (mild, moderate, severe) of urinary tract infection was similar in patients treated with empagliflozin and placebo. Urinary tract infection was reported more frequently in females treated with empagliflozin compared to placebo; there was no difference in males.

### Volume depletion

The overall frequency of volume depletion (including the predefined terms blood pressure (ambulatory) decreased, blood pressure systolic decreased, dehydration, hypotension, hypovolaemia, orthostatic hypotension, and syncope) was similar in patients treated with empagliflozin (empagliflozin 10 mg: 0.5%, empagliflozin 25 mg: 0.3%) and placebo (0.3%). The frequency of volume depletion events was increased in patients 75 years and older treated with empagliflozin 10 mg (2.3%) or empagliflozin 25 mg (4.4%) compared to placebo (2.1%).

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

## **4.9 Overdose**

### Symptoms

In controlled clinical studies single doses of up to 800 mg empagliflozin (equivalent to 32 times the highest recommended daily dose) in healthy volunteers and multiple daily doses of up to 100 mg empagliflozin (equivalent to 4 times the highest recommended daily dose) in patients with type 2 diabetes did not show any toxicity. Empagliflozin increased urine glucose excretion leading to an increase in urine volume. The observed increase in urine volume was not dose-dependent and is not clinically meaningful. There is no experience with doses above 800 mg in humans.

### Therapy

In the event of an overdose, treatment should be initiated as appropriate to the patient's clinical status. The removal of empagliflozin by haemodialysis has not been studied.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Drugs used in diabetes, Other blood glucose lowering drugs, excl. insulins, ATC code: A10BX12

## Mechanism of action

Empagliflozin is a reversible, highly potent (IC<sub>50</sub> of 1.3 nmol) and selective competitive inhibitor of sodium-glucose co-transporter 2 (SGLT2). Empagliflozin does not inhibit other glucose transporters important for glucose transport into peripheral tissues and is 5000 times more selective for SGLT2 versus SGLT1, the major transporter responsible for glucose absorption in the gut. SGLT2 is highly expressed in the kidney, whereas expression in other tissues is absent or very low. It is responsible, as the predominant transporter, for the reabsorption of glucose from the glomerular filtrate back into the circulation. In patients with type 2 diabetes and hyperglycaemia a higher amount of glucose is filtered and reabsorbed.

Empagliflozin improves glycaemic control in patients with type 2 diabetes by reducing renal glucose reabsorption. The amount of glucose removed by the kidney through this glucuretic mechanism is dependent on blood glucose concentration and GFR. Inhibition of SGLT2 in patients with type 2 diabetes and hyperglycaemia leads to excess glucose excretion in the urine.

In patients with type 2 diabetes, urinary glucose excretion increased immediately following the first dose of empagliflozin and is continuous over the 24 hour dosing interval. Increased urinary glucose excretion was maintained at the end of the 4-week treatment period, averaging approximately 78 g/day. Increased urinary glucose excretion resulted in an immediate reduction in plasma glucose levels in patients with type 2 diabetes.

Empagliflozin improves both fasting and post-prandial plasma glucose levels. The mechanism of action of empagliflozin is independent of beta cell function and insulin pathway and this contributes to a low risk of hypoglycaemia. Improvement of surrogate markers of beta cell function including Homeostasis Model Assessment- (HOMA- ) was noted. In addition, urinary glucose excretion triggers calorie loss, associated with body fat loss and body weight reduction. The glucosuria observed with empagliflozin is accompanied by mild diuresis which may contribute to sustained and moderate reduction of blood pressure.

## Clinical efficacy and safety

A total of 11,250 patients with type 2 diabetes were treated in 10 double-blind, placebo- and active-controlled clinical studies, of which 6,015 received empagliflozin (empagliflozin 10 mg: 3021 patients; empagliflozin 25 mg: 3994 patients). Four studies had treatment durations of 24 weeks; extensions of those and other studies had patients exposed to empagliflozin for up to 102 weeks.

Treatment with empagliflozin as monotherapy and in combination with metformin, pioglitazone, a sulphonylurea, DPP-4 inhibitors, and insulin lead to clinically relevant improvements in HbA1c, fasting plasma glucose (FPG), body weight, and systolic and diastolic blood pressure. Administration of empagliflozin 25 mg resulted in a higher proportion of patients achieving HbA1c goal of less than 7% and fewer patients needing glycaemic rescue compared to empagliflozin 10 mg and placebo. Higher baseline HbA1c was associated with a greater reduction in HbA1c.

### *Monotherapy*

The efficacy and safety of empagliflozin as monotherapy was evaluated in a double-blind, placebo- and active-controlled study of 24 weeks duration in treatment-naïve patients. Treatment with empagliflozin resulted in a statistically significant (p<0.0001) reduction in HbA1c compared to placebo (Table 2) and a clinically meaningful decrease in FPG.

In a prespecified analysis of patients (N=201) with a baseline HbA1c 8.5%, treatment resulted in a reduction in HbA1c from baseline of -1.44% for empagliflozin 10 mg, -1.43% for empagliflozin 25 mg, -1.04% for sitagliptin, and an increase of 0.01% for placebo.

In the double-blind placebo-controlled extension of this study, reductions of HbA1c, body weight and blood pressure were sustained up to Week 52.

Table 2: Efficacy results of a 24 week placebo-controlled study of empagliflozin as monotherapy<sup>a</sup>

	Placebo	Jardiance		Sitagliptin
		10 mg	25 mg	100 mg
N	228	224	224	223
<b>HbA1c (%)</b>				
Baseline (mean)	7.91	7.87	7.86	7.85
Change from baseline <sup>1</sup>	0.08	-0.66	-0.78	-0.66
Difference from placebo <sup>1</sup> (97.5% CI)		-0.74* (-0.90, -0.57)	-0.85* (-1.01, -0.69)	-0.73 (-0.88, -0.59) <sup>3</sup>
N	208	204	202	200
<b>Patients (%) achieving HbA1c &lt;7% with baseline HbA1c ≥7%<sup>2</sup></b>	12.0	35.3	43.6	37.5
N	228	224	224	223
<b>Body Weight (kg)</b>				
Baseline (mean)	78.23	78.35	77.80	79.31
Change from baseline <sup>1</sup>	-0.33	-2.26	-2.48	0.18
Difference from placebo <sup>1</sup> (97.5% CI)		-1.93* (-2.48, -1.38)	-2.15* (-2.70, -1.60)	0.52 (-0.04, 1.00) <sup>3</sup>
N	228	224	224	223
<b>SBP (mmHg)<sup>4</sup></b>				
Baseline (mean)	130.4	133.0	129.9	132.5
Change from baseline <sup>1</sup>	-0.3	-2.9	-3.7	0.5
Difference from placebo <sup>1</sup> (95% CI)		-2.6* (-5.2, -0.0)	-3.4* (-6.0, -0.9)	0.8 (-1.4, 3.1) <sup>3</sup>

<sup>a</sup> Full analysis set (FAS) using last observation carried forward (LOCF) prior to glycaemic rescue therapy

<sup>1</sup> Mean adjusted for baseline value

<sup>2</sup> Not evaluated for statistical significance as a result of the sequential confirmatory testing procedure

<sup>3</sup> 95% CI

<sup>4</sup> LOCF, values after antihypertensive rescue censored

\*p-value <0.0001

#### Combination therapy

##### *Empagliflozin as add on to metformin, sulphonylurea, pioglitazone*

Empagliflozin as add-on to metformin, metformin and a sulphonylurea, or pioglitazone with or without metformin resulted in statistically significant (p<0.0001) reductions in HbA1c and body weight compared to placebo (Table 3). In addition it resulted in a clinically meaningful reduction in FPG, systolic and diastolic blood pressure compared to placebo.

In the double-blind placebo-controlled extension of these studies, reduction of HbA1c, body weight and blood pressure were sustained up to Week 52.

Table 3: Efficacy results of 24 week placebo-controlled studies<sup>a</sup>

<b>Add-on to metformin therapy</b>			
	<b>Placebo</b>	<b>Jardiance</b>	
		<b>10 mg</b>	<b>25 mg</b>
N	207	217	213
<b>HbA1c (%)</b>			
Baseline (mean)	7.90	7.94	7.86
Change from baseline <sup>1</sup>	-0.13	-0.70	-0.77
Difference from placebo <sup>1</sup> (97.5% CI)		-0.57* (-0.72, -0.42)	-0.64* (-0.79, -0.48)
N	184	199	191
<b>Patients (%) achieving HbA1c &lt;7% with baseline HbA1c 7%<sup>2</sup></b>	12.5	37.7	38.7
N	207	217	213
<b>Body Weight (kg)</b>			
Baseline (mean)	79.73	81.59	82.21
Change from baseline <sup>1</sup>	-0.45	-2.08	-2.46
Difference from placebo <sup>1</sup> (97.5% CI)		-1.63* (-2.17, -1.08)	-2.01* (-2.56, -1.46)
N	207	217	213
<b>SBP (mmHg)<sup>2</sup></b>			
Baseline (mean)	128.6	129.6	130.0
Change from baseline <sup>1</sup>	-0.4	-4.5	-5.2
Difference from placebo <sup>1</sup> (95% CI)		-4.1* (-6.2, -2.1)	-4.8* (-6.9, -2.7)
<b>Add-on to metformin and a sulphonylurea therapy</b>			
	<b>Placebo</b>	<b>Jardiance</b>	
		<b>10 mg</b>	<b>25 mg</b>
N	225	225	216
<b>HbA1c (%)</b>			
Baseline (mean)	8.15	8.07	8.10
Change from baseline <sup>1</sup>	-0.17	-0.82	-0.77
Difference from placebo <sup>1</sup> (97.5% CI)		-0.64* (-0.79, -0.49)	-0.59* (-0.74, -0.44)
N	216	209	202
<b>Patients (%) achieving HbA1c &lt;7% with baseline HbA1c 7%<sup>2</sup></b>	9.3	26.3	32.2
N	225	225	216
<b>Body Weight (kg)</b>			
Baseline (mean)	76.23	77.08	77.50
Change from baseline <sup>1</sup>	-0.39	-2.16	-2.39
Difference from placebo <sup>1</sup> (97.5% CI)		-1.76* (-2.25, -1.28)	-1.99* (-2.48, -1.50)
N	225	225	216
<b>SBP (mmHg)<sup>2</sup></b>			
Baseline (mean)	128.8	128.7	129.3
Change from baseline <sup>1</sup>	-1.4	-4.1	-3.5
Difference from placebo <sup>1</sup> (95% CI)		-2.7 (-4.6, -0.8)	-2.1 (-4.0, -0.2)

<b>Add-on to pioglitazone +/- metformin therapy</b>			
	<b>Placebo</b>	<b>Jardiance</b>	
		<b>10 mg</b>	<b>25 mg</b>
N	165	165	168
<b>HbA1c (%)</b>			
Baseline (mean)	8.16	8.07	8.06
Change from baseline <sup>1</sup>	-0.11	-0.59	-0.72
Difference from placebo <sup>1</sup> (97.5% CI)		-0.48* (-0.69, -0.27)	-0.61* (-0.82, -0.40)
N	155	151	160
<b>Patients (%) achieving HbA1c &lt;7% with baseline HbA1c 7%<sup>2</sup></b>	7.7	24	30
N	165	165	168
<b>Body Weight (kg)</b>			
Baseline (mean)	78.1	77.97	78.93
Change from baseline <sup>1</sup>	0.34	-1.62	-1.47
Difference from placebo <sup>1</sup> (97.5% CI)		-1.95* (-2.64, -1.27)	-1.81* (-2.49, -1.13)
N	165	165	168
<b>SBP (mmHg)<sup>3</sup></b>			
Baseline (mean)	125.7	126.5	126
Change from baseline <sup>1</sup>	0.7	-3.1	-4.0
Difference from placebo <sup>1</sup> (95% CI)		-3.9 (-6.23, -1.50)	-4.7 (-7.08, -2.37)

<sup>a</sup> Full analysis set (FAS) using last observation carried forward (LOCF) prior to glycaemic rescue therapy

<sup>1</sup> Mean adjusted for baseline value

<sup>2</sup> Not evaluated for statistical significance as a result of the sequential confirmatory testing procedure

<sup>3</sup> LOCF, values after antihypertensive rescue censored

\* p-value <0.0001

#### *Empagliflozin 24 months data, as add on to metformin in comparison to glimepiride*

In a study comparing the efficacy and safety of empagliflozin 25 mg versus glimepiride (up to 4 mg per day) in patients with inadequate glycaemic control on metformin alone, treatment with empagliflozin daily resulted in superior reduction in HbA1c (Table 4), and a clinically meaningful reduction in FPG, compared to glimepiride. Empagliflozin daily resulted in a statistically significant reduction in body weight, systolic and diastolic blood pressure and a statistically significantly lower proportion of patients with hypoglycaemic events compared to glimepiride (1.6% for empagliflozin, 20.4% for glimepiride, p<0.0001).



Table 4: Efficacy results at 104 week in an active controlled study comparing empagliflozin to glimepiride as add on to metformin<sup>a</sup>

	<b>Empagliflozin 25 mg</b>	<b>Glimepiride<sup>b</sup></b>
N	765	780
<b>HbA1c (%)</b>		
Baseline (mean)	7.92	7.92
Change from baseline <sup>1</sup>	-0.66	-0.55
Difference from glimepiride <sup>1</sup> (97.5% CI)	-0.11* (-0.20, -0.01)	
N	690	715
<b>Patients (%) achieving HbA1c &lt;7% with baseline HbA1c 7%<sup>2</sup></b>	33.6	30.9
N	765	780
<b>Body Weight (kg)</b>		
Baseline (mean)	82.52	83.03
Change from baseline <sup>1</sup>	-3.12	1.34
Difference from glimepiride <sup>1</sup> (97.5% CI)	-4.46** (-4.87, -4.05)	
N	765	780
<b>SBP (mmHg)<sup>2</sup></b>		
Baseline (mean)	133.4	133.5
Change from baseline <sup>1</sup>	-3.1	2.5
Difference from glimepiride <sup>1</sup> (97.5% CI)	-5.6** (-7.0, -4.2)	

<sup>a</sup> Full analysis set (FAS) using last observation carried forward (LOCF) prior to glycaemic rescue therapy

<sup>b</sup> Up to 4 mg glimepiride

<sup>1</sup> Mean adjusted for baseline value

<sup>2</sup> LOCF, values after antihypertensive rescue censored

\* p-value <0.0001 for non-inferiority, and p-value = 0.0153 for superiority

\*\* p-value <0.0001

#### *Add-on to insulin therapy*

##### *Empagliflozin as add on to multiple daily insulin*

The efficacy and safety of empagliflozin as add-on to multiple daily insulin with or without concomitant metformin therapy was evaluated in a double-blind, placebo-controlled trial of 52 weeks duration. During the initial 18 weeks and the last 12 weeks, the insulin dose was kept stable, but was adjusted to achieve pre-prandial glucose levels <100 mg/dl [5.5 mmol/l], and post-prandial glucose levels <140 mg/dl [7.8 mmol/l] between Weeks 19 and 40.

At Week 18, empagliflozin provided statistically significant improvement in HbA1c compared with placebo (Table 5).

At Week 52, treatment with empagliflozin resulted in a statistically significant decrease in HbA1c and insulin sparing compared with placebo and a reduction in FPG and body weight.

Table 5: Efficacy results at 18 and 52 weeks in a placebo-controlled study of empagliflozin as add on to multiple daily doses of insulin with or without metformin

	<b>Placebo</b>	<b>Jardiance</b>	
		<b>10 mg</b>	<b>25 mg</b>
N	188	186	189
<b>HbA1c (%) at week 18</b>			
Baseline (mean)	8.33	8.39	8.29
Change from baseline <sup>1</sup>	-0.50	-0.94	-1.02
Difference from placebo <sup>1</sup> (97.5% CI)		-0.44* (-0.61, -0.27)	-0.52* (-0.69, -0.35)
N	115	119	118
<b>HbA1c (%) at week 52<sup>2</sup></b>			
Baseline (mean)	8.25	8.40	8.37
Change from baseline <sup>1</sup>	-0.81	-1.18	-1.27
Difference from placebo <sup>1</sup>		-0.38*** (-0.62, -0.13)	-0.46* (-0.70, -0.22)

(97.5% CI)			
N	113	118	118
<b>Patients (%) achieving HbA1c &lt;7% with baseline HbA1c 7% at week 52</b>	26.5	39.8	45.8
N	115	118	117
<b>Insulin dose (IU/day) at week 52<sup>2</sup></b>			
Baseline (mean)	89.94	88.57	90.38
Change from baseline <sup>1</sup>	10.16	1.33	-1.06
Difference from placebo <sup>1</sup> (97.5% CI)		-8.83 <sup>#</sup> (-15.69, -1.97)	-11.22** (-18.09, -4.36)
N	115	119	118
<b>Body Weight (kg) at week 52<sup>2</sup></b>			
Baseline (mean)	96.34	96.47	95.37
Change from baseline <sup>1</sup>	0.44	-1.95	-2.04
Difference from placebo <sup>1</sup> (97.5% CI)		-2.39* (-3.54, -1.24)	-2.48* (-3.63, -1.33)

<sup>1</sup> Mean adjusted for baseline value

<sup>2</sup> Week 19-40: treat-to-target regimen for insulin dose adjustment to achieve pre-defined glucose target levels (pre-prandial <100 mg/dl (5.5 mmol/l), post-prandial <140 mg/dl (7.8 mmol/l))

\* p-value <0.0001

\*\* p-value = 0.0003

\*\*\* p-value = 0.0005

# p-value = 0.0040

#### *Empagliflozin as add on to basal insulin*

The efficacy and safety of empagliflozin as add on to basal insulin with or without metformin and/or a sulphonylurea was evaluated in a double-blind, placebo-controlled trial of 78 weeks duration. During the initial 18 weeks the insulin dose was kept stable, but was adjusted to achieve a FPG <110 mg/dl in the following 60 weeks.

At week 18, empagliflozin provided statistically significant improvement in HbA1c (Table 6).

At 78 weeks, empagliflozin resulted in a statistically significant decrease in HbA1c and insulin sparing compared to placebo. Furthermore, empagliflozin resulted in a reduction in FPG, body weight, and blood pressure.

Table 6 Efficacy results at 18 and 78 weeks in a placebo-controlled study of empagliflozin as add on to basal insulin with or without metformin or a sulphonylurea<sup>a</sup>

	Placebo	Empagliflozin 10 mg	Empagliflozin 25 mg
N	125	132	117
<b>HbA1c (%) at week 18</b>			
Baseline (mean)	8.10	8.26	8.34
Change from baseline <sup>1</sup>	-0.01	-0.57	-0.71
Difference from placebo <sup>1</sup> (97.5% CI)		-0.56* (-0.78, -0.33)	-0.70* (-0.93, -0.47)
N	112	127	110
<b>HbA1c (%) at week 78</b>			
Baseline (mean)	8.09	8.27	8.29
Change from baseline <sup>1</sup>	-0.02	-0.48	-0.64
Difference from placebo <sup>1</sup> (97.5% CI)		-0.46* (-0.73, -0.19)	-0.62* (-0.90, -0.34)
N	112	127	110
<b>Basal insulin dose (IU/day) at</b>			

<b>week 78</b>			
Baseline (mean)	47.84	45.13	48.43
Change from baseline <sup>1</sup>	5.45	-1.21	-0.47
Difference from placebo <sup>1</sup> (97.5% CI)		-6.66** (-11.56, -1.77)	-5.92** (-11.00, -0.85)

<sup>a</sup> Full analysis set (FAS) - Completers using last observation carried forward (LOCF) prior to glycaemic rescue therapy

<sup>1</sup> mean adjusted for baseline value

\* p-value <0.0001

\*\* p-value <0.025

*Patients with renal impairment, 52 week placebo controlled data*

The efficacy and safety of empagliflozin as add on to antidiabetic therapy was evaluated in patients with renal impairment in a double-blind, placebo-controlled study for 52 weeks. Treatment with empagliflozin led to a statistically significant reduction of HbA1c (Table 7) and clinically meaningful improvement in FPG compared to placebo at Week 24. The improvement in HbA1c, body weight, and blood pressure was sustained up to 52 weeks.

Table 7 Results at 24 week in a placebo-controlled study of empagliflozin in renally impaired type 2 diabetes patients<sup>a</sup>

	Placebo	Empagliflozin 10 mg	Empagliflozin 25 mg	Placebo	Empagliflozin 25 mg
	eGFR 60 to <90 ml/min/1.73 m <sup>2</sup>			eGFR 45 to <60 ml/min/1.73 m <sup>2</sup>	
N	95	98	97	89	91
<b>HbA1c (%)</b>					
Baseline (mean)	8.09	8.02	7.96	8.08	8.12
Change from baseline <sup>1</sup>	0.06	-0.46	-0.63	-0.08	-0.54
Difference from placebo <sup>1</sup> (95% CI)		-0.52* (-0.72, -0.32)	-0.68* (-0.88, -0.49)		-0.46 (-0.66, -0.27)
N	89	94	91	84	86
<b>Patients (%) achieving HbA1c &lt;7% with baseline HbA1c 7%<sup>2</sup></b>					
	6.7	17.0	24.2	10.7	15.1
N	95	98	97	89	91
<b>Body Weight (kg)<sup>2</sup></b>					
Baseline (mean)	86.00	92.05	88.06	83.20	84.90
Change from baseline <sup>1</sup>	-0.33	-1.76	-2.33	-0.11	-1.39
Difference from placebo <sup>1</sup> (95% CI)		-1.43 (-2.09, -0.77)	-2.00 (-2.66, -1.34)		-1.28 (-2.08, -0.48)
N	95	98	97	89	91
<b>SBP (mmHg)<sup>2</sup></b>					
Baseline (mean)	134.69	137.37	133.68	137.29	135.04
Change from baseline <sup>1</sup>	0.65	-2.92	-4.47	0.37	-5.69
Difference from placebo <sup>1</sup> (95% CI)		-3.57 (-6.86, -0.29)	-5.12 (-8.41, -1.82)		-6.07 (-9.79, -2.34)

<sup>a</sup> Full analysis set (FAS) using last observation carried forward (LOCF) prior to glycaemic rescue therapy

<sup>1</sup> Mean adjusted for baseline value

<sup>2</sup> Not evaluated for statistical significance as a result of the sequential confirmatory testing procedure

\* p<0.0001

### Cardiovascular safety

In a prospective, pre-specified meta-analysis of independently adjudicated cardiovascular events from 12 phase 2 and 3 clinical studies involving 10,036 patients with type 2 diabetes, empagliflozin did not increase cardiovascular risk.

### Fasting plasma glucose

In four placebo-controlled studies, treatment with empagliflozin as monotherapy or add-on therapy to metformin, pioglitazone, or metformin plus a sulphonylurea resulted in mean changes from baseline in FPG of -20.5 mg/dl [-1.14 mmol/l] for empagliflozin 10 mg and -23.2 mg/dl [-1.29 mmol/l] for empagliflozin 25 mg compared to placebo (7.4 mg/dl [0.41 mmol/l]). This effect was observed after 24 weeks and maintained for 76 weeks.

### 2-hour post-prandial glucose

Treatment with empagliflozin as add-on to metformin or metformin and a sulphonylurea resulted in a clinically meaningful reduction of 2-hour post-prandial glucose (meal tolerance test) at 24 weeks (add-on to metformin: placebo +5.9 mg/dl, empagliflozin 10 mg: -46.0 mg/dl, empagliflozin 25 mg: -44.6 mg/dl, add-on to metformin and a sulphonylurea: placebo -2.3 mg/dl, empagliflozin 10 mg: -35.7 mg/dl, empagliflozin 25 mg: -36.6 mg/dl).

### Patients with high baseline HbA1c >10%

In a pre-specified pooled analysis of three phase 3 studies, treatment with open-label empagliflozin 25 mg in patients with severe hyperglycaemia (N=257, mean baseline HbA1c 11.26%) resulted in a clinically meaningful reduction in HbA1c from baseline of 3.27%; no placebo or empagliflozin 10 mg arms were included in these studies.

### Body weight

In a pre-specified pooled analysis of 4 placebo controlled studies, treatment with empagliflozin resulted in body weight reduction (-0.24 kg for placebo, -2.04 kg for empagliflozin 10 mg and -2.26 kg for empagliflozin 25 mg) at week 24 that was maintained up to week 52 (-0.16 kg for placebo, -1.96 kg for empagliflozin 10 mg and -2.25 kg for empagliflozin 25 mg).

### Blood pressure

The efficacy and safety of empagliflozin was evaluated in a double-blind, placebo controlled study of 12 weeks duration in patients with type 2 diabetes and high blood pressure on different antidiabetic and up to 2 antihypertensive therapies. Treatment with empagliflozin once daily resulted in statistically significant improvement in HbA1c, and 24 hour mean systolic and diastolic blood pressure as determined by ambulatory blood pressure monitoring (Table 8). Treatment with empagliflozin provided reductions in seated SBP and DBP.

Table 8 Efficacy results at 12 week in a placebo-controlled study of empagliflozin in patients with type 2 diabetes and uncontrolled blood pressure<sup>a</sup>

	Placebo	Jardiance	
		10 mg	25 mg
N	271	276	276
<b>HbA1c (%) at week 12<sup>1</sup></b>			
Baseline (mean)	7.90	7.87	7.92
Change from baseline <sup>2</sup>	0.03	-0.59	-0.62
Difference from placebo <sup>2</sup> (95% CI)		-0.62* (-0.72, -0.52)	-0.65* (-0.75, -0.55)
<b>24 hour SBP at week 12<sup>3</sup></b>			
Baseline (mean)	131.72	131.34	131.18

Change from baseline <sup>4</sup>	0.48	-2.95	-3.68
Difference from placebo <sup>4</sup> (95% CI)		-3.44* (-4.78, -2.09)	-4.16* (-5.50, -2.83)
<b>24 hour DBP at week 12<sup>3</sup></b>			
Baseline (mean)	75.16	75.13	74.64
Change from baseline <sup>5</sup>	0.32	-1.04	-1.40
Difference from placebo <sup>5</sup> (95% CI)		-1.36** (-2.15, -0.56)	-1.72* (-2.51, -0.93)

<sup>a</sup> Full analysis set (FAS)

<sup>1</sup> LOCF, values after taking antidiabetic rescue therapy censored

<sup>2</sup> Mean adjusted for baseline HbA1c, baseline eGFR, geographical region and number of antihypertensive medicinal products

<sup>3</sup> LOCF, values after taking antidiabetic rescue medications or changing antihypertensive rescue therapy censored

<sup>4</sup> Mean adjusted for baseline SBP, baseline HbA1c, baseline eGFR, geographical region and number of antihypertensive medicinal products

<sup>5</sup> Mean adjusted for baseline DBP, baseline HbA1c, baseline eGFR, geographical region and number of antihypertensive medicinal products

\* p-value <0.0001

\*\* p-value <0.001

In a pre-specified pooled analysis of 4 placebo-controlled studies, treatment with empagliflozin resulted in a reduction in systolic blood pressure (empagliflozin 10 mg: -3.9 mmHg; empagliflozin 25 mg: -4.3 mmHg) compared with placebo (-0.5 mmHg) and in diastolic blood pressure (empagliflozin 10 mg: -1.8 mmHg; empagliflozin 25 mg: -2.0 mmHg) compared with placebo (-0.5 mmHg) at week 24 that were maintained up to week 52.

### Paediatric population

The European Medicines Agency has deferred the obligation to submit the results of studies with Jardiance in one or more subsets of the paediatric population in type 2 diabetes mellitus (see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

### Absorption

The pharmacokinetics of empagliflozin have been extensively characterised in healthy volunteers and patients with type 2 diabetes. After oral administration, empagliflozin was rapidly absorbed with peak plasma concentrations occurring at a median  $t_{max}$  of 1.5 hours post-dose. Thereafter, plasma concentrations declined in a biphasic manner with a rapid distribution phase and a relatively slow terminal phase. The steady state mean plasma AUC and  $C_{max}$  were 1870 nmol.h and 259 nmol/l with empagliflozin 10 mg and 4740 nmol.h and 687 nmol/l with empagliflozin 25 mg once daily. Systemic exposure of empagliflozin increased in a dose-proportional manner. The single-dose and steady-state pharmacokinetic parameters of empagliflozin were similar suggesting linear pharmacokinetics with respect to time. There were no clinically relevant differences in empagliflozin pharmacokinetics between healthy volunteers and patients with type 2 diabetes.

Administration of empagliflozin 25 mg after intake of a high-fat and high calorie meal resulted in slightly lower exposure; AUC decreased by approximately 16% and  $C_{max}$  by approximately 37% compared to fasted condition. The observed effect of food on empagliflozin pharmacokinetics was not considered clinically relevant and empagliflozin may be administered with or without food.

### Distribution

The apparent steady-state volume of distribution was estimated to be 73.8 l based on the population pharmacokinetic analysis. Following administration of an oral [<sup>14</sup>C]-empagliflozin solution to healthy

volunteers, the red blood cell partitioning was approximately 37% and plasma protein binding was 86%.

### Biotransformation

No major metabolites of empagliflozin were detected in human plasma and the most abundant metabolites were three glucuronide conjugates (2-, 3-, and 6-O glucuronide). Systemic exposure of each metabolite was less than 10% of total drug-related material. *In vitro* studies suggested that the primary route of metabolism of empagliflozin in humans is glucuronidation by the uridine 5'-diphospho-glucuronosyltransferases UGT2B7, UGT1A3, UGT1A8, and UGT1A9.

### Elimination

Based on the population pharmacokinetic analysis, the apparent terminal elimination half-life of empagliflozin was estimated to be 12.4 hours and apparent oral clearance was 10.6 l/hour. The inter-subject and residual variabilities for empagliflozin oral clearance were 39.1% and 35.8%, respectively. With once-daily dosing, steady-state plasma concentrations of empagliflozin were reached by the fifth dose. Consistent with the half-life, up to 22% accumulation, with respect to plasma AUC, was observed at steady-state. Following administration of an oral [<sup>14</sup>C]-empagliflozin solution to healthy volunteers, approximately 96% of the drug-related radioactivity was eliminated in faeces (41%) or urine (54%). The majority of drug-related radioactivity recovered in faeces was unchanged parent drug and approximately half of drug related radioactivity excreted in urine was unchanged parent drug.

### Special populations

#### *Renal impairment*

In patients with mild, moderate or severe renal impairment (eGFR <30 - <90 ml/min/1.73 m<sup>2</sup>) and patients with kidney failure/end stage renal disease (ESRD), AUC of empagliflozin increased by approximately 18%, 20%, 66%, and 48%, respectively compared to subjects with normal renal function. Peak plasma levels of empagliflozin were similar in subjects with moderate renal impairment and kidney failure/ESRD compared to patients with normal renal function. Peak plasma levels of empagliflozin were roughly 20% higher in subjects with mild and severe renal impairment as compared to subjects with normal renal function. The population pharmacokinetic analysis showed that the apparent oral clearance of empagliflozin decreased with a decrease in eGFR leading to an increase in drug exposure.

#### *Hepatic impairment*

In subjects with mild, moderate, and severe hepatic impairment according to the Child-Pugh classification, AUC of empagliflozin increased approximately by 23%, 47%, and 75% and C<sub>max</sub> by approximately 4%, 23%, and 48%, respectively, compared to subjects with normal hepatic function.

#### *Body Mass Index*

Body mass index had no clinically relevant effect on the pharmacokinetics of empagliflozin based on the population pharmacokinetic analysis. In this analysis, AUC was estimated to be 5.82%, 10.4%, and 17.3% lower in subjects with BMI of 30, 35, and 45 kg/m<sup>2</sup>, respectively, compared to subjects with a body mass index of 25 kg/m<sup>2</sup>.

#### *Gender*

Gender had no clinically relevant effect on the pharmacokinetics of empagliflozin based on the population pharmacokinetic analysis.

#### *Race*

In the population pharmacokinetic analysis, AUC was estimated to be 13.5% higher in Asians with a body mass index of 25 kg/m<sup>2</sup> compared to non-Asians with a body mass index of 25 kg/m<sup>2</sup>.

### *Elderly patients*

Age did not have a clinically meaningful impact on the pharmacokinetics of empagliflozin based on the population pharmacokinetic analysis.

### *Paediatric patients*

Studies characterising the pharmacokinetics of empagliflozin in paediatric patients have not been performed.

## **5.3 Preclinical safety data**

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, genotoxicity, fertility and early embryonic development.

In long term toxicity studies in rodents and dogs, signs of toxicity were observed at exposures greater than or equal to 10-times the clinical dose of empagliflozin. Most toxicity was consistent with secondary pharmacology related to urinary glucose loss and electrolyte imbalances including decreased body weight and body fat, increased food consumption, diarrhea, dehydration, decreased serum glucose and increases in other serum parameters reflective of increased protein metabolism and gluconeogenesis, urinary changes such as polyuria and glucosuria, and microscopic changes including mineralisation in kidney and some soft and vascular tissues. Microscopic evidence of the effects of exaggerated pharmacology on the kidney observed in some species included tubular dilatation, and tubular and pelvic mineralisation at approximately 4-times the clinical AUC exposure of empagliflozin associated with the 25 mg dose.

Empagliflozin is not genotoxic.

In a 2 year carcinogenicity study, empagliflozin did not increase the incidence of tumors in female rats up to the highest dose of 700 mg/kg/day, which corresponds to approximately 72-times the maximal clinical AUC exposure to empagliflozin. In male rats, treatment-related benign vascular proliferative lesions (haemangiomas) of the mesenteric lymph node were observed at the highest dose, but not at 300 mg/kg/day, which corresponds to approximately 26-times the maximal clinical exposure to empagliflozin. Interstitial cell tumors in the testes were observed with a higher incidence in rats at 300 mg/kg/day and above, but not at 100 mg/kg/day which corresponds to approximately 18-times the maximal clinical exposure to empagliflozin. Both tumors are common in rats and are unlikely to be relevant to humans.

Empagliflozin did not increase the incidence of tumors in female mice at doses up to 1000 mg/kg/day, which corresponds to approximately 62-times the maximal clinical exposure to empagliflozin. Empagliflozin induced renal tumors in male mice at 1000 mg/kg/day, but not at 300 mg/kg/day, which corresponds to approximately 11-times the maximal clinical exposure to empagliflozin. The mode of action for these tumors is dependent on the natural predisposition of the male mouse to renal pathology and a metabolic pathway not reflective of humans. The male mouse renal tumors are considered not relevant to humans.

At exposures sufficiently in excess of exposure in humans after therapeutic doses, empagliflozin had no adverse effects on fertility or early embryonic development. Empagliflozin administered during the period of organogenesis was not teratogenic. Only at maternally toxic doses, empagliflozin also caused bent limb bones in the rat and increased embryofetal loss in the rabbit.

In pre- and postnatal toxicity studies in rats, reduced weight gain of offspring was observed at maternal exposures approximately 4-times the maximal clinical exposure to empagliflozin. No such effect was seen at systemic exposure equal to the maximal clinical exposure to empagliflozin. The relevance of this finding to humans is unclear.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Lactose monohydrate  
Microcrystalline cellulose  
Hydroxypropylcellulose  
Croscarmellose sodium  
Colloidal anhydrous silica  
Magnesium stearate

#### Film coating

Hypromellose  
Titanium dioxide (E171)  
Talc  
Macrogol (400)  
Iron oxide yellow (E172)

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

3 years

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

PVC/aluminium perforated unit dose blisters.  
Pack sizes of 7 x 1, 10 x 1, 14 x 1, 28 x 1, 30 x 1, 60 x 1, 70 x 1, 90 x 1, and 100 x 1 film-coated tablets.

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim International GmbH  
Binger Str. 173  
D-55216 Ingelheim am Rhein  
Germany

## **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/14/930/001  
EU/1/14/930/002



EU/1/14/930/003  
EU/1/14/930/004  
EU/1/14/930/005  
EU/1/14/930/006  
EU/1/14/930/007  
EU/1/14/930/008  
EU/1/14/930/009

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

## **ANNEX II**

- A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer responsible for batch release

Boehringer Ingelheim Pharma GmbH & Co. KG  
Binger Strasse 173  
55216 Ingelheim am Rhein  
Germany

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to medical prescription.

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

### **• Periodic Safety Update Reports**

The marketing authorisation holder shall submit the first periodic safety update report for this product within 6 months following authorisation. Subsequently, the marketing authorisation holder shall submit periodic safety update reports for this product in accordance with the requirements set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

### **• Risk Management Plan (RMP)**

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the Marketing Authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

If the dates for submission of a PSUR and the update of a RMP coincide, they can be submitted at the same time.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Jardiance 10 mg film-coated tablets  
Empagliflozin

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each tablet contains 10 mg of empagliflozin.

**3. LIST OF EXCIPIENTS**

Contains lactose, see leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

7 x 1 film-coated tablets  
10 x 1 film-coated tablets  
14 x 1 film-coated tablets  
28 x 1 film-coated tablets  
30 x 1 film-coated tablets  
60 x 1 film-coated tablets  
70 x 1 film-coated tablets  
90 x 1 film-coated tablets  
100 x 1 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
Oral use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim International GmbH  
D-55216 Ingelheim am Rhein  
Germany

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/14/930/010 7 tablets  
EU/1/14/930/011 10 tablets  
EU/1/14/930/012 14 tablets  
EU/1/14/930/013 28 tablets  
EU/1/14/930/014 30 tablets  
EU/1/14/930/015 60 tablets  
EU/1/14/930/016 70 tablets  
EU/1/14/930/017 90 tablets  
EU/1/14/930/018 100 tablets

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Jardiance 10 mg

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**Blisters (perforated)**

**1. NAME OF THE MEDICINAL PRODUCT**

Jardiance 10 mg tablets  
Empagliflozin

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**



**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Jardiance 25 mg film-coated tablets  
Empagliflozin

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each tablet contains 25 mg of empagliflozin.

**3. LIST OF EXCIPIENTS**

Contains lactose, see leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

7 x 1 film-coated tablets  
10 x 1 film-coated tablets  
14 x 1 film-coated tablets  
28 x 1 film-coated tablets  
30 x 1 film-coated tablets  
60 x 1 film-coated tablets  
70 x 1 film-coated tablets  
90 x 1 film-coated tablets  
100 x 1 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
Oral use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim International GmbH  
D-55216 Ingelheim am Rhein  
Germany

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/14/930/001 7 tablets  
EU/1/14/930/002 10 tablets  
EU/1/14/930/003 14 tablets  
EU/1/14/930/004 28 tablets  
EU/1/14/930/005 30 tablets  
EU/1/14/930/006 60 tablets  
EU/1/14/930/007 70 tablets  
EU/1/14/930/008 90 tablets  
EU/1/14/930/009 100 tablets

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Jardiance 25 mg

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**Blisters (perforated)**

**1. NAME OF THE MEDICINAL PRODUCT**

Jardiance 25 mg tablets  
Empagliflozin

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

**B. PACKAGE LEAFLET**

## Package leaflet: Information for the patient

### Jardiance 10 mg film-coated tablets

### Jardiance 25 mg film-coated tablets

empagliflozin

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

#### **Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Jardiance is and what it is used for
2. What you need to know before you take Jardiance
3. How to take Jardiance
4. Possible side effects
5. How to store Jardiance
6. Contents of the pack and other information

#### **1. What Jardiance is and what it is used for**

Jardiance contains the active substance empagliflozin which works by blocking a protein in the kidneys called sodium-glucose co-transporter 2 (SGLT2). SGLT2 prevents glucose from being excreted in urine by absorbing glucose into the bloodstream as blood is being filtered in the kidneys. By blocking this protein, the medicine causes glucose (blood sugar), to be removed via the urine and blood glucose levels, which are too high because of your type 2 diabetes, are thereby reduced.

Your doctor has prescribed this medicine to help lower your blood sugar.

- Jardiance is used to treat type 2 diabetes in adult patients (aged 18 years and older) that cannot be controlled by diet and exercise alone.
- Jardiance can be used without other medicines in patients who cannot take metformin (another diabetes medicine).
- Jardiance can also be used with other medicines. These may be medicines taken by mouth or insulin given by injection.

It is important that you continue with your diet and exercise plan as told by your doctor, pharmacist or nurse.

#### **What is type 2 diabetes?**

Type 2 diabetes is a disease that comes from both your genes and your lifestyle. If you have type 2 diabetes, your pancreas does not make enough insulin to control the level of glucose in your blood, and your body is unable to use its own insulin effectively. This results in high levels of glucose in your blood which can lead to medical problems like heart disease, kidney disease, blindness, and poor circulation in your limbs.

## **2. What you need to know before you take Jardiance**

### **Do not take Jardiance:**

- if you are allergic to empagliflozin or any of the other ingredients of this medicine (listed in section 6).

### **Warnings and precautions**

Talk to your doctor, pharmacist or nurse before taking Jardiance:

- if you have “type 1 diabetes”. This type usually starts when you are young and your body does not produce any insulin.
- if you have increased levels of “ketone bodies” in your urine or blood, seen in laboratory tests. This is a sign of “diabetic ketoacidosis” – a problem you can get with diabetes whose signs include rapid weight loss, feeling sick or being sick, a sweet smell to your breath, a sweet or metallic taste in your mouth or a strange odour to your urine or sweat.
- if you have serious kidney problems – your doctor may ask you to take a different medicine.
- if you are 75 years old or older, as increased passing of urine due to the medicine may affect fluid balance in your body and increase your risk of dehydration. Possible signs are listed in section 4, ‘Possible side effects’ under ‘dehydration’.
- if you are 85 years old or older as you should not start taking Jardiance.
- if you are being sick, have diarrhoea or fever, or if you are not able to eat or drink. These conditions can cause dehydration. Your doctor may ask you to stop taking Jardiance until you recover to prevent loss of too much body fluid.
- if you have a serious infection of the kidney or the urinary tract with fever. Your doctor may ask you to stop taking Jardiance until you have recovered.

### **Urine glucose**

Because of how this medicine works, your urine will test positive for sugar while you are taking this medicine.

### **Children and adolescents**

Jardiance is not recommended for children and adolescents under 18 years, because it has not been studied in these patients.

### **Other medicines and Jardiance**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

It is important to tell your doctor:

- if you are taking a medicine used to remove water from the body (diuretic). Your doctor may ask you to stop taking Jardiance. Possible signs of losing too much fluid from your body are listed in section 4 ‘Possible side effects’.
- if you are taking other medicines that lower the amount of sugar in your blood such as insulin or a “sulphonylurea” medicine. Your doctor may want to lower the dose of these other medicines, to prevent your blood sugar levels from getting too low (hypoglycaemia).

### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. Do not use Jardiance if you are pregnant. It is unknown if Jardiance is harmful to the unborn child. Do not use Jardiance if you are breast-feeding. It is not known if Jardiance passes into human breast milk.

### **Driving and using machines**

Jardiance has minor influence on the ability to drive and use machines.

Taking this medicine in combination with medicines called sulphonylureas or with insulin can cause blood sugar levels to drop too low (hypoglycaemia), which may cause symptoms such as shaking, sweating and change in vision, and may affect your ability to drive and use machines. Do not drive or use any tools or machines, if you feel dizzy while taking Jardiance.

**Jardiance contains lactose**

Jardiance contains lactose (milk sugar). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

**3. How to take Jardiance**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

**How much to take**

- The starting dose of Jardiance is one 10 mg tablet once a day. Your doctor will decide whether to increase your dose to 25 mg once a day.
- Your doctor may limit your dose to 10 mg once a day if you have a kidney problem.
- Your doctor will prescribe the strength that is right for you. Do not change your dose unless your doctor has told you to.

**Taking this medicine**

- Swallow the tablet whole with water
- You can take the tablet with or without food
- You can take the tablet at any time of the day. However, try to take it at the same time each day. This will help you to remember to take it.

Your doctor may prescribe Jardiance together with another diabetes medicine. Remember to take all medicines as directed by your doctor to achieve the best results for your health.

Diet and exercise can help your body use its blood sugar better. It is important to stay on the diet and exercise program recommended by your doctor while taking Jardiance.

**If you take more Jardiance than you should**

If you take more Jardiance than you should, talk to a doctor immediately or go to a hospital immediately. Take the medicine pack with you.

**If you forget to take Jardiance**

What to do if you forget to take a tablet depends on how long it is until your next dose.

- If it is 12 hours or more until your next dose, take Jardiance as soon as you remember. Then take your next dose at the usual time.
- If it is less than 12 hours until your next dose, skip the missed dose. Then take your next dose at the usual time.
- Do not take a double dose of Jardiance to make up for a forgotten dose.

**If you stop taking Jardiance**

Do not stop taking Jardiance without first consulting your doctor. Your blood sugar levels may increase when you stop taking Jardiance.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

**4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Contact your doctor as soon as possible if you notice the following side effects:**

**Low blood sugar (hypoglycemia), seen very commonly (may affect more than 1 in 10 people)**

If you take Jardiance with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin, your risk of getting low blood sugar is higher. The signs of low blood sugar may include:

- shaking, sweating, feeling very anxious or confused, fast heart beat
- excessive hunger, headache

Your doctor will tell you how to treat low blood sugar levels and what to do if you get any of the signs above. If you have symptoms of low blood sugar, eat glucose tablets, a high sugar snack or drink fruit juice. Measure your blood sugar if possible and rest.

**Urinary tract infection, seen commonly (may affect up to 1 in 10 people)**

The signs of urinary tract infection are:

- burning sensation when passing urine
- urine that appears cloudy
- pain in the pelvis, or mid-back pain (when kidneys are infected)

An urge to pass urine or more frequent urination may be due to the way Jardiance works, but as they can also be signs of urinary tract infection, if you note an increase in such symptoms, you should also contact your doctor.

**Dehydration, seen uncommonly (may affect up to 1 in 100 people)**

The signs of dehydration are not specific, but may include:

- unusual thirst
- lightheadedness or dizziness upon standing
- fainting or loss of consciousness

**Other side effects while taking Jardiance:**

Common

- thrush or genital yeast infection
- passing more urine than usual or needing to pass urine more often
- itching

Uncommon

- straining or pain when emptying the bladder

**Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Jardiance**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister and the carton after 'EXP'. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not use this medicine if you notice that the packaging is damaged or shows signs of tampering.



Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Jardiance contains**

- The active substance is empagliflozin.
  - Each tablet contains 10 mg or 25 mg empagliflozin.
- The other ingredients are:
  - tablet core: lactose monohydrate (see end of section 2 under ‘Jardiance contains lactose’), cellulose microcrystalline, hydroxypropylcellulose, croscarmellose sodium, colloidal anhydrous silica, magnesium stearate
  - film-coating: hypromellose, titanium dioxide (E171), talc, macrogol (400), iron oxide yellow (E172)

### **What Jardiance looks like and contents of the pack**

Jardiance 10 mg film-coated tablets are round, pale yellow, biconvex and bevel-edged. They have “S10” on one side and the Boehringer Ingelheim logo on the other side. The tablets are 9.1 mm in diameter.

Jardiance 25 mg film-coated tablets are oval, pale yellow and biconvex. They have “S25” on one side and the Boehringer Ingelheim logo on the other side. The tablet is 11.1 mm long and has a width of 5.6 mm.

Jardiance tablets are available in PVC/aluminium perforated unit dose blisters. The pack sizes are 7 x 1, 10 x 1, 14 x 1, 28 x 1, 30 x 1, 60 x 1, 70 x 1, 90 x 1, and 100 x 1 film-coated tablets.

Not all pack sizes may be marketed in your country.

### **Marketing Authorisation Holder**

Boehringer Ingelheim International GmbH  
Binger Strasse 173  
55216 Ingelheim am Rhein  
Germany

### **Manufacturer**

Boehringer Ingelheim Pharma GmbH & Co. KG  
Binger Strasse 173  
55216 Ingelheim am Rhein  
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

SCS Boehringer Ingelheim Comm.V  
Tél/Tel: +32 2 773 33 11

Eli Lilly Benelux S.A./N.V.  
Tél/Tel: +32 2 548 84 84

: +359 2 958 79 98

" " . . -  
. +359 2 491 41 40

**eská republika**

Boehringer Ingelheim spol. s r.o.  
Tel: +420 234 655 111

ELI LILLY R, s.r.o.  
Tel: +420 234 664 111

**Danmark**

Boehringer Ingelheim Danmark A/S  
Tlf: +45 39 15 88 88

Eli Lilly Danmark A/S  
Tlf: +45 45 26 60 00

**Deutschland**

Boehringer Ingelheim Pharma GmbH & Co. KG  
Tel: +49 (0) 800 77 90 900

Lilly Deutschland GmbH  
Tel. +49 (0) 6172 273 2222

**Eesti**

Boehringer Ingelheim RCV GmbH & Co KG  
Eesti filiaal  
Tel: +372 612 8000

Eli Lilly Holdings Limited Eesti filiaal  
Tel: +372 68 17 280

Boehringer Ingelheim Ellas A.E.  
T : +30 2 10 89 06 300

- . . . .  
: +30 2 10 62 94 600

**Lietuva**

Boehringer Ingelheim RCV GmbH & Co KG  
Lietuvos filialas  
Tel.: +370 37 473 922

Eli Lilly Holdings Limited atstovyb  
Tel. +370 52 649 600

**Luxembourg/Luxemburg**

- SCS Boehringer Ingelheim Comm.V  
Tél/Tel: +32 2 773 33 11

Eli Lilly Benelux S.A./N.V.  
Tél/Tel: +32 2 548 84 84

**Magyarország**

Boehringer Ingelheim RCV GmbH & Co KG  
Magyarországi Fióktelepe  
Tel.: +36 1 299 89 00

Lilly Hungária Kft.  
Tel: +36 1 328 5100

**Malta**

Boehringer Ingelheim Ltd.  
Tel: +44 1344 424 600

Charles de Giorgio Ltd.  
Tel: +356 2 560 05 00

**Nederland**

Boehringer Ingelheim b.v.  
Tel: +31 (0) 800 22 55 889

Eli Lilly Nederland B.V.  
Tel: +31 (0) 306 02 58 00

**Norge**

Boehringer Ingelheim Norway KS  
Tlf: +47 66 76 13 00

Eli Lilly Norge A.S.  
Tlf: +47 22 88 18 00

**Österreich**

Boehringer Ingelheim RCV GmbH & Co KG  
Tel: +43 1 80 105-0

Eli Lilly Ges.m.b.H.  
Tel: +43 1 71 1780

**España**

Boehringer Ingelheim España S.A.  
Tel: +34 93 404 51 00

Lilly S.A.

Tel: +34 91 663 50 00

**France**

Boehringer Ingelheim France S.A.S.  
Tél: +33 3 26 50 45 33

Lilly France SAS

Tél: +33 1 55 49 34 34

**Hrvatska**

Boehringer Ingelheim Zagreb d.o.o.  
Tel: +385 1 2444 600

Eli Lilly Hrvatska d.o.o.

Tel: +385 1 2350 999

**Ireland**

Boehringer Ingelheim Ireland Ltd.  
Tel: +353 1 295 9620

Eli Lilly and Company (Ireland) Limited

Tel: +353 1 661 4377

**Ísland**

Vistor hf.

Sími: +354 535 7000

Icepharma hf.

Sími + 354 540 8000

**Italia**

Boehringer Ingelheim Italia S.p.A.  
Tel: +39 02 5355 1

Eli Lilly Italia S.p.A.

Tel: +39 055 42571

Boehringer Ingelheim Ellas A.E.

T : +30 2 10 89 06 300

Phadisco Ltd

: +357 2 271 5000

**Polska**

Boehringer Ingelheim Sp.zo.o.  
Tel.: +48 22 699 0 699

Eli Lilly Polska Sp. z o.o.

Tel: +48 22 440 3 300

**Portugal**

Boehringer Ingelheim, Lda.  
Tel: +351 21 313 53 00

Lilly Portugal Produtos Farmacêuticos, Lda

Tel: +351 21 412 66 00

**România**

Boehringer Ingelheim RCV GmbH & Co KG  
Viena - Sucursala Bucure ti  
Tel: +40 21 302 28 00

Eli Lilly România S.R.L.

Tel: +40 21 402 30 00

**Slovenija**

Boehringer Ingelheim RCV GmbH & Co KG  
Podružnica Ljubljana  
Tel: +386 1 586 40 00

Eli Lilly farmacevtska družba, d.o.o.

Tel: +386 1 580 00 10

**Slovenská republika**

Boehringer Ingelheim RCV GmbH & Co KG  
organizacia ná zložka  
Tel: +421 2 5810 1211

Eli Lilly Slovakia, s.r.o.

Tel: +421 2 2066 3111

**Suomi/Finland**

Boehringer Ingelheim Finland Ky  
Puh/Tel: +358 10 3102 800

Oy Eli Lilly Finland Ab

Puh/Tel: +358 98 5452 50

**Sverige**

Boehringer Ingelheim AB  
Tel: +46 8 721 21 00

Eli Lilly Sweden AB

Tel: +46 8 737 88 00

**Latvija**

Boehringer Ingelheim RCV GmbH & Co KG  
Latvijas filiāle  
Tel: +371 67 240 011

**United Kingdom**

Boehringer Ingelheim Ltd.  
Tel: +44 1344 424 600

Eli Lilly Holdings Limited pārstāvis Latvijā  
Tel: +371 67 364 000

Eli Lilly and Company Limited  
Tel: +44 1256 315 000

**This leaflet was last revised in {MM/YYYY}.**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>.