

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

1. NAME OF THE MEDICINAL PRODUCT

Defitelio 80 mg/mL concentrate for solution for infusion

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One mL contains defibrotide* 80mg corresponding to a quantity of 200 mg in 2.5 mL in a vial and corresponding to a concentration in the range of 4 mg/mL to 20 mg/mL after dilution

* produced from porcine intestinal mucosa.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Concentrate for solution for infusion (sterile concentrate).

The solution is clear light yellow to brown, free from particulate matter or turbidity.

4. CLINICAL PARTICULARS

4.1 Therapeutic indication

Defitelio is indicated for the treatment of severe hepatic veno-occlusive disease (VOD) also known as sinusoidal obstructive syndrome (SOS) in haematopoietic stem-cell transplantation (HSCT) therapy.

It is indicated in adults and in adolescents, children and infants over 1 month of age .

4.2 Posology and method of administration

Defitelio must be prescribed and administered to patients by specialised physicians experienced in the diagnosis and treatment of complications of HSCT.

Posology

The recommended dose is 6.25 mg/kg body weight every 6 hours (25 mg/kg/day).

There is limited efficacy and safety data on doses above this level and consequently it is not recommended to increase the dose above 25 mg/kg/day.

Defitelio should be administered for a minimum of 21 days and continued until the symptoms and signs of severe VOD resolve.

Renal and Hepatic Impairment

No formal pharmacokinetic studies have been performed in patients with renal or hepatic impairment, however, the medicinal product has been used in clinical trials in patients developing renal and hepatic impairment without dose adjustment with no safety issues identified. No dose adjustment is therefore recommended but careful monitoring of patients should be undertaken (see section 5.2).

Paediatric population

The recommended dose for children aged 1 month to 18 years is the same mg/kg dose as for adults i.e. 6.25 mg/kg body weight every 6 hours.

Method of administration

Defitelio is administered by intravenous infusion, over two hours.

Defitelio should always be diluted prior to use. Defitelio can be diluted with 5% glucose solution for infusion or sodium chloride 9 mg/mL (0.9%) solution for infusion, to a suitable concentration to permit infusion over 2 hours. The total volume of infusion should be determined based on the individual's patient weight. The final concentration of Defitelio should be in the range of 4 mg/mL to 20 mg/mL.

Vials are intended for a single use and unused solution from a single dose must be discarded (see section 6.6)

For detailed instructions on dilution of the medicinal product before administration, see section 6.6.

4.3 Contraindications

- Hypersensitivity to defibrotide or to any of the excipients listed in section 6.1
- Concomitant use of thrombolytic therapy (e.g. t-PA) (see section 4.5).

4.4 Special warnings and precautions for use

Use of medicinal products that increase the risk of haemorrhage within 24 hours of Defitelio administration (within 12 hours in the case of unfractionated heparin) is not recommended.

Concomitant systemic anticoagulant therapy (e.g. heparin, warfarin, direct thrombin inhibitors and direct factor Xa inhibitors) (see section 4.5), except for routine maintenance or reopening of central venous line, requires careful monitoring. Consideration should be given to discontinuation of Defitelio during use of such therapy.

Medicinal products that affect platelet aggregation (e.g. non-steroidal anti-inflammatory agents) should be administered with care, under close medical supervision, during Defitelio administration.

In patients who have or develop clinically significant acute bleeding requiring blood transfusion, Defitelio is not recommended or should be discontinued. Temporary discontinuation of Defitelio is recommended in patients who undergo surgery or invasive procedures at significant risk of major bleeding.

Administration of Defitelio to patients who have haemodynamic instability, defined as inability to maintain mean arterial pressure with single pressor support, is not recommended.

The safety and efficacy of Defitelio in children aged less than 1 month has not yet been established. No data are available. The use of Defitelio in children aged less than one month is not recommended.

A bolus administration of Defitelio may cause flushing or a sensation of "generalised heat".

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially "sodium-free".

4.5 Interaction with other medicinal products and other forms of interaction

Potential Interactions with recombinant t-PA

In a mouse model of thromboembolism, recombinant t-PA potentiated the antithrombotic effect of defibrotide when given intravenously and thus co-administration may present an increased risk of haemorrhage and is contraindicated (see section 4.3).

Potential Interactions with antithrombotic fibrinolytic agents

Defibrotide has a profibrinolytic effect (see section 5.1) and this may potentially enhance the activity of antithrombotic/fibrinolytic medicinal products.

There is currently no reported experience in patients on the concomitant treatment with Low Molecular Weight Heparins (LMWHs), warfarin or the concomitant treatment with direct thrombin inhibitors (e.g., dabigatran) or direct Factor Xa inhibitors (e.g., rivaroxaban and apixaban). Therefore, the use of defibrotide with antithrombotic/fibrinolytic medicinal products is not recommended. However, if used, in exceptional cases, caution should be exercised by closely monitoring the coagulation parameters (see section 4.4).

Potential Interactions with other medicinal products

Defitelio does not inhibit or induce CYP450s (see section 5.2).

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no studies using defibrotide in pregnant women. Embryo-foetal developmental toxicology studies in pregnant rats and rabbits of defibrotide doses close to the recommended therapeutic human dose, revealed a high rate of haemorrhagic abortion (see section 5.3).

Defitelio should not be used during pregnancy unless the clinical condition of the woman requires treatment with Defitelio.

Contraception in males and females

Effective contraception is required for patients and partners of patients during exposure to Defitelio and for one week subsequent to discontinuation.

Breast-feeding

It is not known whether defibrotide is excreted in human milk. Considering the nature of the product, a risk to the newborns/infants is not expected. Defitelio may be used during breastfeeding.

Fertility

There are no studies investigating the effects of defibrotide on human fertility.

4.7 Effects on ability to drive and use machines

Defitelio is expected to have no or negligible influence on the ability to drive and operate machinery. However, patients would not be expected to drive or operate machinery due to the nature of the underlying disease.

4.8 Undesirable effects

Summary of the Safety Profile

In the Phase 3 pivotal treatment study (2005-01 Study), the overall incidence of adverse events was similar in the defibrotide treatment group and in the control group (historical).

Any events reported as possibly related on at least two occasions have been defined as ADRs and included in the table below.

The most frequent adverse reactions observed during the treatment of hepatic VOD in pre-marketing use are haemorrhage (including but not limited to gastrointestinal haemorrhage, pulmonary haemorrhage and epistaxis), hypotension and coagulopathy.

In addition, although in the defibrotide studies in VOD there have been no reports of hypersensitivity, cases of hypersensitivity including anaphylaxis were reported from a previously marketed formulation of defibrotide, consequently hypersensitivity is included as an ADR

Tabulated list of adverse reactions

Adverse reactions observed are listed below, by system organ class and frequency. Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness. Frequencies are defined as: very common ($\geq 1/10$), common ($> 1/100$ to $< 1/10$), uncommon ($> 1/1,000$ to $< 1/100$), rare ($> 1/10,000$ to $< 1/1,000$), very rare ($< 1/10,000$).

<i>Blood and lymphatic system disorders</i>	
Common	Coagulopathy
<i>Immune system disorders</i>	
Uncommon	Hypersensitivity
	Anaphylactic reaction
<i>Nervous system disorders</i>	
Common	Cerebral haemorrhage
Uncommon	Cerebral haematoma
<i>Eye disorders</i>	
Uncommon	Conjunctival haemorrhage
<i>Vascular disorders</i>	
Common	Hypotension
	Haemorrhage
<i>Respiratory, thoracic and mediastinal disorders</i>	
Common	Pulmonary haemorrhage
	Epistaxis
Uncommon	Haemothorax
<i>Gastrointestinal disorders:</i>	
Common	Gastrointestinal haemorrhage
	Vomiting
Uncommon	Haematemesis
	Melaena
	Mouth haemorrhage
	Diarrhoea
	Nausea
<i>Skin and subcutaneous tissue disorders</i>	
Uncommon	Ecchymosis
	Petechiae
	Rash
	Pruritus
<i>Renal and urinary disorders</i>	
Common	Haematuria
<i>General disorders and administration site conditions</i>	
Common	Catheter site haemorrhage

Uncommon	Injection site haemorrhage
	Pyrexia

Paediatric population

In the treatment studies over 50% of the patients were children. In doses above the recommended dose of 25 mg/kg/day there was a higher proportion of patients with bleeding events in the high dose group but since many events occurred in the follow-up period, a clear relationship with defibrotide treatment could not be determined. In the paediatric prevention study at 25 mg/kg/day there was an increased incidence of any bleeding events in the defibrotide group compared with the treatment group. However there was no difference in incidence of serious bleeding or bleeding events with fatal outcome.

The frequency nature and severity of adverse reactions in children are otherwise the same as in adults. No special precautions are indicated.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorization of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions **via the national reporting system listed in [Appendix V](#)**.

4.9 Overdose

There is no specific antidote for overdose and treatment should be symptomatic.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: **not yet assigned**; ATC code: **not yet assigned**.

Mechanism of action

In vitro, defibrotide has been shown to bind to various sites on vascular endothelium that are involved in cell regulation, providing a stimulus that promotes protection of activated endothelial cells. Defibrotide has also been shown to protect endothelial cells from fludarabine-mediated apoptosis, while not impacting its anti-leukemic effect. Defibrotide also inhibits the expression of heparanase contributing to extracellular matrix integrity and thereby tissue homeostasis. It is postulated that these actions protect endothelial cells.

Also, *in vitro*, defibrotide has been shown to increase tissue-type plasminogen activator (t-PA) function and decrease plasminogen activator inhibitor-1 (PAI-1) activity resulting in a decrease in procoagulant activity and an increase in the fibrinolytic potential of endothelial cells. Defibrotide also has been shown to have a weak profibrinolytic activity *in vitro*.

The pathophysiology of VOD is multifactorial and complex. Both endothelial cell damage and prothrombotic-hypofibrinolytic state are critical factors in the pathophysiology of this disease. Whilst the mechanism of action of defibrotide has not been fully elucidated, *in vitro* data support a role for defibrotide in both endothelial cell protection and the restoration of the thrombo-fibrinolytic balance. However no pharmacodynamics effects from defibrotide have been identified *in vivo*.

Clinical efficacy and safety

The efficacy and safety of Defitelio in the treatment of severe VOD were studied in a pivotal Phase 3 historical-controlled study (2005-01). Forty-four children and 58 adult patients with severe VOD

post-HSCT, were treated with Defitelio 25 mg/kg/day intravenous by infusion, and compared with 32 historical control patients. Median length of therapy in those treated with Defitelio was 22 days. A significantly higher proportion of patients in the Defitelio treated group achieved a complete response defined as total bilirubin less than 2 mg/dL and resolution of MOF (multiple organ failure); Day+100 complete response was 23.5% (24/102) with Defitelio versus 9.4% (3/32) in the historical control (p=0.013). In addition, Day+100 survival rate was improved in the Defitelio group with 38.2% (39/102) of the patients surviving versus 25.0% (8/32) in the historical control group (p=0.034).

The efficacy data from this pivotal study are supported and confirmed with data from a dose-finding study (25 mg/kg arm) and the interim analysis of an ongoing Treatment IND study (severe VOD subset), as presented in Table 1 and 2.

Table 1: Treatment Study Results: Complete Response of Severe VOD at Day+100

	Individual Studies			
	Dose-Finding (25mg/kg/day arm)	Open Label Treatment IND (25mg/kg/day)	Historically Controlled Trial (25mg/kg/day)	
			Defibrotide treated group	Historical Control
Complete Response by Day+100	43% (32/75)	25.9% (57/220)	23.5% (24/102)	9.4% (3/32)
p= 0.0131				

Table 2: Treatment Study Results: Day+100 Survival

	Individual Studies			
	Dose-Finding (25mg/kg/day arm)	Open Label Treatment IND (25mg/kg/day)	Historically Controlled Trial (25mg/kg/day)	
			Defibrotide treated group	Historical Control
Survival by Day+100	43.9%*	44.8%*	38.2%*	25.0%*
p=0.0341				

*=Kaplan Meier estimates for time-to-event analysis by Day100

Outcome data available from 611 patients treated with Defitelio on a compassionate use basis for non-severe and severe VOD post-transplant, are consistent with the controlled clinical studies, with complete response rate 24% (51/212) and survival 37% (78/212) in the subset of patients with severe VOD.

A controlled randomised prophylaxis study (Study 2004-000592-33) was conducted in the paediatric patients undergoing HSCT. Patients (n=356) were randomised to receive 25 mg/kg/day from the start of conditioning or were randomised to receive no prophylaxis.

A 40% reduction in the overall incidence of VOD in the Defitelio prophylaxis arm (from 19.9% in the control arm to 12.2% in the Defitelio arm), has been shown. The use of Defitelio rescue treatment for all patients who developed VOD meant that the study was not designed to assess any survival advantage and none was seen in this study.

In secondary analyses on the subset of patients undergoing allogeneic transplants, Defitelio prophylaxis was also associated with a lower incidence and less Grade 2 to 4 severity of acute graft versus host disease (aGvHD) by Day+100.

Coppell et al in 2010 reported data from a large meta-analysis of 235 patients with severe VOD showing a background mortality rate of severe VOD of 84.3% and that this mortality rate has remained constant over several decades.

Data derived from an independent US registry have shown a beneficial effect of Defitelio in routine clinical practice. At an interim analysis of the on-going registry, data from 96 patients with severe VOD were available.

The Day+100 all-cause mortality in patients with severe VOD who were not treated with defibrotide was 69%, and 61% in those patients who received defibrotide. These data are from an open label registry and the subjects were not randomised.

Additional information is shown in the following Table 3

Table 3: US Registry data

	Non-defibrotide treated	Defibrotide treated
	55	41
Alive at Day +100	17 (31%)	16 (39%)
VOD resolved by Day +100	16 (29%)	21 (51%)

Paediatric population

In each of the clinical studies performed in the treatment of VOD, over 50% of patients were under the age of 18 years. Safety information in children are available from the prevention study conducted solely in children. Safety and efficacy in children aged less than 1 month have not yet been established.

Cardiac electrophysiology

Based on the results of the QTc study, conducted in healthy subjects at therapeutic and supra-therapeutic doses, it can be concluded that Defitelio has no significant or clinically relevant QTc-prolonging potential at doses up to 4 times higher than therapeutically indicated. Defitelio might be considered free of proarrhythmic toxicity related to QT changes.

This medicinal product has been authorised under ‘exceptional circumstances’. This means that due to the rarity of the disease and for ethical reasons preventing to perform a placebo-controlled study, it has not been possible to obtain complete information on this medicinal product.

The European Medicines Agency will review any new information which may become available every year and this SmPC will be updated as necessary

5.2 Pharmacokinetic properties

Absorption and Distribution

In healthy volunteers, after a single 6.25 mg/kg dose of Defitelio given as a 2-hour infusion, the pharmacokinetic parameters were as follows:

Table 4. Defitelio pharmacokinetic parameters after intravenous infusion of 6.25 mg/kg to healthy subjects.

Parameter	Defitelio PK Parameters Mean ± SD
C _{max} (µg/mL)	17.3 ± 3.83
t _{max} (h)#	2.00 (1.00-2.00)
AUC _t (µg/mL*h)	26.9 ± 8.53
AUC (µg/mL*h)	48.1 ± 6.49
V _d (mL)	9934 ± 3807
CL (L/h)	10.4 ± 1.77
Kel (1/h)	1.25 ± 0.66
t _{1/2} (h)	0.71 ± 0.35

median (min-max)

Maximum plasma concentrations peaked at the end of the infusion period and declined thereafter with a rapid clearance and most of samples were undetectable 3.5 hours after the start of the infusion. Pharmacokinetic modelling simulation analysis showed that Defitelio plasma concentrations do not accumulate upon multiple doses administration and with doses up to 4-fold the therapeutic dose. Volume of distribution is around 10 L and Defitelio is not bound to plasma proteins.

Elimination

98% of Defitelio is excreted unchanged in the urine in the first 4 hours after the start of the infusion. The remaining 2% is excreted within 24 hours.

Metabolism

Defitelio does not inhibit or induce CYP450s.

Special Populations

No formal pharmacokinetic studies have been performed in special populations. Defitelio has been used in clinical trials in patients developing renal and hepatic impairment without dose adjustment with no major safety issues identified (see section 4.2).

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity or carcinogenicity.

In both species, the main findings were accumulation of vacuolated macrophages in liver of dogs and in liver, kidneys and lymph nodes of rats. Macrophages are considered the main target organ.

Embryo-foetal development

In the Segment II reproductive studies in rats and rabbits, defibrotide has shown maternal toxicity by inducing a high rate of haemorrhagic abortion when infused intravenously over two hours at all dose levels tested including doses close to the human dose. Due to this maternal toxicity, no conclusion can be drawn regarding the effects of defibrotide on embryo-foetal development. PAI-2 is known to be uniquely up-regulated in the placenta.

Juvenile Toxicity

Repeated intravenous administration of defibrotide, at doses below and close to the human therapeutic dose, to juvenile rats resulted in a delay in the mean age of preputial separation, suggesting a delay in the onset of male puberty in rats. However, the clinical relevance of these findings is unknown.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sodium citrate, dihydrate
Hydrochloric acid (for pH adjustment)
Sodium hydroxide (for pH adjustment)
Water for injection

6.2 Incompatibilities

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

6.3 Shelf life

Unopened vials: 2 years

In-use stability after first opening and/or dilution: from a microbiological point of view, after dilution, the reconstituted medicinal product should be used immediately. However, chemical and physical in-use stability has been demonstrated for 72 hours at 15-25°C for a concentration range of 4 mg/mL to 20 mg/mL in sodium chloride 9 mg/mL (0.9%) solution for infusion or 5% glucose solution for infusion at 15-25°C for 72 hours.

If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would not normally be expected to exceed 24 hours at 2-8°C..

6.4 Special precautions for storage

Do not freeze.

For storage conditions of the diluted medicinal product see section 6.3.

6.5 Nature and contents of container

2.5 mL vials (Ph. Eur. Type I clear glass), closed with a stopper (butyl rubber) and seal (aluminium) .

Pack size of 10 vials.

6.6 Special precautions for disposal and other handling

Defitelio is for single use only.

The concentrate solution for infusion has to be diluted using aseptic technique..

Defitelio should be diluted with sodium chloride 9 mg/mL (0.9%) solution for infusion or 5% glucose solution for infusion (see section 6.3 for concentration range and stability of the diluted solution) to a suitable concentration to permit 2 hours infusion time (see section 4.2).

Preparation of Defitelio (use aseptic technique):

1. The number of vials to be diluted should be determined based on the individual patient's weight (see section 4.2).
2. Before dilution, each vial should be inspected for particles. If particles are observed and/or the liquid in the vial is not clear, the vial must not be used.
3. The total volume of infusion should be determined based on the individual patient's weight. The final concentration of Defitelio should be in the concentration range of 4 mg/mL – 20 mg/mL (see section 6.3).
4. A volume of the sodium chloride 9 mg/mL (0.9%) solution for infusion or glucose 5% solution for infusion from the infusion bag should be withdrawn and discarded, equal to the total volume of Defitelio solution to be added.
5. The required volume from the Defitelio vials should be withdrawn and combined.
6. The combined volumes of Defitelio should be added to the sodium chloride 9 mg/mL (0.9%) solution for infusion or glucose 5% solution for infusion.
7. The solution for infusion should be mixed gently.
8. Prior to use the solution should be visually inspected for particulate matter. Only clear solutions without visible particles should be used. Depending on the type and amount of diluent the colour of the diluted solution may vary from colourless to light yellow. It is recommended that the diluted Defitelio solution be administered to patients using an infusion set equipped with a 0.2 µm in-line filter.

9. After the infusion is complete, the intravenous line should be flushed with sodium chloride 9 mg/mL (0.9%) solution for infusion or glucose 5% solution for infusion.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Gentium SpA
Piazza XX Settembre 2,
Villa Guardia,
22079 Italy
Phone: +39 031 5373200
Fax: +39 031 5373241
info@gentium.it

8. MARKETING AUTHORISATION NUMBER(S)

EU/01/13/878/001

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation:

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

ANNEX II

- A. MANUFACTURER OF THE BIOLOGICAL ACTIVE SUBSTANCE AND MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**
- E. SPECIFIC OBLIGATION TO COMPLETE POST-AUTHORISATION MEASURES FOR THE MARKETING AUTHORISATION UNDER EXCEPTIONAL CIRCUMSTANCES**

A. MANUFACTURER OF THE BIOLOGICAL ACTIVE SUBSTANCE AND MANUFACTURER RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer of the biological active substance

Gentium S.p.A.
Piazza XX Settembre, 2
22079 Villa Guardia (Como)
Italy

Name and address of the manufacturer responsible for batch release

Gentium S.p.A.
Piazza XX Settembre, 2
22079 Villa Guardia (Como)
Italy

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal products on “restricted” medical prescription, reserved for use in certain specialised areas (see Annex I: Summary of Product Characteristics, section 4.2).

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

• **Periodic safety update reports**

The marketing authorisation holder shall submit the first periodic safety update report for this product within 6 months following authorisation. Subsequently, the marketing authorisation holder shall submit periodic safety update reports for this product in accordance with the requirements set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and published on the European medicines web-portal.

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

• **Risk Management Plan (RMP)**

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the Marketing Authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

If the submission of a PSUR and the update of a RMP coincide, they can be submitted at the same time.

- **Additional risk minimisation measures**

The MAH shall agree the following with the National Competent Authority in each member state from which patients will be enrolled:

The format and content of the Healthcare professional material that highlights the existence of the Registry as well as the means to enter patients into the registry.

- **Obligation to conduct post-authorisation measures**

The MAH shall complete, within the stated timeframe, the below measures:

Description	Due date
Results of the validation of the SK-HEP-1 cell based assay should be provided. Together with the results a proposal to include the assay as an additional routine Quality Control test for batch release and stability testing for both defibrotide active substance and finished product should be provided.	March 2014

E. SPECIFIC OBLIGATION TO COMPLETE POST-AUTHORISATION MEASURES FOR THE MARKETING AUTHORISATION UNDER EXCEPTIONAL CIRCUMSTANCES

This being a marketing authorisation under exceptional circumstances and pursuant to Article 14(7) of Regulation (EC) No 726/2004, the MAH shall complete, within the stated timeframe, the following measures:

Description	Due date
Prior to launch, the Marketing Authorisation Holder (MAH) shall set up a patient registry to investigate the long term safety, health outcomes and patterns of utilisation of defibrotide during normal use. It shall be a multi-centre, multinational and prospective observational disease registry of patients diagnosed with severe hepatic VOD following haematopoietic stem cell plantation (HSCT) and enroll patients treated with defibrotide, other treatments or supportive care. The MAH shall ensure that information regarding all safety concerns identified in the most recent version of the Risk Management Plan is being collected. The MAH shall also ensure that all health care professionals who might prescribe defibrotide are provided with information on the importance of, and how to enter patients in, the registry.	Annual reports within the annual re-assessment

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING
BOX**

1. NAME OF THE MEDICINAL PRODUCT

Defitelio 80 mg/mL concentrate for solution for infusion
defibrotide

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each 2.5 mL vial contains 200 mg defibrotide.

3. LIST OF EXCIPIENTS

Also contains: sodium citrate dihydrate, hydrochloric acid and sodium hydroxide (for pH adjustment),
water for injections

4. PHARMACEUTICAL FORM AND CONTENTS

Concentrate for solution for infusion
10 vials

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
For intravenous use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT
OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP:

9. SPECIAL STORAGE CONDITIONS

Do not freeze

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS
OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF
APPROPRIATE**

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Gentium SpA
Piazza XX Settembre 2
Villa Guardia
22079 Italy
P: +39 031 5373200
F: +39 031 5373241
info@gentium.it

12. MARKETING AUTHORISATION NUMBER(S)

EU/01/13/878/001

13. BATCH NUMBER

Lot:

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

VIAL

1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION

Defitelio 80 mg/mL sterile concentrate
Defibrotide
Intravenous use

2. METHOD OF ADMINISTRATION

3. EXPIRY DATE

EXP:

4. BATCH NUMBER

Lot

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT

2.5 mL

6. OTHER

Gentium S.p.A.

B. PACKAGE LEAFLET

PACKAGE LEAFLET: INFORMATION FOR THE USER

Defitelio 80 mg/mL concentrate for solution for infusion

Defibrotide

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor.
- If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Defitelio is and what it is used for
2. What you need to know before you are administered Defitelio
3. How you will be given Defitelio.
4. Possible side effects.
5. How to store Defitelio.
6. Contents of the pack and other information.

1. What Defitelio is and what it is used for

Defitelio is a medicine that contains the active substance defibrotide.

It is used to treat a condition called hepatic veno-occlusive disease, in which the blood vessels in the liver become damaged and obstructed by blood clots. This can be caused by medicines that are given prior to a stem cell transplantation.

Defibrotide works by protecting the cells of the blood vessels and preventing or breaking down the blood clots.

This medicine can be used in adults, and in adolescents, children and infants over one month of age.

2. What you need to know before you take Defitelio

Do not use Defitelio

- if you are allergic to defibrotide or other ingredients of this medicine (listed in section 6)
- if you are using other medicines to break down blood clots such as tissue plasminogen activator

Warnings and precautions

Talk to your doctor before using Defitelio:

- if you have heavy bleeding and need a blood transfusion
- if you are undergoing surgery
- if you have problems with blood circulation because your body cannot maintain a constant blood pressure.

Children and adolescents

Defitelio is not recommended in children less than 1 month of age.

Other medicines and Defitelio

Tell your doctor if you are taking medicines to prevent blood clotting such as acetylsalicylic acid, heparins, warfarin, dabigatran, rivaroxaban or apixaban or if you are taking anti-inflammatory medicines (e.g., ibuprofen, naproxen, diclofenac and other non-steroidal anti-inflammatory medicines)

Pregnancy and breast-feeding

Do not use Defitelio if you are pregnant unless your disease requires treatment with Defitelio.

If you are sexually active and you or your partner could become pregnant, you both must use effective contraception during treatment with Defitelio and for 1 week after stopping the treatment.

Driving and using machines

It is not expected that Defitelio will affect your ability to drive and operate machinery.

Important information about some of the Ingredients of Defitelio

This medicine contains less than 23 mg of sodium, which means it is essentially “sodium-free”.

3. How you will be given Defitelio

The treatment with Defitelio can be initiated and continuously supervised only by an experienced doctor in an hospital or in a specialised centre for stem cells transplantation.

It will be slowly injected (over a 2-hour period) into one of your veins. This is called and ‘intravenous infusion’ or drip.

You will receive this treatment four times a day for at least 21 days or until your symptoms resolve. The recommended dose in children from one month to 18 years of age is the same as in adults.

If a dose of Defitelio has been forgotten:

As you will be given this medicine by a doctor or a nurse it is unlikely that a dose will be missed. However, tell your doctor or healthcare professional if you think that a dose has been forgotten. You must not be given a double dose to make up for a missed dose.

If you have any further questions on the use of this medicine, ask your doctor, nurse or pharmacist.

4. Possible side effects

Like all medicines, Defitelio can cause side effects, although not everybody gets them.

Common (may affect up to 1 in 10 people)

- bleeding in general
- bleeding from the nose
- bleeding in the brain
- bleeding in the gut
- bleeding in the lungs
- bleeding from the infusion line
- blood in the urine
- coagulopathy (disturbance of blood clotting)
- vomiting
- low blood pressure.

If you experience any of these events, you should **contact your doctor immediately**.

Uncommon (may affect up to 1 in 100 people)

- bleeding from the eye

- vomiting blood
- blood in the stool
- bleeding from the mouth
- bleeding at the site of injection
- localized blood collection out of the vessel (hematoma) in the brain
- Diarrhoea
- Nausea (feeling sick)
- Haemothorax (accumulation of blood in the area between the heart and the lung)
- Rash
- Itching
- Bruising
- Red spots on the body caused by broken blood vessels in skin
- Fever
- Severe allergic reaction (you might experience swelling of the hands, face, lips, tongue or throat, difficulty in breathing).

Children and adolescents

Side effects in children (1 month to 18 years old) are expected to be similar in type, severity and frequency and no other special precautions are needed.

Reporting of side effects

If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Defitelio

Keep this medicine out of the sight and reach of children.

Do not use Defitelio after the expiry date which is stated on the carton and label after EXP. The expiry date refers to the last day of that month.

Do not freeze.

Once diluted for use the infusion storage should not exceed 24 hours at 2-8°C unless dilution has taken place in controlled and validated aseptic conditions.

Defitelio should not be used if the solution is cloudy or contains particles.

6. Contents of the pack and other information

What Defitelio contains

- The active substance is defibrotide. Each 2.5 mL vial contains 200 mg defibrotide and each mL solution contains 80 mg defibrotide.
- The other ingredients are sodium citrate dihydrate, hydrochloric acid and sodium hydroxide (both for pH-adjustment) and water for injections.

What Defitelio looks like and contents of the pack

Defitelio is a clear light yellow to brown concentrate for solution for infusion, free from particulate matter or turbidity.

Carton containing 10 glass vials with 2.5 mL of concentrate.

Marketing Authorisation Holder and Manufacturer

Gentium SpA
Piazza XX Settembre 2
Villa Guardia
22079 Italy
P:+39 031 5373200

F: +39 031 5373241

info@gentium.it

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

België/Belgique/Belgien – Deutschland – España - France – Ireland – Italia – Luxembourg/Luxemburg – Malta – Nederland – Österreich – Portugal – United Kingdom

Gentium SpA
Piazza XX Settembre 2
Villa Guardia
22079 Italy
Tel: +39 031 5373200
info@gentium.it

България

Фармасуис ЕООД
ж.к. Лагера, ул.Троянски проход 16
BG-1612 София
Тел.: + 359 2 895 21 10
Bulgaria.Info@valeant.com

Česká republika

PharmaSwiss ČR s.r.o.
Jankovcova 1569/2c
17000 Praha 7
Tel.: +420-234 719 600
czech.info@valeant.com

Κύπρος

Phadisco Ltd
Λεωφόρος Γιάννου Κρασιδιώτη 185, CY-2234
Λατσιά,
Κύπρος
Τηλ.: +357 22 71500
Fax: +357 22 715100

Danmark/Ísland

Swedish Orphan Biovitrum A/S
Wilders Plads 5
DK-1403 København K
Tlf: + 45 32 96 68 69
mail.dk@sobi.com

Eesti

Swedish Orphan Biovitrum International AB
c/o CentralPharma Communications OÜ
Selise 26-11
EE-13522 Tallinn
Tel: + 372 6 015 540
centralpharma@centralpharma.ee

Lietuva

Swedish Orphan Biovitrum International AB
c/o CentralPharma Communications OÜ
J. Savickio g. 4-1
LT-01108 Vilnius
Tel: +370 5 2430444
centralpharma@centralpharma.lt

Magyarország

Valeant Pharma Magyarország Kft.
Csatárka u. 82-84, H-1025 Budapest
Hungary
Tel: +36-1-345-5900
Fax: +36-1-345-5918

Norge

Swedish Orphan Biovitrum AS
Østensjøveien 18, Bryn
N-0661 OSLO
Tlf: + 47 66 82 34 00
mail.no@sobi.com

Polska

ICN Polfa Rzeszów S.A.
2 Przemysłowa Street, 35-959 Rzeszow
Poland
Tel.: +48 17 865 5100
ICN_Polfa@valeant.com

România

PharmaSwiss Medicines S.R.L.
Str. Gheorghe Țițeica nr. 121 C, etaj 1
020295 Sector 2, București, România
Tel.: +40 374 102 600
Romania.info@valeant.com

Slovenija

PharmaSwiss d.o.o.
Wolfova ulica 1
SI-1000 Ljubljana
Slovenia
Tel: +386 1 236 47 00
slovenia.inforegulatory@valeant.com

Ελλάδα

Pharmaswiss Hellas A.E.
Λ. Πεντέλης, 15235 Βρυλήσσια
Ελλάδα
Τηλ.: +30-2108108460

Hrvatska

PharmaSwiss d.o.o.
Miramarska 23
10000 Zagreb
Hrvatska
Tel: +385 1 6311 833
croatia.info@valeant.com

Latvija

Swedish Orphan Biovitrum International AB
c/o CentralPharma Communications OÜ
Baložu iela 28-13
Rīga, LV-1048-13
Tel: + 371 67 450 497
centralpharma@centralpharma.lv

Slovenská republika

Valeant Slovakia s.r.o.
Polianky 5/A | 841 01 Bratislava
Tel : + 421 2 6920 3922
Fax : + 421 2 6920 3926

Suomi/Finland

Oy Swedish Orphan Biovitrum AB
Äyritie 18
FIN-01510 Vantaa
Puh/Tel: + 358 201 558 840
mail.fi@sobi.com

Sverige

Swedish Orphan Biovitrum AB (publ)
S-112 76 Stockholm
Tel: + 46 8 697 20 00
mail.se@sobi.com

This leaflet was last approved in

This medicine has been authorised under ‘exceptional circumstances’. This means that because of the rarity of this disease and for ethical reasons > it has been impossible to perform a placebo-controlled clinical trial and to get complete information on this medicine.

The European Medicines Agency will review any new information on this medicine every year and this leaflet will be updated as necessary.

Detailed information on this medicine is available on the European Medicines Agency web site:

<http://www.ema.europa.eu>

There are also links to other websites about rare diseases and treatments.

<-----