



**EUROPEAN COMMISSION**  
HEALTH & CONSUMER PROTECTION DIRECTORATE-GENERAL

Directorate C - Public Health and Risk Assessment  
**C7 - Risk assessment**

## **SCIENTIFIC COMMITTEE ON CONSUMER PRODUCTS**

### **SCCP**

#### **Preliminary Opinion on**

**Biological effects of ultraviolet radiation relevant to  
health with particular reference to sun beds for cosmetic purposes.**

Adopted by the SCCP  
during the 6<sup>th</sup> plenary of 13 December 2005

**TABLE OF CONTENTS**

1.	BACKGROUND	.....	3
2.	TERMS OF REFERENCE	.....	4
3.	OPINION	.....	6
4.	CONCLUSION	.....	23
5.	MINORITY OPINION	.....	27
6.	REFERENCES	.....	27
7.	ACKNOWLEDGEMENTS	.....	38
8.	APPENDIX A: TECHNIQUE FOR DETERMINING THE ERYTHEMAL-WEIGHTED IRRADIANCE		39

## 1. BACKGROUND

The main source of exposure to ultraviolet radiation (UVR) is the sun, but for some individuals substantial exposure occurs from artificial sources including sun beds for cosmetic purposes, industrial lamps, arc welding and medical UVR therapies.

There is evidence that UVR can cause damage to health.

In the context of a notification under the safeguard procedure in accordance with Article 9 of the Low Voltage Directive (LVD) 73/23/EEC<sup>1</sup>, a shortcoming in the European harmonised standard EN 60335-2-27:1997<sup>2</sup> has been brought to the attention of the European Commission by the Spanish authorities.

The LVD, a harmonisation Directive based on Article 95 of the EC Treaty, regulates the placing on the market of electrical appliances with a voltage rating between 50 and 1000 V (AC) and 75 and 1500 V (DC) with respect to health and safety. According to Article 95(3) of the EC Treaty<sup>3</sup> the LVD takes as its basis a high level of protection. Electrical appliances that comply with European harmonised standards under the LVD are presumed to comply with the corresponding essential health and safety requirements of the LVD.

The above mentioned shortcoming in the European harmonised standard EN 60335-2-27:1997, which has been reported to the European Commission, relates to the fact that the standard does not entirely cover the health and safety aspects which have to be considered during the design phase of the electrical appliance. In particular, it does not provide limit values on the maximum effective irradiance of UV radiation for the types of tanning devices that are covered by the scope of the standard.

In reaction to the notification and after consultation of governmental experts of Member States in the LVD ADCO<sup>4</sup> working group the Commission Services decided to request a scientific opinion from the “Non-Food Scientific Committees”.

The scientific opinion will be used when preparing a Commission mandate to the European standardisation organisations regarding:

- the revision of the above mentioned standard EN 60335-2-27:1997;
- drafting or revising product related standards covering risks associated with the exposure of persons to ultraviolet radiation (UVR).

---

<sup>1</sup> Council Directive 73/23/EEC of 19 February 1973 relating to electrical equipment designed for use within certain voltage limits (OJ L 77, 26.3.1973); Directive as amended by Directive 93/68/EEC (OJ L 220, 30.8.1993)

<sup>2</sup> EN 60335-2-27:1997 “Safety of household and similar electrical appliances - Part 2-27: Particular requirements for appliances for skin exposure to ultraviolet and infrared radiation”

<sup>3</sup> Article 95 of the EC Treaty see: [www.europa.eu.int/eur-lex/en/treaties/selected/livre221.html](http://www.europa.eu.int/eur-lex/en/treaties/selected/livre221.html)

<sup>4</sup> “Administrative co-operation” working group in the area of the LVD, consisting of the Market Surveillance representatives from all Member States and the European Commission

## 2. TERMS OF REFERENCE

The scientific committee is requested to answer the following questions in relation to the sun beds for cosmetic purposes:

1. What are the general health and safety implications (negative and positive) relating to the exposure of persons to ultraviolet radiation (UVR)<sup>5</sup>?
2. What are the differences between risks associated with exposure of persons to natural UVR and those risks from artificial UVR? What are the differences regarding the health and safety risks with respect to exposure of persons to UVA, UVB and UVC radiation respectively?
3. Is the total dose value of UVR the only effective health and safety parameter with regard to the risks associated with exposure of persons to both natural and artificial UVR? What is the validity of the Bunsen-Roscoe law<sup>6</sup> over the range of irradiances and wavelengths associated with exposure of persons to both natural and artificial UVR?
4. What are the specific health and safety implications (negative and positive) relating to the exposure of persons to UVR from tanning devices for cosmetic purposes?
5. Are limit values necessary for the irradiance of UVR from artificial sources, in particular from tanning devices for cosmetic purposes, with respect to health and safety? Is it necessary to give different values for the irradiance of UV-A, UV-B and UV-C radiation respectively? If so, please specify the limit values for the irradiance of artificial UVR above which adverse health effects will occur. What are the uncertainties of these limit values?
6. Please specify the limit values of total dose of artificial UV-A, UV-B and UV-C radiation above which adverse health effects will occur, taking into account skin phototype, intensity of exposure, duration of exposure and associated uncertainties.

### Supporting documents

- Spanish formal objection against European harmonised standard EN 60335-2-27
- [ICNIRP statement \(2003\)](http://www.icnirp.org/documents/sunbed.pdf) (<http://www.icnirp.org/documents/sunbed.pdf>)
- [WHO guidance brochure: artificial tanning sunbeds \(2003\)](http://www.who.int/uv/publications/en/sunbeds.pdf) (<http://www.who.int/uv/publications/en/sunbeds.pdf>)
- ESA (European Sunlight Association) Position paper
- [ESA \(European Sunlight Association\) frequently asked questions](http://www.europeansunlight.org/test/esa/html/faq.htm) (<http://www.europeansunlight.org/test/esa/html/faq.htm>)
- [NRPB: Health effects from Ultraviolet Radiation V13 No.1 2002](http://www.nrpb.org/publications/documents_of_nrpb/pdfs/doc_13_1.pdf) ([http://www.nrpb.org/publications/documents\\_of\\_nrpb/pdfs/doc\\_13\\_1.pdf](http://www.nrpb.org/publications/documents_of_nrpb/pdfs/doc_13_1.pdf))

<sup>5</sup> The International Commission on Illumination (CIE) defines ultraviolet radiation (UVR) as optical radiation between 100 and 400 nm. The spectral region is divided into three photo-biological spectral regions: UVC (100-280 nm), UVB (280-315 nm) and UVA (315-400 nm).

<sup>6</sup> The Bunsen-Roscoe law (law of reciprocity) states that a certain biological effect is directly proportional to the total energy dose irrespective of the administered regime. Dose is the product of intensity and the duration of exposure. (Bunsen R, Roscoe HE, *Photochemische Untersuchungen*, Poggendorff's Annalen 1855: 96: 373-394, 1857: 100: 43-88 and 481-516, 1857: 101:235-263, 1859: 108: 193-273.).

- **NRPB: Advice on Protection Against Ultraviolet Radiation V13 No.3 2002** ([http://www.nrpb.org/publications/documents\\_of\\_nrpb/pdfs/doc\\_13\\_3.pdf](http://www.nrpb.org/publications/documents_of_nrpb/pdfs/doc_13_3.pdf))
- **SSK: Schutz des Menschen vor den Gefahren der UV-Strahlung in Solarien** (<http://www.ssk.de/2001/ssk0101w.pdf>)
- Scientific investigations from **ECOFYS**
- Other documents

### **National and international organisations contributing to the discussion**

- ICNIRP - International Commission on Non-Ionizing Radiation Protection (<http://www.icnirp.org>)
- WHO - World Health Organization (<http://www.who.int>)
- IARC - International Agency for Research on Cancer (<http://www.iarc.fr>)
- UNEP - United Nations Environment Programme ([http://www.unep.org/PDF/Solar\\_Index\\_Guide.pdf](http://www.unep.org/PDF/Solar_Index_Guide.pdf))
- NRPB - National Radiological Protection Board (UK) (<http://www.nrpb.org>)
- SSK – Strahlenschutzkommission (Germany) (<http://www.ssk.de>)
- IMM – Institute of Environmental Medicine (Sweden) (<http://www.imm.ki.se>)
- EPA - U.S. Environmental Protection Agency (US) (<http://www.epa.gov>)
- FDA - U. S. Food and Drug Administration (US) (<http://www.fda.gov>)
- NIES - National Institute for Environmental Studies (JP) ([www@nies.go.jp](http://www.nies.go.jp))
- List of experts provided by ESA (European Sunlight Association)

### 3. OPINION

#### 1. What are the general health and safety implications (negative and positive) relating to the exposure of persons to ultraviolet radiation (UVR)?

##### 1.1 Negative Effects

##### 1.1.1 Acute

##### Skin

Exposure of the skin to UVR results in inflammation (erythema/sunburn) that is usually maximal about 24 hours later (Farr et al, 1988). Erythema is associated with increased blood flow (Young et al, 1985), increased sensitivity to thermal and mechanical stimuli (Harrison et al, 2004), a dermal inflammatory infiltrate (Gilchrest et al, 1983; Hawk et al, 1988) and the presence of apoptotic keratinocytes known as sunburn cells (Sheehan and Young, 2002). Individual sensitivity to erythema can be assessed by determining the minimal erythema dose (MED) that increases with skin type as shown in Table 1 but MED is not predictive of skin type because there is considerable variation of MED within different white skin types (Harrison and Young, 2002). Within a few days of exposure delayed melanogenesis (tanning) occurs that is dependent on skin type. This results from the synthesis of melanin in melanocytes: specialized pigment producing cells in the epidermis that transfer melanin to keratinocytes. Many people expose themselves to UVR for the sole purpose of obtaining a tan that becomes more intense with repeated exposure. This repeated exposure also results in thickening of the epidermis, especially the *stratum corneum*, the outermost dead layer, which results in the skin feeling dry. A UVB tan is photoprotective against erythema but the level of photoprotection is modest and equivalent to a sunscreen with a sun protection factor (SPF) of 2-3 (Agar and Young, 2005). However, tans primarily induced by UVA are not photoprotective against erythema (Gange et al, 1985). UVR exposure, in particular UVA, results in transient immediate pigment darkening (IPD) the function of which is not known (Routaboul et al, 1999).

**Table 1 Classification of different skin phototypes**

Skin Photo Type	Sunburn Susceptibility	Tanning Ability	Classes Of Individuals	Indicative MED (range in parentheses) SED <sup>§</sup>
I	Always burn	No tan	Melano-compromised	2 (1 - 3)
II	High	Light tan		
III	Moderate	Medium tan	Melano-competent	5 (3 - 7)
IV	Low	Dark tan		
V	Very low	Natural brown skin	Melano-protected	8 (7 - >12)
VI	Extremely low	Natural black skin		

<sup>§</sup> The unit of erythemal radiation is the Standard Erythema Dose (SED), where 1 SED is equivalent to an erythemal effective radiant exposure of 100 Jm<sup>-2</sup> (CIE 1998). It requires an exposure of about 3 SED to produce just minimal erythema in the unacclimatized white skin of the most common northern European skin types (Harrison & Young 2002). An exposure of 5-8 SED will result in moderate sunburn and 10 SED or more can result in a painful, blistering sunburn.

Solar UVR exposure can aggravate certain skin diseases such as lupus erythematosus and pemphigus (Morison et al, 1999) and induce skin photosensitivity with commonly used UVR-absorbing systemic drugs and topically encountered chemicals (Ferguson et al, 1999). Furthermore, there is a wide range of acquired and genetic UVR and visible radiation photodermatoses that are beyond the scope of this document.

Exposure of the skin to UVR can suppress cell-mediated immunity when assessed with the sensitisation (Kelly et al, 2000) and the elicitation arms (Moyal and Fourtanier, 2003) of the contact hypersensitivity (CHS) response. A single sub-erythral exposure of solar simulating radiation (SSR) suppresses the induction (sensitisation) arm of the CHS response in skin types I/II (Kelly et al, 2000) but erythral exposure is necessary to suppress the elicitation arm (Moyal and Fourtanier, 2003). Suppression of cell-mediated immunity is thought to play a role in UVR-induced skin cancer and infectious diseases, e.g. Herpes simplex infections.

The clinical effects of UVR exposure, whether acute or long-term, are underpinned by many molecular and cellular events (Matsumura and Ananthaswamy, 2002). UVR-induced damage to epidermal DNA, especially cyclobutane pyrimidine dimers (CPD), is thought to be responsible for many adverse effects of solar UVR, including immunosuppression, and can be demonstrated in the skin immediately after exposure to erythral and sub-erythral UVR (Young et al, 1998). DNA integrity is maintained by complex repair processes and the p53 mediated elimination of damaged cells by apoptosis (sunburn cell formation). Failure of these processes is thought to result in skin cancer (Matsumura and Ananthaswamy, 2002). Membrane as well as DNA effects also contribute to UVR-induced skin damage. The relevant cell surface or cytoplasmic chromophores are currently unknown. There is considerable evidence that the photoisomerization of stratum corneum trans-urocanic acid (UCA) to the cis-form also plays an important role in immunosuppression. Exposure to erythral UVR or repeated sub-erythral UVR results in a loss of epidermal antigen presenting Langerhans cells (Novakovic et al, 2001).

## **Eye**

The eye is a complex multi-layered organ that receives visible radiation on its retina. The intermediate layers attenuate UVR to different degrees and thereby protect the retina from UV photodamage. The outermost cornea absorbs UVC and a substantial amount of UVB, which is further attenuated by the lens and the vitreous humor in front of the retina. UVA is less well attenuated by the cornea but is attenuated by the internal structures so it does not reach the retina (Slinney, 2001; Roberts, 2001; Johnson, 2004).

The only acute clinical effect of UVR on the eye is photokeratitis that is also known as snow blindness or welder's flash (Slinney, 2001; Roberts, 2001; Johnson, 2004). This is a painful transient inflammatory condition caused by UVC and UVB-induced damage to the corneal epithelium. Typically it appears 6-12 hours after exposure and resolves, without long-term consequences, within 48 hours. In some ways it can be regarded as sunburn of the eye.

## 1.1.2 Chronic

### Skin Cancer

#### Non-melanoma

Solar exposure is recognized as the main environmental factor in the development of basal cell carcinoma (BCC) and squamous cell carcinoma (SCC) that form the great majority of skin cancers (IARC, 1992). These lesions result in a high level of morbidity with only occasional mortality from infrequent metastatic SCC. UVR is also associated with actinic keratoses (AK) that are regarded as biomarkers for increased risk of SCC.

The evidence for UVR in these lesions has been primarily ecologic (reviewed by Armstrong and Kricger, 2001), supported by mouse studies in the case of SCC (de Gruijl, 1995). More recently, a role for UVR has been supported by the presence of UVR “signature mutations” in tumours (Brash et al, 1996). Skin type is an important determinant of BCC and SCC risk with skin types I and II at greater risk than skin types III and IV, with the lowest risk being in skin types V and VI. SCC is associated with chronic UVR exposure and is more common in people with outdoor occupations. There is evidence that BCC is associated with intermittent exposure (Kricger et al, 1995). Registration of BCC and SCC is often poor compared with melanoma but there is evidence that the incidence rate is increasing substantially in Europe (Boyle et al, 2004). Data from the skin cancer registry in Trentino, Italy showed incidence rates of 88 per 100,000 for BCC and 29 per 100,000 for SCC in the period 1993-1998 in comparison to 14 per 100,000 for melanoma (Boi et al, 2003).

#### Melanoma

Though much less common than BCC and SCC, melanoma is the main cause of death from skin cancer. There were an estimated 35,000 cases of melanoma diagnosed in Europe in 2000 with 9000 deaths (Boyle et al, 2004). Sun exposure is established as the major environmental determinant of melanoma (IARC, 1992; Donawho et al, 1994; Armstrong and Kricger, 1993) and the risk of melanoma depends on the interaction between environmental exposures and the genes which determine susceptibility. Melanoma is rare in black skinned peoples (Parkin et al, 1997) in whom vitamin D levels tend to be lower, presumably as a result of lower levels of cutaneous synthesis (Brazzol et al, 1988; Dawson-Hughes, 2004) despite an apparently similar ability to manufacture vitamin D in blacks and whites treated with artificial UVR (Brazzol, 1988). Vitamin D levels are thought to be important for many aspects of human health and therefore any UVR exposure advice given to European Citizens will depend upon the balance of risk of skin cancer versus possible benefit and will therefore be determined by genes governing skin colour and those that determine susceptibility to skin cancer. The health and safety implications for sun exposure in Europeans, is therefore critically dependent upon skin colour and this is an important component of the health advice messages given. Difficulties relate to the complexities of the relationship between skin colour or type and susceptibility and additional problems caused by some still unresolved issues pertaining to the type of sun exposure, which is causal for melanoma.

There is no doubt that skin colour and sun exposure are potent determinants of risk of melanoma. World incidence figures show that the risk to individuals is greatest where pale skinned peoples live at low latitudes such as Australia and New Zealand (Parkin et al, 1997; Bulliard, 2000). In areas of the world where dark and pale skinned peoples live at high UV exposure levels, such as



Hawaii, then the risk to pale skinned people is much greater than for their darker skinned neighbours (Chuang et al, 1999). Within Europe there is variation in incidence which reflects the interaction between skin colour and latitude as the peak incidence is in the north, in countries such as Sweden, where fair skinned peoples live an outdoor life and have access to sunny holidays in the south, or Switzerland where fair skinned peoples live at high altitude (Parkin et al, 1997). So, in the period 1996-8 the incidence rates (European Standardised Rates) in women were reported to be 17 per 100,000 in Switzerland, 6 per 100,000 in Spain and 16 per 100,000 in Sweden (de Vries and Coebergh, 2004).

Broadly, it would be reasonable to conclude that the risks of melanoma are so low in black-skinned peoples that sun protection advice should be directed only towards white-skinned peoples. The difficulty here is that skin colour is a continuous rather than a discontinuous variable. Some Asian peoples have quite a high tendency to burn and within white skinned peoples there is variation in susceptibility to sunburn and to melanoma which is related to skin colour and whether there are freckles or not. Data from many case control studies have established that phenotypic characteristics associated with vulnerability to the sun are risk factors for melanoma. Gandini et al (2005a) recently summarized these in a meta-analysis of 60 such studies. Her overall conclusions were that skin type I (versus IV) was associated with a relative risk (RR) of 2.1 for melanoma (95% CI 1.7-2.6), where skin type I, is skin which always burns and never tans and skin type IV is skin which never burns. A high density of freckles was associated with a RR=2.1 (95% CI 1.8-2.5), eye colour (Blue vs. Dark: RR=1.5, 1.3-1.7) and hair colour (Red vs. Dark: RR=3.6, 2.6-5.4). Hence, whatever the ethnic origin of Europeans, in terms of skin type, health advice about skin cancer should be directed to those individuals with a tendency to burn rather than to tan, those who have freckles and those with fair (particularly red) hair. It is clear from the level of these risk factors that the relative risk is significant but the absolute risk associated with these phenotypic characteristics is relatively small in European countries with incidence rates of between 5 and 17 per 100,000 per annum (European Standardised Rates) (de Vries and Coebergh, 2004). The prevalence of individuals with these risk factors will vary considerably between populations. In a study of healthy women in the UK, 8% had red hair and 6% had very high freckle scores on the back (Bertram et al, 2002).

Risk of melanoma is also greater in patients with larger numbers of melanocytic naevi whether banal or clinically atypical, where an atypical naevus is defined as a mole 5mm or greater in diameter, with an irregular or ill-defined edge and variable pigmentation. Numerous case-control studies have addressed this, and a second meta-analysis by Gandini et al (2005b) showed that the number of common naevi was confirmed as an important risk factor for melanoma with a substantially increased risk associated with the presence of 101-120 naevi compared with <15 (pooled Relative Risk (RR) = 6.9; 95% Confidential Interval (CI): 4.6, 10.3) as was the number of atypical naevi (RR = 6.4 95%; CI: 3.8, 10.3; for 5 versus 0). Twin studies have provided strong evidence that naevus number is genetically determined (Wachsmuth et al, 2001; Zhu et al, 1999; Easton et al, 1991) and the association of the phenotype with melanoma risk therefore implies the presence of naevus genes, which are also low penetrance melanoma susceptibility genes. Thus, persons with this atypical naevus phenotype have an increased risk of melanoma, which is significantly higher than that associated with red hair or freckles. The prevalence of this phenotype also varies between populations but was reported in 2% of individuals in the UK (Bataille et al, 1996).

The phenotypes described above are genetically determined and therefore it is not surprising that family history is a risk factor for melanoma. Familial melanoma was reported in the 19<sup>th</sup> century

in the UK (Norris, 1820), and a strong family history of melanoma is the most potent risk factor for melanoma (Kefford et al, 1999). Any family history of melanoma is associated with a doubling of risk for close relatives. A study from the Utah population database estimates risk to first-degree relatives of melanoma cases to be 2.1 (95% CI 1.4-2.9). A similar study from the Swedish Cancer Registry estimated the standardized incidence ratio for melanoma to be 2.4 (95% CI 2.1-2.7) for offspring if one parent had a melanoma, 3.0 (95% CI 2.5-3.5) for an affected sibling and 8.9 (95% CI 4.3-15.3) if a parent and a sibling were both affected. The highest ratio was 61.8 (95% CI 5.8-227.2) for offspring when a parent had multiple melanomas (Hemminki et al, 2003). Such patterns of risk are indicative of a significant hereditary component, which is most probably inherited as an autosomal dominant trait with incomplete penetrance. The risk of melanoma increases with age although in Europe the age distribution curve is relatively flat and in Europe the incidence is commonly higher in women than in men (Parkin et al, 1997).

Sun exposure is clearly the major environmental risk factor for melanoma as discussed above. A third meta-analysis reported by Gandini et al (2005c) has supported the conclusions of many individual case-control studies that intermittent sun exposure remains the most predictive environmental risk factor for melanoma (random effects model RR=1.6 (95% CI 1.3-2.0) and that sunburn, especially in childhood is a significant risk factor, although there was much heterogeneity between studies. A random effects model suggested a highly significant effect for sunburn at any age (RR=2.0 95% CI 1.7-2.4). The pooled analysis provided no evidence for a causal effect of chronic sun exposure on melanoma risk, RR=1.0 (95% CI 0.9-1.0). Further evidence for a role of sun exposure in melanoma comes from penetrance studies for the melanoma susceptibility gene CDKN2A in which there was evidence for an interaction between susceptibility genes and latitude of residence so that penetrance was highest in families with germline CDKN2A mutations living in Australia when compared with those in Europe (Bishop et al, 2002).

A meta-analysis, incorporating latitude, showed that phenotypic indicators of excessive sun exposure (representing gene/environment interaction) in fair-skinned individuals are risk factors for melanoma (Gandini et al, 2005a). Pre-malignant and malignant lesions were associated with a RR=4.3 (2.8-6.6) and actinic damage indicators with a RR=2.0 (1.2-3.3). This is of note despite the lack of epidemiological evidence from case-control studies for chronic sun exposure as a risk factor for melanoma.

In summary, there is strong evidence that excessive sun exposure is causal for melanoma. Evidence persists that the exposure pattern is important, e.g. intermittent, although the observation in some studies that actinic skin damage is a risk factor provides some evidence that chronic over-exposure is also causal in some patients. The evidence is also strong that excessive sun exposure increases the risk of melanoma in those with a strong family history. There is an emerging view, based upon epidemiological and biological studies that there may be more than one route to melanoma: one associated with low or intermittent sun exposure and for which numerous naevi is a risk factor, and another with chronic over exposure (Whiteman et al, 2003). All of the risk factors quoted above are independent risk factors in individual case control studies and therefore the presence of multiple risk factors in an individual increases the relative risk of melanoma.

Health education is postulated to be most effective when targeted at those at greatest risk. Thus, UVR risk communication to European citizens is probably best directed at those with established

risk factors (e.g. family history, fair skin and multiple naevi). There is a need to communicate these complex issues to the European citizen in a way that is easily understood.

### **Photoageing**

Exposure of the skin to UVR results in UVR-induced skin ageing known as photoageing, which is very evident, when one compares normally sun-exposed (face) and sun-protected (buttock) sites. Clinical symptoms of photoageing include wrinkling, laxity and disturbances of the distribution of pigmentation (Glogau, 1996). Photoageing is thought to at least partially arise from the induction of matrix metalloproteinases (MMPs) that degrade collagen, the major structural protein of the dermis (Fisher et al, 2002). Photoageing, assessed by elastosis, is an indicator of non-melanoma skin cancer risk (Kricger et al, 1991).

### **Effects on the Eye**

There is limited epidemiological evidence that solar UVR exposure results in cataracts of the lens, anterior lens capsular change and pterygium (Johnson, 2004). *In vivo* and *ex vivo* acute studies on mammalian lens (Pitts et al, 1977; Merriam et al, 2000; Oriowo et al, 2001) and a chronic *in vivo* study (Jose and Pitts, 1985) have indicated that the UVB part of the solar spectrum is most likely to be responsible for any long term effects that solar UVR has on the lens.

## **1.2 Positive Effects**


Exposure to solar UVR initiates the synthesis of vitamin D, in the skin, that is vital for musculo-skeletal health (Vieth, 2005) and there is evidence that large numbers of people are vitamin D insufficient (Holick, 2005). There is also emerging evidence, as yet mainly ecologic (i.e. by association), that vitamin D is important in the prevention of autoimmune disorders (Ponsonby et al, 2005) and several internal malignancies (Berwick and Kesler, 2005).

Exposure to UVR has been associated with a sense of “well being” (Diffey, 1986) but this has not been linked to any change in circulating serotonin, melatonin or opioid peptides (Gambishler et al, 2002a, 2002b). UVB induces  $\beta$ -endorphin production in keratinocytes *in vitro*, which, at least in theory, may have a systemic effect (Gilchrest et al, 1996).

## **2(a) What are the differences between risks associated with exposure of persons to natural UVR and those risks from artificial UVR?**

There are no physical differences between natural and artificial UVR *per se*. However, there are important differences in the spectral distribution and absolute and relative irradiances of UVR from the sun and artificial sources, and between different artificial sources as shown in Figure 1. There is no standard solar spectrum because this varies with factors such as season, latitude and time of day. From a physical standpoint, the sun is primarily a UVA source with a maximal UVB content of about 5%. Artificial UVR emission spectra are not similar to solar UVR from a physical point of view.

It is relatively easy to compare the acute risks of exposure to natural and artificial UVR, which are similar, the details of which are discussed in Section 4. It is much more difficult to compare chronic effects which, in the sun, also depend on patterns of exposure.



<sup>¶</sup> Measured in Melbourne (38° S) at solar noon on 17 January 1990. Measurements were made at the Australian Radiation Laboratory with a Spex 1680B double monochromator with a resolution of 1 nm

**Figure 1 Emission spectra of solar UVR and two tanning lamps (i) Cleo natural and (ii) Cleo performance**

Data on the risk of skin cancer associated with artificial UVR sources (see Section 4) are few compared with those related to sun exposure. Furthermore the tanning device studies are often uninformative because of small numbers of cases and controls, and low usage of the devices. There are also great difficulties in collecting adequate exposure data because of recall bias and lack of user knowledge of the type of UVR emitted by the devices. Many studies therefore have recorded only whether a tanning device has been used “ever” or “never” so that the power to address dose or age effects is limited.

Furthermore, tanning device users are often those who sunbathe frequently and it is likely that case-control studies are seriously confounded. There are some data published on the effects of medical use of artificial UVR sources. Although the UVR dose is considerably lower than that to which users of tanning devices are potentially exposed, such studies do have the merit of much more accurate dosage estimation.

Photo(chemo)therapy is used in the treatment of skin diseases. The use of psoralen plus UVA (PUVA) to treat psoriasis is known to cause skin cancer (Stern and Laird, 1994) but PUVA is mechanistically quite different from UVA and UVB and therefore is not relevant to the current discussion. In the PUVA cohort study reported by Stern from the United States, there was no discernable additional effect of exposure to UVB (Stern and Laird, 1994). In a study of psoriatics treated with coal tar and UVB in the 1950s followed up for 25 years there was no demonstrable increased risk of skin cancer but the numbers treated were relatively small (280) (Pittelkow et al, 1981). In an even smaller study of 195 German psoriatics treated with broadband (n=69) or narrow band UVB (n= 126) from 1994 to 2000 only one skin cancer had occurred by 2004. This was an *in situ* melanoma which developed in the same year that narrow band UVB therapy was begun (Weischer et al, 2004). A study in Scotland with a median follow

up period of 4 years has shown a small increase in BCC after treatment with narrow band UVB phototherapy (Man et al, 2005).

Overall, the risks of skin cancer from the medicinal use of artificial UVR (in the absence of photosensitizers) appear to be small but the data are few and the dose to which the patients are exposed tends to be significantly smaller than users of commercial sun beds are potentially exposed to. It is likely, from our knowledge of skin cancer and solar UVR, that the skin cancer risk attributable to artificial UVR will be greater in those who are genetically susceptible such as the fair skinned.

Household lights emit significant amounts of UVR (Sayre et al, 2004) and several case-control studies have addressed risk for melanoma associated with such exposure. The earliest study suggested an elevated risk associated with exposure to fluorescent lights at work (Beral et al, 1982) but all subsequent studies failed to identify such a risk (Osterlind et al, 1988; Rigel et al, 1983; Walter et al, 1992; Holly et al, 1995).

## **2(b) What are the differences regarding the health and safety risks with respect to exposure of persons to UVA, UVB and UVC radiation respectively?**

Coblentz introduced the concept of the spectral regions UVA, UVB and UVC at the Second International Congress on Light in Copenhagen in 1932. These regions were determined by the transmission properties of three common glass filters; a barium-flint filter defined the UVA (315-400nm); a barium-flint-pyrex filter the UVB (280-315nm); and a pyrex filter defined the UVC (wavelengths shorter than 280nm). So the basis of these divisions has its grounding in physics, and not biology. Although these are the official designations of the Commission Internationale de l'Éclairage (CIE), other authorities, especially in the biological and clinical sciences, use different definitions such as UVA (320-400nm), UVB (290-320nm) and UVC (190-280nm).

### **2(b).1 Acute Effects**

The wavelength dependency of a given photobiological effect is demonstrated by its action spectrum, which depends on a variety of factors but is based on the absorption spectrum of the chromophore (UVR absorbing biomolecule) and the optical properties of the skin. Action spectroscopy and studies with different broad-spectrum sources show that UVB is much more effective than UVA for most acute endpoints studied in human skin. This includes erythema (CIE 1998; Young et al, 1998), delayed pigmentation (Parrish et al, 1982), DNA photodamage (Young et al, 1998) and UCA photoisomerization (McLoone et al, 2005). In general, UVB is 3 to 4 orders of magnitude more effective per unit physical dose ( $J/cm^2$ ) than UVA, but this difference depends on the specific wavelengths/wavebands being compared. Action spectra for immunosuppression in human skin are not available. UVB is known to be immunosuppressive but the role of UVA is still not clear (Phan et al, 2006). The action spectrum for IPD shows that UVA is more effective than UVB (Irwin et al, 1993).

UVC is not an issue for terrestrial solar UVR because it is completely absorbed by the ozone layer. In any case, UVC is strongly attenuated by chromophores in the upper epidermis (Young, 1997) and UVC-induced DNA damage in the dividing basal layer of human epidermis is not readily detected (Campbell et al, 1993; Chadwick et al, 1995) which may explain why the dose response curve for UVC erythema in human skin is very much less steep than for UVB (Diffey

and Farr, 1991). It is unlikely that UVC from artificial sources presents an acute or long-term hazard to human skin.

Wavelength dependency is crucial in determining the biological effect of a given spectral region of a UVR source. For example, the 0.8% UVB content of a tanning lamp accounted for 75% of the CPD that it induced in human keratinocytes *in vitro* (Woollons et al, 1999). Thus action spectra are essential as weighting functions to determine the biological effects of different UVR emission spectra (see Section 5). Action spectra are only valid if there is no interaction between different spectral regions. However, there is evidence that such interactions do occur at the cellular level (Schieke et al, 2005).

### **2(b).2 Chronic Effects**

Wavelength dependency for chronic effects has been determined in hairless mouse models and these studies have shown that the action spectra for SCC (de Gruijl, 1995) and photoageing (elastosis) (Kligman and Sayre, 1991) are very similar to the human action spectrum for erythema (CIE, 1998; Young et al, 1998). Thus, one might conclude that sunburn, primarily caused by UVB, can be regarded as a surrogate risk factor for SCC and photoageing. There is no animal model for UVR-induced BCC.

Sunburn, an important risk factor for melanoma, has therefore implicated UVB in its pathogenesis (Wang et al, 2001). The incidence of melanoma, as well as BCC and SCC, is very high in xeroderma pigmentosum (XP) with defective excision repair of UVB-type DNA damage, e.g CPD. The wavelength dependency for melanoma however is not yet established because of the lack of a good animal model (Noonan et al, 2003). Melanomas have proved extremely difficult to induce by UVR alone in mice. Wavelength dependency has been determined in a fish model (*Xiphophorus*) (Schartl et al, 1997) the value of which is limited because its melanoma-like lesions arise from the dermis instead of the epidermis and fish are phylogenetically very different from humans. Studies in these fish however showed that visible and UVA radiation, as well as UVB (Setlow et al, 1993) induced lesions that raised concern that UVA might be causal for human melanoma as well or instead of UVB. A mammalian opossum model also developed melanoma-like lesions after broad-band UVA exposure but with low potency compared to broad-band UVB (Robinson et al, 2000). A mouse model was described in 2003 (the hepatocyte growth factors/scatter factor transgenic mouse) in which melanomas with a strong epidermal component were induced (Nonnan et al, 2003). Neonatal UV irradiation was necessary and sufficient to induce melanoma although adult irradiation increased the number of lesions. In 2004 the same group reported studies using the mouse in which UVB but not UVA induced melanoma, providing perhaps more persuasive evidence that UVB exposure is causal rather than UVA (De Fabo et al, 2004).

Studies of somatic mutations in a variety of genes have been reported in the search for evidence to support a role for UVB exposure. Genes such as p53 have, however, failed to show the characteristic UVB signature C to T transitions and CC to TT mutations, providing additional concern that UVB may not be the key causal waveband. Recently, mutations in BRAF (downstream of RAS) were found in a majority of naevi and melanoma. The dominant point mutation (T1796A) is not characteristic of UVB radiation, but this does not exclude a causal role for UVR (de Gruijl, 2003).

It is more difficult to determine UVA induced mutagenesis because DNA does not significantly absorb UVA at doses obtained with solar exposure. It is thought that UVA induced mutagenesis is mainly mediated by photosensitising reactions that generate reactive oxygen species. In one system it was suggested that T to G transversions are typical of UVA induced damage (Drobetsky et al, 1995) but in another G to T transversions were seen as well as small tandem base deletions (Pfeifer et al, 2005). There is no consensus on UVA signature somatic mutations in tumours. Furthermore, it is possible that UVA may have an indirect adverse effects on the micro-environment in the dermis and dermo-epidermal junction by inducing growth factor release which may have a proliferative effect on melanocytes (Brenner et al, 2005).

In summary, UVB is likely to be the main cause of photoageing and SSC. Sunburn, a marker for excessive UVR exposure, is a risk factor for melanoma. UVB is the main cause of sunburn but this does not necessarily mean that it is the prime cause of melanoma, the spectral dependence of which remains unknown. The conservative approach is to restrict UVB and UVA exposure in susceptible phenotypes until wavelength dependency is established.


**3 (a) Is the total dose value of UVR the only effective health and safety parameter with regard to the risks associated with exposure of persons to both natural and artificial UVR? (b) What is the validity of the Bunsen-Roscoe law over the range of irradiances and wavelengths associated with exposure of persons to both natural and artificial UVR?**

Experiments in which the photoresponse of a material is investigated as a function of radiant flux (dose rate or irradiance) are commonly called *reciprocity law experiments*. Bunsen and Roscoe (1859) are credited with conducting the first reciprocity law experiments. Reciprocity holds in photobiology when the observable response depends only on the total administered radiant exposure (commonly referred to as *dose*) and is independent of the two factors that determine total dose, that is, irradiance and exposure time.

Since the reciprocity law only depends on total dose, its validation for a particular end-point can have many experimental manifestations. Assuming that the reciprocity law is valid, then each manifestation should be equivalent to the others as long as the integrated total dose is the same. Thus, when the reciprocity law is obeyed, the same photobiological response is observed when specimens receive the same integrated total dose regardless as to whether the exposure is performed:

- a) At a high radiant flux for a short period of time.
- b) At a low radiant flux for a long period of time.
- c) By repeatably switching a light source on-or-off and controlling both the on-off frequency of the light and the length of time that the light remains in the on and the off state. Experiments in which the light is turned on-and-off at an extremely high frequency are called *flash photolysis experiments*, while experiments in which the light is turned on-and-off at a low frequency are called *intermittency experiments*.
- d) By ramping the radiant flux to a high level, holding the flux for a specified period of time, and then ramping it back down to a lower level or any variant of these stress regimes.

These exposure regimes are depicted graphically in Figure 2 (a-d) (from Martin *et al* 2003).



**Figure 2 Different temporal patterns for delivering a given UVR dose**

A summary of reciprocity experiments in human and mice skin is given in Table 2. In almost every case, reciprocity was shown to hold. Of particular relevance to sunbed use, where exposure times will vary between a few minutes up to half-an-hour or so depending on the power and spectral output of the lamps, Meanwell & Diffey (1989) showed that exposure to polychromatic radiation for time periods ranging from 1s to 1h induced degrees of delayed erythema ranging from minimal to marked that depended only on dose and not dose rate. These findings both support and extend those of previous studies in which the end point was confined to minimal erythema. There are no quantitative human data for long-term effects of UVR on the risk of skin cancer, but patterns of exposure may be important, especially for melanoma (see Question 1) that suggests a failure of the reciprocity law



**Table 2 UVR reciprocity studies in human and mouse skin. Mouse tumorigenesis studies are for SCC**

Response	Source	Spectrum	Irradiation	Range in irradiance	Reciprocity?	Reference
<b>Human skin</b>						
Erythema	Mercury	Monochromatic	Continuous	4	Y	Hausser 1927
Erythema	Mercury	Monochromatic	Continuous	?	Y	Hausser 1928
Erythema	Mercury	Monochromatic	Continuous	8	Y	Luchiesh 1930
Erythema	Mercury	Monochromatic	Continuous	4	Y	Coblentz 1932
Erythema	Mercury	Monochromatic	Continuous	20	Y	Blum 1946
Erythema	Mercury	Monochromatic	Flash	200	Y	Schmidt 1963
Erythema	Mercury	Monochromatic	Continuous	10 <sup>3</sup>	Y	Park 1984
Erythema	Xenon	Monochromatic	Intermittent	?	Y	Everett 1969
Erythema	Xenon	Monochromatic	Continuous	?	Y	Everett 1969
Erythema	Xenon	Polychromatic	Continuous	10 <sup>3</sup>	Y	Meanwell 1989
Erythema	Laser	Monochromatic	Flash	3	Y	Parrish 1976
Erythema	Laser	Monochromatic	Continuous	10 <sup>4</sup>	Y	Anderson 1980
<b>Mice skin</b>						
Tumorigenesis	Mercury	Polychromatic	Continuous	12	Y	Blum 1941
Tumorigenesis	Mercury	Polychromatic	Intermittent	4	Y	Blum 1942
Tumorigenesis	Mercury	Polychromatic	Intermittent	4	N	Bain 1943
Tumorigenesis	Xenon	Monochromatic	Intermittent	3	N	Forbes 1979
Tumorigenesis	Fluorescent	Polychromatic	Continuous	5	N	Forbes 1981
Tumorigenesis	Fluorescent	Polychromatic	Continuous	8	Y	de Gruijl 1982
Immunosuppression	Fluorescent	Polychromatic	Intermittent	1	Y	DeFabo 1979
Immunosuppression	Fluorescent	Polychromatic	Continuous	10	Y	DeFabo 1980
Immunosuppression	Fluorescent	Polychromatic	Continuous	10	Y	Noonan 1981

#### **4 What are the specific health and safety implications (negative and positive) relating to the exposure of persons to UVR from tanning devices for cosmetic purposes?**

##### **4.1 Negative Effects**

###### **4.1.1 Acute and non skin cancer effects**

The use of tanning devices has been associated with acute adverse reactions such as a form of skin fragility known as pseudoporphyria (Murphy et al, 1990; Weiss and Jung, 1990) and lentigines (Salisbury et al, 1989; Kadunce et al, 1990) that have been noted in case reports. There have also been case reports of induction (Fruchter and Edoute, 2004) and exacerbation (Stern and Docken, 1986) of systemic lupus erythematosus.

There is a risk of phototoxic reactions with people using certain medications (Bisland, 1990) or applying topical aromatherapy products, such as bergamot oil, that contain photosensitising chemicals (Kaddu and Wolf, 2001) or eating plants that contain such chemicals (Ljunggren, 1990).

Devices with a higher UVB/UVA ratio or a high UVB irradiance will be more effective at tanning and will require a shorter exposure time. However, this also increases the likelihood of a

burn (doses > 1 MED) because there is a lower margin of error in the determination of exposure time (see Section 5). Burns have also been reported due to equipment failure (Eltigani and Mathews, 1994).

Studies in the late 1980s showed that the use of tanning devices has an adverse effect on human immune function (Hersey et al, 1987; Rivers et al, 1989). More recently, Whitmore and Morison (2000) reported that 10 full-body exposures over a two-week period suppressed immunity as assessed by the induction and elicitation arms of the contact hypersensitivity response. These authors also studied the effect of 10 full-body tanning exposures in 11 volunteers and, not surprisingly, reported the presence of CPD and p53 protein expression in keratinocytes *in vivo* (Whitmore et al, 2001).

#### **4.1.2 Chronic Skin Cancer**

##### Non-melanoma

Very few studies have been done on the relationship between sunbed use and non-melanoma skin cancer risk. Two hospital-based case-control studies in Ireland, in the mid to late 1980s, did not show any relationship between the use of tanning devices and non-melanoma skin cancer (O'Loughlin et al, 1985; Herity et al, 1989). A similar conclusion, at about the same time, was reached by Bajdik et al (1996) in British Columbia, Canada, who evaluated 406 controls (population based) against 180 SCC cases and 226 BCC cases. About 10% of each group had "ever" used a sunlamp. The adjusted OR for BCC and SCC for "ever" having used a sunlamp were 1.2 (0.7-2.2) and 1.4 (0.7-2.7) respectively, which are clearly non-significant. One 2002 study, using the "generalized estimating equation method" reported no significant effect of tanning devices for BCC, even though the total lifetime exposure to tanning devices was almost twice as high in patients compared with controls (Boyd et al, 2002). In the same year, Karagas et al (2002) assessed the relationship between use of tanning devices and BCC and SCC in a population-based case control study. In this study there was greater use of tanning devices ranging from 9.2% (male controls) to 28.4% (female patients). The OR for BCC and SCC were 1.5 (1.1-2.1) and 2.5 (1.7-3.8) respectively and adjustment for a variety of factors made no difference to these results. The results of Karagas et al (2002) suggest that the use of tanning devices is a risk factor for non-melanoma skin cancer.

##### Melanoma

Sun bed usage has increased considerably in recent years (Rafnsson et al, 2004) but the data on melanoma risk are scanty. There are a number of case-control studies but the details on exposure for the majority was small and all, as case-control studies, were subject to bias of recall and the effect of confounders. There is a single cohort study (Verierod et al, 2003) in which risk of melanoma was addressed.

A number of case-control studies reported no evidence of sun bed use as a risk factor for melanoma (Osterlind et al, 1988; Holly et al, 1995; Westerdal et al, 1994; Zanetti et al, 1988; Chen et al, 1998; Dunn-lane et al, 1993; Naldi et al, 2000; Bataille et al 2004, 2005). The majority of these studies were, however, small and the prevalence of sun bed usage in cases and controls was very low. Others were supportive of weak evidence or evidence in "at risk" groups (Walter et al, 1990; Westerdahl et al, 2000). Walter et al (1990) showed some suggestion of a trend to increased risk of melanoma with longer duration of use. In the study by Westerdahl et al (2000) an increased risk of melanoma was demonstrated only for use of sun beds before the age

of 35 years (OR, 2.3; CI, 1.2–4.2). Swerdlow et al (1988) showed a significantly increased risk for any use of sun beds OR 2.94 (95% CI 1.4-6.17) with a significant trend for increased duration of use. Autier et al (1994) showed little evidence of risk overall when corrected for skin type etc but did show evidence of increased risk for usage of sun beds for 10 hours or more, when burning was reported after use of the sun bed or when the users reported use of the sun bed to tan.

The only cohort study to address risk associated with solarium followed more than 100,000 Norwegian women for an average of 8 years, and 187 melanomas developed. This study identified use of a solarium for 1 or more times per month as a risk factor for melanoma (Veierod et al, 2003). This is probably the most persuasive evidence for a role for sun beds in causing melanoma but the data are as yet relatively weak and support the view only that frequent use is deleterious.

Gallagher et al (2005) carried out a meta-analysis of 9 case-control studies and the one cohort study and came to the conclusion that sunbed use significantly increased the risk of melanoma with an OR of 1.25 (1.1-1.5) “ever” versus “never” used. This increased to 1.69 (1.3 –2.2) using the metric “first exposure as a young adult”.

### **Photoageing**

There seems to be no published literature on the photoageing effects of sunbed use but this would be expected from the long-term use of sunbeds because photoageing is associated with solar exposure (Fisher et al, 2002). Some studies have looked at the effect of repeated suberythral exposure of UVB and UVA in human skin and reported some changes that are associated with photoageing (Lavker et al, 1995a, 1995b; Lavker and Kaidby, 1997).

As with the sun, tanning devices emit infrared radiation (IR: 760nm to 1mm). The effects of IR on skin are poorly understood but *in vitro* studies suggest that it may play a role in photoageing, which has been suggested by animal studies (Schieke et al, 2003).

## **4.2 Positive Effects**

### **Tanning**

The use of a sunbed results in a tan that is the desired outcome sought by the user. This has been demonstrated and quantified in a study that followed a Food and Drug Administration (FDA) protocol (Caswell, 2000) with 3 weekly exposures for 8 weeks. A significant tanning effect was evident after 6 exposures and the level of tan increased steadily over the 8-week assessment period. Another study, with twice weekly exposure for 6 weeks, demonstrated tanning (Ruegger et al, 2002). There are few data on the photoprotective properties of a sunbed-induced tan but one study showed a protection factor of about 3 against UVB-induced erythema (Rivers et al, 1989). This is consistent with a wide range of laboratory and field studies (Agar and Young, 2005). The effect of such protection on skin cancer is not known. The use of a sunscreen in combination with an induced tan would theoretically result in an overall protection factor that would approximate the SPF multiplied by the protection factor of the tan.

**Vitamin D status**

Tanning with UVB-emitting sunbeds would be expected to improve vitamin D status and this has been reported in a recent study (Tangpricha et al, 2004) that showed that people who used a sunbed at least once a week for at least 6 months had a mean serum concentration of 25 hydroxyvitamin D (25(OH)D) of  $115.5 \pm 8.0$  (SEM) nmol/L compared with the controls who had levels of  $60.3 \pm 3.0$  nmol/L ( $P < 0001$ ). The tanners also had significantly higher hipbone mineral density. However, this study has several flaws; (i) it relied on recall of sunbed use without establishing serum 25(OH)D before sunbed use, (ii) the tanning group had much greater sunlight exposure and (iii) there was a much greater proportion of white-skinned people in the tanning group. Furthermore, there were no data on the spectral output of the tanning devices used.

**Feel good factor**

Many people claim to feel better after sunbed use (Diffey 1986) but studies using primarily UVA emitting sunbeds showed that mood effects could not be attributed to circulating serotonin or melatonin (Gambichler et al, 2002a) or opioid peptides (Gambichler et al, 2002b). The possible role of UVB-induced keratinocyte-derived  $\beta$ -endorphin (Gilchrest et al, 1996) has yet to be investigated.

- 5 (a) Are limit values necessary for the irradiance of UVR from artificial sources, in particular from tanning devices for cosmetic purposes, with respect to health and safety? (b) Is it necessary to give different values for the irradiance of UVA, UVB and UVC radiation respectively? (c) If so, please specify the limit values for the irradiance of artificial UVR above which adverse health effects will occur. What are the uncertainties of these limit values?**

From the above discussion on reciprocity, it is clear that acute clinical effects resulting from sunbed use (i.e. erythema) are likely to depend only on total dose and not dose rate. It is not possible to make any statements on the risk of skin cancer, especially melanoma. Since all tanning devices emit a broad UVR spectrum, the spectral profile of which determines the device's effectiveness to elicit clinical effects, it is irrelevant to specify irradiance limits in different spectral wavebands, especially since the spectral regions UVA, UVB and UVC were originally based on the optical properties of different glasses (see section 2b). A more appropriate way to speak of tanning devices than using the terms UVA, UVB and UVC is to compare their erythemal power, as a percentage of total UVR power, with sunlight. This is expressed mathematically as:

$$100 \times \frac{\sum_{290}^{400} E(\lambda) \cdot \varepsilon(\lambda) \cdot \Delta\lambda}{\sum_{290}^{400} E(\lambda) \cdot \Delta\lambda}$$

$E(\lambda)$  is the relative spectral power distribution of the UV source and  $\varepsilon(\lambda)$  is the erythemal effectiveness of radiation of wavelength  $\lambda$  nm (CIE 1998). An example of this calculation is given in Appendix A. For the 3 sources illustrated in the Figure 1, the erythemally effective percentages are 0.44%, 0.51% and 0.13% and for summer sunlight, the "Cleo Natural" and "Cleo Performance" lamps, respectively. Clearly, the "Cleo Natural" lamp more closely resembles sunlight "biologically" than the "Cleo Performance" lamp.


In specifying an upper limit of irradiance, the important quantity is the erythemally-weighted irradiance, obtained by weighting each spectral irradiance component of the lamp by its relative effectiveness to induce erythema and summing over all wavelengths present in the source spectrum (see equation above). To minimise the risk of timing errors, which might result in “sunburn” it is desirable that the prescribed sunbed exposure session should be no less than 5 minutes. The avoidance of “sunburn” may also reduce the risk of melanoma.

Depending on an individual’s phototype, the exposure in SED during this 5-minute period should not exceed the subject’s estimated indicative MED (see Table 1). For people with sun-sensitive skin this exposure is around 2 SED, leading to a maximum erythemal-weighted irradiance of 25 SED/h ( $0.7 \text{ W/m}^2$ ).

The main conclusion from this analysis is that the erythemally weighted properties of a given sunbed emission spectrum (as demonstrated in Appendix A) are more important than its physical properties *per se*. This is because UVB is orders of magnitude more erythemogenic than UVA which, as shown in Figure 1, is the main UVR component of sunlight and tanning device spectra. At present, certainty can only be reasonably assured for acute effects such as erythema.

**6 Please specify the limit values of total dose of artificial UVA, UVB and UVC radiation above which adverse health effects will occur, taking into account skin phototype, intensity of exposure, duration of exposure and associated uncertainties.**

The clinical effects of UVR exposure can be either *deterministic*, where the magnitude of the effect is related to exposure and a threshold dose is possible, or *stochastic* in which the probability of the effect is related to exposure and there is no threshold dose. Erythema is an example of a deterministic effect and SCC is a stochastic effect, as illustrated in Figure 3.



**Figure 3 Differences between deterministic and stochastic effects of UVR**

There is no need to specify total dose separately for UVA, UVB and UVC for the reasons given in Section 5. It is not possible to specify uncertainties for long-term effects.

For a single exposure on a sunbed it is important to avoid marked or severe erythema but necessary (for the desired cosmetic effect) to receive a sufficient UVR dose to stimulate

melanogenesis. Experience has shown that an exposure at or just below that to induce a just perceptible MED 8 to 24 hours after exposure approximates the optimum.

A classification of skin phototypes based on susceptibility to sunburn in sunlight (WHO 2003), together with indicative MEDs that might be expected following exposure on unacclimatized skin, is given in Table 1. Depending on an individual's phototype, the exposure in SED on any single occasion should not exceed their estimated indicative MED.

With a stochastic effect like SCC skin cancer there is no threshold dose below which the effect will not occur. Consequently, any recommendation about a limit value of total dose accumulated over a specific time period (e.g. year, lifetime) is arbitrary and subjective. A limit of 20 sessions per year (equivalent to an exposure of approximately 40 SED or 20 MED in melano-compromised individuals) was proposed by the British Photodermatology Group (BPG) in 1990 (Diffey *et al* 1990) and was subsequently adopted by the UK Health & Safety Executive (HSE 1995).

The International Electrotechnical Commission (1995) recommends that the maximum annual exposure should not exceed an erythemal-weighted dose of 15 kJ/m<sup>2</sup> (150 SED), equivalent to around 50 MED in white-skinned people. Not surprisingly, it is this higher limit adopted by The Sunbed Association in the UK in their code of practice for operators (TSA 2004).

Other agencies have simply advised against sunbed use and not specified an "acceptable" maximum annual usage (AGNIR 2002; WHO 2003; ICNIRP 2003).

Estimates have been made of the risk of basal and squamous cell skin cancers arising from sunbed use (AGNIR 2002) and what constitutes an "acceptable" risk is a matter of judgment. For most people, who may use sunbeds 10 or 20 times a year for 10 years or so in young adulthood, the estimated additional lifetime risk of non-melanoma skin cancer, compared with non-users, is up to 10% (AGNIR 2002).

Case-control studies, particularly more recent ones, have generally found an association between sunbed use and melanoma (Young 2004) with an odds ratio of around 1.5. In communicating this risk to policy makers (Heller *et al* 2003), it may be helpful to estimate the potential number of cases and deaths prevented each year in a population if sunbeds were eliminated. These are given in Table 3 for the UK, where the relative risk of incidence and mortality of sunbed users is taken to be 1.5 relative to non-users. Melanoma incidence and mortality data refer to the year 2002 (Cancer Research UK, 2005).

Sex	Millions in UK Population in 2002	Melanoma per 10 <sup>5</sup>		% using sunbeds	Relative risk of incidence & mortality	Population attributable risk	No. affected in 2002		Population impact by eliminating sunbeds	
		Incidence	Mortality				Incidence	Mortality	Incidence	Mortality
Male	28.7	11.2	3.0	5	1.5	0.024	3193	874	78	21
Female	30.3	13.7	2.5	9	1.5	0.043	4128	770	179	33
						<b>Total</b>	<b>7321</b>	<b>1644</b>	<b>257</b>	<b>54</b>

**Table 3 Non-use of tanning devices might have resulted in about 54 fewer deaths from melanoma in 2002 than the 1644 that were observed in the UK**

An alternative approach to estimating the mortality associated with sunbeds is through modeling population exposure to both sunlight and sunbeds, assuming that the patterns of exposure from these two sources are equally carcinogenic, that the melanomas that result are equally fatal, and that the fraction of deaths due to sunbed use is equal to the population exposure from sunbeds expressed as a fraction of the total population exposure from sunlight and sunbeds. Using this approach Diffey (2003) estimates the mortality due to sunbed use each year in the UK is around 100, with a range of about 50 to 200. The estimates from this approach and that illustrated in Table 3 are not inconsistent given the many uncertainties and assumptions involved. However, from the above discussion it is clear that there is no limit value of total dose of artificial UVR below which adverse health effects will not occur and that any limit is subjective and arbitrary

#### 4. CONCLUSION

**Question 1: What are the general health and safety implications (negative and positive) relating to the exposure of persons to ultraviolet radiation (UVR)<sup>7</sup>?**

- Clinically relevant UV-radiation is UVB (280 – 315 nm) and UVA (315-400 nm)
- Solar UVB (~295 – 315nm) is primarily responsible for inducing erythema (sunburn) and tanning
- UVA has similar acute clinical effects to UVB if the physical doses (J/cm<sup>2</sup>) given are approximately 1000 times greater
- Human skin may be phenotypically classified into phototypes I – VI which are determined by acute sensitivity to sunlight, melanin content and tanning ability
- Solar exposure is associated with basal cell carcinoma, squamous cell carcinoma and malignant melanoma.

<sup>7</sup> The International Commission on Illumination (CIE) defines ultraviolet radiation (UVR) as optical radiation between 100 and 400 nm. The spectral region is divided into three photo-biological spectral regions: UVC (100-280 nm), UVB (280-315 nm) and UVA (315-400 nm).

- The risk of a given type of skin cancer is influenced by patterns of UVR exposure
- Phototype is a good indicator of skin cancer risk which reflects acute sensitivity to sunlight with phototype I being the most sensitive and phototype VI being the most resistant
- Moles and freckles are good indicators of susceptibility to malignant melanoma
- Moles and freckles are independent risk factors for skin cancer
- UVR is immunosuppressive in humans, the consequences of which are unknown but may be important in skin cancer and infectious diseases
- Public health messages should be directed to those people at greatest risk of skin cancer in order to promote behaviour which is appropriate to the balance of risk
- Solar UVR, especially UVB, causes photokeratitis (snow blindness) of the eye and has been implicated in cataract formation

**Question 2: What are the differences between risks associated with exposure of persons to natural UVR and those risks from artificial UVR? What are the differences regarding the health and safety risks with respect to exposure of persons to UVA, UVB and UVC radiation respectively?**

- There are no intrinsic differences between the physical and biological properties of natural and artificial UVR but there are differences in spectral profile that may have biological consequences
- It is relatively easy to compare the acute effects of natural and artificial UVR but much more difficult to compare the long-term effects.
- UVR, with and without photosensitizers, is used in the phototherapy of skin diseases and skin cancer is an accepted risk in such treatment
- Wavelength dependency (action spectrum) studies show that UVB is the most harmful part of the solar UVR spectrum for both acute and term-effects but wavelength dependency data on UVA are more limited than for UVB
- We lack data to make conclusive statements on the wavelength dependency of melanoma



**Question 3: Is the total dose value of UVR the only effective health and safety parameter with regard to the risks associated with exposure of persons to both natural and artificial UVR? What is the validity of the Bunsen-Roscoe law<sup>8</sup> over the range of irradiances and wavelengths associated with exposure of persons to both natural and artificial UVR?**

- The reciprocity law applies for human erythema
- There are no quantitative human data for long-term effects but patterns of exposure may be important, especially for melanoma (see Question 1) that suggests a failure of the reciprocity law

**Question 4: What are the specific health and safety implications (negative and positive) relating to the exposure of persons to UVR from tanning devices for cosmetic purposes?**

- There are some case reports for adverse clinical effects (other than skin cancer) from sun bed use but it is not possible to estimate the frequency of these
- Several studies, and two meta-analyses, have shown a significant association between sunbed use and malignant melanoma. Typically, the measure of association is less than 2
- Sunbed use can result in the desired cosmetic outcome which is tanning
- There is no evidence to support a pharmacological basis for the “feel good” factor of sunbed use
- Use of sunbeds that contain UVB may enhance vitamin D status but there are few data available on this relationship. The emission spectrum of the source is likely to be important

**Question 5: Are limit values necessary for the irradiance of UVR from artificial sources, in particular from tanning devices for cosmetic purposes, with respect to health and safety? Is it necessary to give different values for the irradiance of UV-A, UV-B and UV-C radiation respectively? If so, please specify the limit values for the irradiance of artificial UVR above which adverse health effects will occur. What are the uncertainties of these limit values?**

- The biological consequences of a given sunbed emission spectrum are much more relevant than its specific irradiances within different wavebands which were originally defined by physical rather than biological parameters
- A biologically effective dose can be obtained by weighting a given emission spectrum with a relevant action spectrum

---

<sup>8</sup> The Bunsen-Roscoe law (law of reciprocity) states that a certain biological effect is directly proportional to the total energy dose irrespective of the administered regime. Dose is the product of intensity and the duration of exposure. (Bunsen R, Roscoe HE, Photochemische Untersuchungen, Poggendorff's Annalen 1855: 96: 373-394, 1857: 100: 43-88 and 481-516, 1857: 101:235-263, 1859: 108: 193-273.)

- This weighting should be done with the human erythema action spectrum, which is similar to the tanning action spectrum. This gives an erythemally weighted irradiance of the emission spectrum of the sunbed as demonstrated in Appendix A
- A sunbed session should be no less than 5 minutes to minimize errors of timing and therefore inadvertent burning. This period should deliver less than 1 minimal erythema dose (MED) for a sun-sensitive skin type (i.e. I or II), which is equivalent to an erythemally weighted irradiance of 0.7W/m<sup>2</sup>, or 25 standard erythema doses (SED) per hour
- Certainty can only be reasonably assumed for acute effects

**Question 6: Please specify the limit values of total dose of artificial UV-A, UV-B and UV-C radiation above which adverse health effects will occur, taking into account skin phototype, intensity of exposure, duration of exposure and associated uncertainties.**

- There is no need to specify different dose limits for UVC, UVB and UVA for the same reasons given in Question 5
- The dose limits for adverse acute effects are dealt with in Question 5. In the context of risk assessment, it is not possible to give dose limits for skin cancer because of lack of human dose-response data. However, SCC is a stochastic effect for which there is no assumed threshold dose. Any annual dose limits given are arbitrary
- The human erythema action spectrum is similar to the mouse SCC action spectrum and is likely to represent the wavelength dependency of human SCC, and possibly BCC. However, we lack mammalian data on the wavelength dependency of malignant melanoma. Broad spectrum studies in mice indicate that as with non melanoma skin cancer, UVB is much more important than UVA
- The important biological risk factors for malignant melanoma are age, sex (in some populations), skin phenotype, moles, freckles and family history. Behavioural/environmental risk factors include intermittent sunburning UVR exposure, especially in youth.

#### Overall Conclusion

The SCCP is of the opinion that the use of sunbeds to achieve and maintain cosmetic tanning, whether by UVB and/or UVA, is likely to increase the risk of malignant melanoma.

People with known risk factors for skin cancer, especially malignant melanoma, should be advised not to use sunbeds. Specifically, these are (i) age under 18, (ii) skin phenotypes I and II and the presence of freckles, (iii) multiple and/or moles and (iv) a family history of melanoma.

## 5. MINORITY OPINION

Not applicable

## 6. REFERENCES

- Agar, N. and A. R. Young. Melanogenesis: a photoprotective response to DNA damage? *Mutat. Res.* 571:121-132, 2005.
- AGNIR. Advisory Group on Non-ionising Radiation Health. Effects from Ultraviolet Radiation. Documents of the NRPB, 2002, 13 (1)
- Anderson RR, Parrish JA. A survey of the acute effects of UV lasers on human and animal skin, in: R. Pratesi, C.A. Sacchi (Eds.), *Lasers in Photomedicine and Photobiology*, Springer, New York, 1980, p. 109
- Armstrong, B.K. and A. Kricger, How much melanoma is caused by sun exposure? *Melanoma Res*, 1993. **3**(6): p. 395-401.
- Autier, P., et al., Cutaneous malignant melanoma and exposure to sunlamps or sunbeds: an EORTC multicenter case-control study in Belgium, France and Germany. EORTC Melanoma Cooperative Group. *Int J Cancer*, 1994. **58**(6): p. 809-13.
- Bain JA, Rusch HP. Carcinogenesis with ultraviolet radiation of wavelength 2800–3400 Å. *Cancer Res* 1943; **3**: 425
- Bajdik, C. D., R. P. Gallagher, G. Astrakianakis, G. B. Hill, S. Fincham and D. I. McLean. Non-solar ultraviolet radiation and the risk of basal and squamous cell skin cancer. *Br. J. Cancer* 73:1612-1614, 1996.
- Bataille, V., et al., Risk of cutaneous melanoma in relation to the numbers, types and sites of naevi: a case-control study. *Br J Cancer*, 1996. **73**(12): p. 1605-11.
- Bataille, V., et al., Exposure to the sun and sunbeds and the risk of cutaneous melanoma in the UK: a case-control study. *Eur J Cancer*, 2004. **40**(3): p. 429-35.
- Bataille, V., et al., A multicentre epidemiological study on sunbed use and cutaneous melanoma in Europe. *Eur J Cancer*, 2005. **41**(14): p. 2141-9.
- Beral, V., et al., Malignant melanoma and exposure to fluorescent lighting at work. *Lancet*, 1982. **2**(8293): p. 290-3.
- Bertram, C.G., et al., An Assessment of the CDKN2A Variant Ala148Thr as a Nevus/Melanoma Susceptibility Allele. *J Invest Dermatol*, 2002. **119**(4): p. 961-965.
- Berwick, M. and D. Kesler. Ultraviolet Radiation Exposure, Vitamin D, and Cancer. *Photochem. Photobiol.* 2005.

- 
- Bilsland, D. and W. S. Douglas. Sunbed pseudoporphyria induced by nalidixic acid. *Br. J. Dermatol.* 123:547, 1990.
- Bishop, D.T., et al., Geographical variation in the penetrance of CDKN2A mutations for melanoma. *J Natl Cancer Inst*, 2002. **94**(12): p. 894-903
- Blum HF, Kirby-Smith JS, Grady HG. Quantitative induction of tumors in mice with ultraviolet radiation. *J Natl Cancer Inst* 1941; **2**: 259
- Blum HF, Grady HG, Kirby-Smith. Relationships between dosage and rate of tumor induction by ultraviolet radiation. *J Natl Cancer Inst* 1942; **3**: 91
- Blum HF, Terus WS. Inhibition of the erythema of sunburn by large doses of ultraviolet radiation. *Am J Physiol* 1946; **146**: 97
- Boi, S., M. Cristofolini, R. Micciolo, E. Polla and P. P. Dalla. Epidemiology of skin tumors: data from the cutaneous cancer registry in Trentino, Italy. *J. Cutan. Med. Surg.* 7:300-305, 2003.
- Boyd, A. S., Y. Shyr and L. E. King, Jr. Basal cell carcinoma in young women: an evaluation of the association of tanning bed use and smoking. *J. Am. Acad. Dermatol.* 46:706-709, 2002.
- Boyle, P., J. F. Dore, P. Autier and U. Ringborg. Cancer of the skin: a forgotten problem in Europe. *Ann. Oncol.* 15:5-6, 2004.
- Brash, D. E., A. Ziegler, A. S. Jonason, J. A. Simon, S. Kunala and D. J. Leffell. Sunlight and sunburn in human skin cancer: p53, apoptosis, and tumor promotion. *J. Investig. Dermatol. Symp. Proc.* 1:136-142, 1996.
- Brazerol, W.F., et al., Serial ultraviolet B exposure and serum 25 hydroxyvitamin D response in young adult American blacks and whites: no racial differences. *J Am Coll Nutr*, 1988. **7**(2): p. 111-8.
- Brenner, M., et al., Differential expression of melanoma-associated growth factors in keratinocytes and fibroblasts by ultraviolet A and ultraviolet B radiation. *Br J Dermatol*, 2005. **153**(4): p. 733-9.
- Bulliard, J.L., Site-specific risk of cutaneous malignant melanoma and pattern of sun exposure in New Zealand. *Int J Cancer*, 2000. **85**(5): p. 627-32.
- Bunsen RW, Roscoe HE. Photochemische untersuchungen. *Ann Phys* 1859; **108**: 193
- Campbell, C., A. G. Quinn, B. Angus, P. M. Farr and J. L. Rees. Wavelength specific patterns of p53 induction in human skin following exposure to UV radiation. *Cancer Res.* 53:2697-2699, 1993.
- Cancer Research UK, <http://info.cancerresearchuk.org/cancerstats/>, accessed [January][2006].
- Caswell, M. The kinetics of the tanning response to tanning bed exposures. *Photodermatol. Photoimmunol. Photomed.* 16:10-14, 2000.

- 
- Chadwick, C. A., C. S. Potten, O. Nikaido, T. Matsunaga, C. Proby and A. R. Young. The detection of cyclobutane thymine dimers, (6-4) photolesions and the Dewar photoisomers in sections of UV-irradiated human skin using specific antibodies, and the demonstration of depth penetration effects. *J. Photochem. Photobiol. B* 28:163-170, 1995.
- Chen, Y.T., et al., Sunlamp use and the risk of cutaneous malignant melanoma: a population-based case-control study in Connecticut, USA. *Int J Epidemiol*, 1998. **27**(5): p. 758-65.
- Chuang, T.Y., et al., Melanoma in Kauai, Hawaii, 1981-1990: the significance of in situ melanoma and the incidence trend. *Int J Dermatol*, 1999. **38**(2): p. 101-7.
- CIE Standard. Erythema reference action spectrum and standard erythema dose. CIE S 007/E-1998. Vienna: Commission Internationale de l'Éclairage, 1998
- Coblentz WW, Stair R, Hogue JM. The spectral erythemic reaction of the untanned human skin to ultra-violet radiation. *Bur Stand J Res* 1932; **8**: 541 (Research Paper No. 433)
- Dawson-Hughes, B., Racial/ethnic considerations in making recommendations for vitamin D for adult and elderly men and women. *Am J Clin Nutr*, 2004. **80**(6 Suppl): p. 1763S-6S.
- DeFabo EC, Kripke ML. Dose-response characteristics of immunologic unresponsiveness to UV-induced tumors produced by UV irradiation of mice. *Photochem Photobiol* 1979; **30**: 385
- DeFabo EC, Kripke ML. Wavelength dependence and dose-rate independence of UV radiation induced suppression of immunologic unresponsiveness of mice of a UV-induced fibrosarcoma. *Photochem Photobiol* 1980; **32**: 183
- DeFabo, E.C., et al., Ultraviolet B but not ultraviolet A radiation initiates melanoma. *Cancer Res*, 2004. **64**(18): p. 6372-6.
- de Gruijl FR, van der Leun JC. Effect of chronic UV exposure on epidermal transmission in mice. *Photochem Photobiol* 1982; **36**: 433
- de Gruijl, F. R. Action spectrum for photocarcinogenesis. *Recent Results Cancer Res*. 139:21-30, 1995.
- de Gruijl, F.R., IL-40 UV Carcinogenesis and Melanocytes. *Pigment Cell Res*, 2003. **16**(5): p. 591.
- de Vries, E. and Coebergh, J.W. Cutaneous malignant melanoma in Europe. *European J Cancer*. 2004, 40:2355-2366
- Diffey, B. L. Use of UV-A sunbeds for cosmetic tanning. *Br. J. Dermatol*. 115:67-76, 1986.
- Diffey, B. L. and P. M. Farr. Sunbed lentiginos. *Br. Med. J. (Clin. Res. Ed)* 296:1468, 1988.
- Diffey BL, Farr PM, Ferguson J, Gibbs NK, de Gruijl FR, Hawk JLM, Johnson BE, Lowe G, MacKie RM, McKinlay AF, Moseley H, Murphy GM, Norris PG, Young AR. Tanning with ultraviolet A sunbeds. *Br Med J* 1990; **301**: 773-4

- 
- Diffey, B. L. and P. M. Farr. Quantitative aspects of ultraviolet erythema. *Clin. Phys. Physiol Meas.* 12:311-325, 1991.
- Diffey BL. A quantitative estimate of melanoma mortality from ultraviolet A sunbed use in the UK. *Br J Dermatol* 2003; **149**: 578-581
- Donawho, C.K., P. Wolf, and M.L. Kripke, Enhanced development of murine melanoma in UV-irradiated skin: UV dose response, waveband dependence, and relation to inflammation. *Melanoma Research*, 1994. **4**: p. 93-100.
- Drobetsky, E.A., J. Turcotte, and A. Chateauneuf, A role for ultraviolet A in solar mutagenesis. *Proc Natl Acad Sci U S A*, 1995. **92**(6): p. 2350-4.
- Dunn-Lane, J., et al., A case control study of malignant melanoma. *Ir Med J*, 1993. **86**(2): p. 57-9.
- Easton, D., et al., Genetic susceptibility to naevi- a twin study. *Br J Cancer*, 1991. **64**: p. 1164-1167
- Eltigani, E. and R. N. Matthews. An unusual cause of sunbed burns. *Burns* 20:87-88, 1994.
- Everett MA, Sayre RM, Olson RL. Physiologic response of human skin to ultraviolet light. in: F. Urbach (Ed.), *The Biologic Effects of Ultraviolet Radiation*, Pergamon Press, New York, 1969, p. 181.
- Farr, P. M., J. E. Besag and B. L. Diffey. The time course of UVB and UVC erythema. *J. Invest Dermatol.* 91:454-457, 1988.
- Ferguson J. Druge and chemical photosensitivity. In: *Photodermatology*, Hawk JLM (ed) London: Arnold, 1999, pp. 155-169.
- Fisher, G. J., S. Kang, J. Varani, Z. Bata-Csorgo, Y. Wan, S. Datta and J. J. Voorhees. Mechanisms of photoaging and chronological skin aging. *Arch. Dermatol.* 138:1462-1470, 2002.
- Forbes PD, Davies RE, Urbach F. Aging, environmental influences, and photocarcinogenesis. *J Invest Dermatol* 1979; **73**: 131
- Forbes PD, Blum HF, Davies RE. Photocarcinogenesis in hairless mice dose–response and the influence of dose-delivery. *Photochem Photobiol* 1981; **34**: 361
- Fruchter, O. and Y. Edoute. First presentation of systemic lupus erythematosus following ultraviolet radiation exposure in an artificial tanning device. *Rheumatology. (Oxford)* 44:558-559, 2005.
- Gallagher, R. P., J. J. Spinelli and T. K. Lee. Tanning beds, sunlamps, and risk of cutaneous malignant melanoma. *Cancer Epidemiol. Biomarkers Prev.* 14:562-566, 2005.
- Gandini, S., et al., Meta-analysis of risk factors for cutaneous melanoma: III. Family history, actinic damage and phenotypic factors. *Eur J Cancer*, 2005a.

- Gandini, S., et al., Meta-analysis of risk factors for cutaneous melanoma: I. Common and atypical naevi. *Eur J Cancer*, 2005b. **41**(1): p. 28-44.
- Gandini, S., et al., Meta-analysis of risk factors for cutaneous melanoma: II. Sun exposure. *Eur J Cancer*, 2005c. **41**(1): p. 45-60.
- Gambichler, T., A. Bader, M. Vojvodic, F. G. Bechara, K. Sauermann, P. Altmeyer and K. Hoffmann. Impact of UVA exposure on psychological parameters and circulating serotonin and melatonin. *BMC. Dermatol.* 2:6, 2002a.
- Gambichler, T., A. Bader, M. Vojvodic, A. Avermaete, M. Schenk, P. Altmeyer and K. Hoffmann. Plasma levels of opioid peptides after sunbed exposures. *Br. J. Dermatol.* 147:1207-1211, 2002b.
- Gange, R. W., A. D. Blackett, E. A. Matzinger, B. M. Sutherland and I. E. Kochevar. Comparative protection efficiency of UVA- and UVB-induced tans against erythema and formation of endonuclease-sensitive sites in DNA by UVB in human skin. *J. Invest Dermatol.* 85:362-364, 1985.
- Gilchrest, B. A., N. A. Soter, J. L. Hawk, R. M. Barr, A. K. Black, C. N. Hensby, A. I. Mallet, M. W. Greaves and J. A. Parrish. Histologic changes associated with ultraviolet A--induced erythema in normal human skin. *J. Am. Acad. Dermatol.* 9:213-219, 1983.
- Gilchrest, B. A., H. Y. Park, M. S. Eller and M. Yaar. Mechanisms of ultraviolet light-induced pigmentation. *Photochem. Photobiol.* 63:1-10, 1996.
- Glogau, R. G. Aesthetic and anatomic analysis of the aging skin. *Semin. Cutan. Med. Surg.* 15:134-138, 1996.
- Harrison, G. I. and A. R. Young. Ultraviolet radiation-induced erythema in human skin. *Methods* 28:14-19, 2002.
- Harrison, G. I., A. R. Young and S. B. McMahon. Ultraviolet radiation-induced inflammation as a model for cutaneous hyperalgesia. *J. Invest Dermatol.* 122:183-189, 2004.
- Hausser KW, Vahle W. Sunburn and suntanning. *Wiss Veröffnugen Siemens Konzern* 1927; **6**: 101–120
- Hausser KW, Vahle W. Die Abhängigkeit des Lichterythems und der Pigmentbildung von der Schwingungszahl (Wellenlänge) der erregenden Strahlung. *Strahlentherapie* 1928; **13**: 41
- Hawk, J. L., G. M. Murphy and C. A. Holden. The presence of neutrophils in human cutaneous ultraviolet-B inflammation. *Br. J. Dermatol.* 118:27-30, 1988.
- Health & Safety Executive, Controlling the health risks from the use of UV tanning equipment. *ING(G)209* 10/95, 1995
- Heller RF, Buchan I, Edwards R et al. Communicating risks at the population level: application of population impact numbers. *Br Med J* 2003; **327**: 1162-5

- 
- Hemminki, K., H. Zhang, and K. Czene, Familial and attributable risks in cutaneous melanoma: effects of proband and age. *J Invest Dermatol*, 2003. **120**(2): p. 217-23.
- Herity, B., G. O'Loughlin, M. J. Moriarty and R. Conroy. Risk factors for non-melanoma skin cancer. *Ir. Med. J.* 82:151-152, 1989.
- Hersey, P., M. MacDonald, C. Burns, S. Schibeci, H. Matthews and F. J. Wilkinson. Analysis of the effect of a sunscreen agent on the suppression of natural killer cell activity induced in human subjects by radiation from solarium lamps. *J. Invest Dermatol.* 88:271-276, 1987.
- Holick, M. F. The vitamin d epidemic and its health consequences. *J. Nutr.* 135:2739S-2748S, 2005.
- Holly, E.A., et al., Cutaneous melanoma in women. I. Exposure to sunlight, ability to tan, and other risk factors related to ultraviolet light. *Am J Epidemiol*, 1995. **141**(10): p. 923-33.
- IARC. Solar and ultraviolet radiation (International Agency for Research on Cancer, Lyon) (1992).
- ICNIRP International Commission on Non-Ionizing Radiation Protection. Health Issues of Ultraviolet Tanning Appliances Used for Cosmetic Purposes. *Health Physics* 2003; **84**: 119-127.
- IEC International Electrotechnical Commission. Safety of household and similar electrical appliances. Part 2: Particular requirements for appliances for skin exposure to ultraviolet and infrared radiation. Geneva:IEC, 1995, 335-2-27.
- Irwin, C., A. Barnes, D. Veres and K. Kaidbey. An ultraviolet radiation action spectrum for immediate pigment darkening. *Photochem. Photobiol.* 57:504-507, 1993.
- Johnson, G. J. The environment and the eye. *Eye* 18:1235-1250, 2004.
- Jose, J. G. and D. G. Pitts. Wavelength dependency of cataracts in albino mice following chronic exposure. *Exp. Eye Res.* 41:545-563, 1985.
- Kaddu, S., H. Kerl and P. Wolf. Accidental bullous phototoxic reactions to bergamot aromatherapy oil. *J. Am. Acad. Dermatol.* 45:458-461, 2001.
- Kadunce, D. P., M. W. Piepkorn and J. J. Zone. Persistent melanocytic lesions associated with cosmetic tanning bed use: "sunbed lentigines". *J. Am. Acad. Dermatol.* 23:1029-1031, 1990.
- Karagas, M. R., V. A. Stannard, L. A. Mott, M. J. Slattery, S. K. Spencer and M. A. Weinstock. Use of tanning devices and risk of basal cell and squamous cell skin cancers. *J. Natl. Cancer Inst.* 94:224-226, 2002.
- Kefford, R.F., et al., Counseling and DNA testing for individuals perceived to be genetically predisposed to melanoma: A consensus statement of the Melanoma Genetics Consortium. *J Clin Oncol*, 1999. **17**(10): p. 3245-51.



- 
- Kelly, D. A., A. R. Young, J. M. McGregor, P. T. Seed, C. S. Potten and S. L. Walker. Sensitivity to sunburn is associated with susceptibility to ultraviolet radiation-induced suppression of cutaneous cell-mediated immunity. *J. Exp. Med.* 191:561-566, 2000.
- Kligman, L. H. and R. M. Sayre. An action spectrum for ultraviolet induced elastosis in hairless mice: quantification of elastosis by image analysis. *Photochem. Photobiol.* 53:237-242, 1991.
- Kricker, A., B. K. Armstrong, D. R. English and P. J. Heenan. Pigmentary and cutaneous risk factors for non-melanocytic skin cancer--a case-control study. *Int. J. Cancer* 48:650-662, 1991.
- Kricker, A., B. K. Armstrong, D. R. English and P. J. Heenan. Does intermittent sun exposure cause basal cell carcinoma? a case-control study in Western Australia. *Int. J. Cancer* 60:489-494, 1995.
- Lavker, R. M., D. A. Veres, C. J. Irwin and K. H. Kaidbey. Quantitative assessment of cumulative damage from repetitive exposures to suberythemogenic doses of UVA in human skin. *Photochem. Photobiol.* 62:348-352, 1995a.
- Lavker, R. M., G. F. Gerberick, D. Veres, C. J. Irwin and K. H. Kaidbey. Cumulative effects from repeated exposures to suberythemal doses of UVB and UVA in human skin. *J. Am. Acad. Dermatol.* 32:53-62, 1995b.
- Lavker, R. and K. Kaidbey. The spectral dependence for UVA-induced cumulative damage in human skin. *J. Invest Dermatol.* 108:17-21, 1997.
- Ljunggren, B. Severe phototoxic burn following celery ingestion. *Arch. Dermatol.* 126:1334-1336, 1990.
- Luchiesh M. *Artificial Sunlight*, New York: Van Nostrand, 1930, p. 77.
- Man, I., I. K. Crombie, R. S. Dawe, S. H. Ibbotson and J. Ferguson. The photocarcinogenic risk of narrowband UVB (TL-01) phototherapy: early follow-up data. *Br. J. Dermatol.* 152:755-757, 2005.
- Martin JW, Chin JW, Nguyen T. Reciprocity law experiments in polymeric photodegradation: a critical review. *Progress in Organic Coatings* 2003; **47**: 292–311
- Matsumura, Y. and H. N. Ananthaswamy. Short-term and long-term cellular and molecular events following UV irradiation of skin: implications for molecular medicine. *Expert. Rev. Mol. Med.* 2002:1-22, 2002.
- McLoone, P., E. Simics, A. Barton, M. Norval and N. K. Gibbs. An action spectrum for the production of cis-urocanic acid in human skin in vivo. *J. Invest Dermatol.* 124:1071-1074, 2005.
- Meanwell EF, Diffey BL. Reciprocity of ultraviolet erythema in human skin. *Photodermatology* 1989; **6**: 146-148

- 
- Merriam, J. C., S. Lofgren, R. Michael, P. Soderberg, J. Dillon, L. Zheng and M. Ayala. An action spectrum for UV-B radiation and the rat lens. *Invest Ophthalmol. Vis. Sci.* 41:2642-2647, 2000.
- Morison WL, Towne LE and Honig B. The photoaggravated dermatoses. In: *Photodermatology*, Hawk JLM (ed) (London: Arnold, 1999, pp. 199-212 .
- Moyal, D. D. and A. M. Fourtanier. Efficacy of broad-spectrum sunscreens against the suppression of elicitation of delayed-type hypersensitivity responses in humans depends on the level of ultraviolet A protection. *Exp. Dermatol.* 12:153-159, 2003.
- Murphy, G. M., J. Wright, D. S. Nicholls, P. H. McKee, A. G. Messenger, J. L. Hawk and G. M. Levene. Sunbed-induced pseudoporphyria. *Br. J. Dermatol.* 120:555-562, 1989.
- Naldi, L., et al., Sunlamps and sunbeds and the risk of cutaneous melanoma. Italian Group for Epidemiological Research in Dermatology. *Eur J Cancer Prev*, 2000. **9**(2): p. 133-4.
- Noonan FP, DeFabo EC, Kripke ML. Suppression of contact hypersensitivity by UV radiation and its relationship to UV-induced suppression of tumor immunity. *Photochem Photobiol* 1981; **34**: 683.
- Noonan, F.P., et al., Animal models of melanoma: an HGF/SF transgenic mouse model may facilitate experimental access to UV initiating events. *Pigment Cell Res*, 2003. **16**(1): p. 16-25.
- Norris, W., A case of fungoid disease. *Edinb Med Surg J*, 1820. **16**: p. 562-565.
- Novakovic, L., S. Lee, G. E. Orchard, J. M. Sheehan, A. R. Young and S. L. Walker. Effects of solar-simulated radiation dose fractionation on CD1a+ Langerhans cells and CD11b+ macrophages in human skin. *Br. J. Dermatol.* 145:237-244, 2001.
- O'Loughlin, C., M. J. Moriarty, B. Herity and L. Daly. A re-appraisal of risk factors for skin carcinoma in Ireland. A case control study. *Ir. J. Med. Sci.* 154:61-65, 1985.
- Oriowo, O. M., A. P. Cullen, B. R. Chou and J. G. Sivak. Action spectrum and recovery for in vitro UV-induced cataract using whole lenses. *Invest Ophthalmol. Vis. Sci.* 42:2596-2602, 2001.
- Ortonne, J. P. The effects of ultraviolet exposure on skin melanin pigmentation. *J. Int. Med. Res.* 18 Suppl 3:8C-17C, 1990.
- Osterlind, A., et al., The Danish case-control study of cutaneous malignant melanoma. II. Importance of UV-light exposure. *Int J Cancer*, 1988. **42**(3): p. 319-24.
- Park YK, Gange RW, Levins PC, Parrish JA. Low and moderate irradiances of UVB and UVC irradiation are equally erythemogenic in human skin. *Photochem Photobiol* 1984; **40**: 667
- Parkin, D., et al., *Cancer Incidence in Five Continents*. IARC Scientific Publications No 143, Lyon, International Agency for Research on Cancer, 1997. **II**.

- 
- Parrish JA, Anderson RR, Ying CY, Pathak MA. Cutaneous effects of pulsed nitrogen gas laser irradiation. *J Invest Dermatol* 1976; **67**: 603
- Parrish, J. A., K. F. Jaenicke and R. R. Anderson. Erythema and melanogenesis action spectra of normal human skin. *Photochem. Photobiol.* 36:187-191, 1982.
- Pfeifer, G.P., Y.H. You, and A. Besaratinia, Mutations induced by ultraviolet light. *Mutat Res*, 2005. **571**(1-2): p. 19-31.
- Phan, T. A., G. M. Halliday, R. S. Barnetso and D. L. Damian. Spectral and dose dependence of ultraviolet radiation-induced immunosuppression. *Front Biosci.* 11:394-411, 2006.
- Pittelkow, M.R., et al., Skin cancer in patients with psoriasis treated with coal tar. A 25-year follow-up study. *Arch Dermatol*, 1981. **117**(8): p. 465-8.
- Pitts, D. G., A. P. Cullen and P. D. Hacker. Ocular effects of ultraviolet radiation from 295 to 365 nm. *Invest Ophthalmol. Vis. Sci.* 16:932-939, 1977.
- Ponsonby, A. L., R. M. Lucas and d. M. van, I. A potential role for UVR and Vitamin D in the induction of Multiple Sclerosis, Type 1 Diabetes, Rheumatoid Arthritis. *Photochem. Photobiol.* 2005.
- Rafnsson, V., et al., Risk factors for malignant melanoma in an Icelandic population sample. *Prev Med*, 2004. **39**(2): p. 247-52.
- Rigel, D.S., et al., Relationship of fluorescent lights to malignant melanoma: another view. *J Dermatol Surg Oncol*, 1983. **9**(10): p. 836-8.
- Rivers, J. K., P. G. Norris, G. M. Murphy, A. C. Chu, G. Midgley, J. Morris, R. W. Morris, A. R. Young and J. L. Hawk. UVA sunbeds: tanning, photoprotection, acute adverse effects and immunological changes. *Br. J. Dermatol.* 120:767-777, 1989.
- Roberts, J. E. Ocular phototoxicity. *J. Photochem. Photobiol. B* 64:136-143, 2001.
- Robinson, E.S., et al., The Monodelphis melanoma model: initial report on large ultraviolet A exposures of suckling young. *Photochem Photobiol*, 2000. **71**(6): p. 743-6.
- Routaboul, C., A. Denis and A. Vinche. Immediate pigment darkening: description, kinetic and biological function. *Eur. J. Dermatol.* 9:95-99, 1999.
- Ruegemer, J., B. Schuetz, K. Hermann, R. Hein, J. Ring and D. Abeck. UV-induced skin changes due to regular use of commercial sunbeds. *Photodermatol. Photoimmunol. Photomed.* 18:223-227, 2002.
- Salisbury, J. R., H. Williams and A. W. du Vivier. Tanning-bed lentigines: ultrastructural and histopathologic features. *J. Am. Acad. Dermatol.* 21:689-693, 1989.
- Sayre, R.M., J.C. Dowdy, and M. Poh-Fitzpatrick, Dermatological risk of indoor ultraviolet exposure from contemporary lighting sources. *Photochem Photobiol*, 2004. **80**: p. 47-51.

- 
- Schartl, A., et al., Analysis of genetic factors and molecular mechanisms in the development of hereditary and carcinogen-induced tumors of Xiphophorus. *Recent Results Cancer Res*, 1997. **143**: p. 225-235.
- Schieke, S.M., Schroeder P and Krutmann J. Cutaneous effects of infrared radiation: from clinical observations to molecular response mechanisms. *Photodermatol. Photoimmuno. Photomed.* 2003, 19: 228-234
- Schieke, S. M., K. Ruwiedel, H. Gers-Barlag, S. Grether-Beck and J. Krutmann. Molecular crosstalk of the ultraviolet a and ultraviolet B signaling responses at the level of mitogen-activated protein kinases. *J. Invest Dermatol.* 124:857-859, 2005.
- Schmidt K. Zur Hauterythemwirkung von UV-Blitzen. *Stahlentherapie* 1963; **124**: 127
- Setlow, R.B., et al., Wavelengths effective in induction of malignant melanoma. *Proc Natl Acad Sci USA*, 1993. **90**: p. 6666-6670.
- Sheehan, J. M. and A. R. Young. The sunburn cell revisited: an update on mechanistic aspects. *Photochem. Photobiol. Sci.* 1:365-377, 2002.
- Sliney, D. H. Photoprotection of the eye - UV radiation and sunglasses. *J. Photochem. Photobiol. B* 64:166-175, 2001.
- Stern, R. S. and W. Docken. An exacerbation of SLE after visiting a tanning salon. *JAMA* 255:3120, 1986.
- Stern, R.S. and N. Laird, The carcinogenic risk of treatments for severe psoriasis. Photochemotherapy Follow-up Study. *Cancer*, 1994. **73**(11): p. 2759-64.
- Swerdlow, A.J., et al., Fluorescent lights, ultraviolet lamps and risk of cutaneous melanoma. *British Medical Journal*, 1988. **297**: p. 647-650.
- Tangpricha, V., A. Turner, C. Spina, S. Decastro, T. C. Chen and M. F. Holick. Tanning is associated with optimal vitamin D status (serum 25-hydroxyvitamin D concentration) and higher bone mineral density. *Am. J. Clin. Nutr.* 80:1645-1649, 2004.
- The Sunbed Association. Code of practice for operators, Chesham: The Sunbed Association, 2004
- Veierod, M.B., et al., A prospective study of pigmentation, sun exposure, and risk of cutaneous malignant melanoma in women. *J Natl Cancer Inst*, 2003. **95**(20): p. 1530-8.
- Vieth, R. The role of vitamin D in the prevention of osteoporosis. *Ann. Med.* 37:278-285, 2005.
- Wachsmuth, R.C., et al., Heritability and gene-environment interactions for melanocytic nevus density examined in a U.K. adolescent twin study. *J Invest Dermatol*, 2001. **117**(2): p. 348-52.
- Walter, S.D., et al., The association of cutaneous malignant melanoma with the use of sunbeds and sunlamps. *Am J Epidemiol*, 1990. **131**(2): p. 232-43.

- 
- Walter, S.D., et al., The association of cutaneous malignant melanoma and fluorescent light exposure. *Am J Epidemiol*, 1992. **135**(7): p. 749-62.
- Wang, S.Q., et al., Ultraviolet A and melanoma: a review. *J Am Acad Dermatol*, 2001. **44**(5): p. 837-46.
- Weischer, M., et al., No evidence for increased skin cancer risk in psoriasis patients treated with broadband or narrowband UVB phototherapy: a first retrospective study. *Acta Derm Venereol*, 2004. **84**(5): p. 370-4.
- Weiss, J. and E. G. Jung. [Solarium pseudoporphyria. Report of a case and review of the literature]. *Hautarzt* 41:671-674, 1990.
- Westerdahl, J., et al., Use of sunbeds or sunlamps and malignant melanoma in Southern Sweden. 1994. **140**: p. 691-699.
- Westerdahl, J., et al., Risk of cutaneous malignant melanoma in relation to use of sunbeds: further evidence for UV-A carcinogenicity. *Br J Cancer*, 2000. **82**(9): p. 1593-9.
- Whiteman, D.C., et al., Melanocytic nevi, solar keratoses, and divergent pathways to cutaneous melanoma. *J Natl Cancer Inst*, 2003. **95**(11): p. 806-12.
- Whitmore, S. E. and W. L. Morison. The effect of suntan parlor exposure on delayed and contact hypersensitivity. *Photochem. Photobiol.* 71:700-705, 2000.
- Whitmore, S. E., W. L. Morison, C. S. Potten and C. Chadwick. Tanning salon exposure and molecular alterations. *J. Am. Acad. Dermatol.* 44:775-780, 2001.
- WHO World Health Organization. Artificial tanning sunbeds - risks and guidance. Geneva, 2003
- Woollons, A., C. Kipp, A. R. Young, C. Petit-Frere, C. F. Arlett, M. H. Green and P. H. Clingen. The 0.8% ultraviolet B content of an ultraviolet A sunlamp induces 75% of cyclobutane pyrimidine dimers in human keratinocytes in vitro. *Br. J. Dermatol.* 140:1023-1030, 1999.
- Young, A. R., R. H. Guy and H. I. Maibach. Laser Doppler velocimetry to quantify UV-B induced increase in human skin blood flow. *Photochem. Photobiol.* 42:385-390, 1985.
- Young, A. R. Chromophores in human skin. *Phys. Med. Biol.* 42:789-802, 1997.
- Young, A. R., C. A. Chadwick, G. I. Harrison, O. Nikaido, J. Ramsden and C. S. Potten. The similarity of action spectra for thymine dimers in human epidermis and erythema suggests that DNA is the chromophore for erythema. *J. Invest Dermatol.* 111:982-988, 1998.
- Young AR. Tanning devices – fast track to skin cancer? *Pigment Cell Res* 2004; **17**: 2-9
- Zanetti, R., et al., [A case-control study of melanoma of the skin in the province of Torino, Italy]. *Rev Epidemiol Sante Publique*, 1988. **36**(4-5): p. 309-17.
- Zhu, G., et al., A major quantitative-trait locus for mole density is linked to the familial melanoma gene CDKN2A: a maximum-likelihood combined linkage and association analysis in twins and their sibs. *Am J Hum Genet*, 1999. **65**(2): p. 483-92.

## 7. ACKNOWLEDGEMENTS

Members of the working group are acknowledged for their valuable contribution to this opinion. The members of the working group are:

### SCCP Members

Prof. J. Krutmann

Prof. V. Rogiers

Prof. T. Sanner

Dr. I.R. White                      chairman

### External experts

Prof. B. Diffey                      Regional Medical Physics Department, Newcastle General Hospital, UK

Prof. J.A. Newton Bishop        Genetic Epidemiology Division, St James's University Hospital, Leeds, UK

Prof. A.R. Young                    St John's Institute of Dermatology, University of London, UK

## 8. APPENDIX A: TECHNIQUE FOR DETERMINING THE ERYTHEMAL-WEIGHTED IRRADIANCE

The erythemal-weighted irradiance should be determined by means of a calibrated spectroradiometer. This type of measurement, known as spectroradiometry, enables the intensity of UV radiation on a wavelength-by-wavelength step to be recorded over all the wavelengths present in the sunbed emission spectrum. A spectroradiometer is an expensive, complex piece of optical equipment, which is found mainly in laboratories with a special interest in photobiology. It is important that users of spectroradiometers have their own standard lamps (either deuterium or tungsten) regularly calibrated by a national standards laboratory so that these can be used to provide an absolute spectral sensitivity calibration of the spectroradiometer.

The technique is to place the input optics of the spectroradiometer on top of the plastic sheet of the sunbed or at the recommended tanning distance from a sun canopy. The spectral irradiance is measured in equal wavelength steps (e.g. 5nm) from 280 to 400nm resulting in a set of numbers like those shown in column 2 in the table below. The numbers in column 3 represent the erythema action spectrum. This is the relative effectiveness of UV radiation of different wavelengths to cause erythema (or redness) in human skin 8-24 hours after exposure. The numbers in each row of columns 2 and 3 are multiplied together in column 4. All the numbers in column 4 are summed to give (in this example this comes to 0.108) and then multiplied by 5nm (the wavelength interval used in scanning the spectrum) to give the erythemal-weighted irradiance in  $W/m^2$ , which in this example is  $0.108 \times 5 = 0.54 W/m^2$ . This number should not exceed  $0.7 W/m^2$ .

nm	Spectral irradiance $W/m^2/nm$	erythema action spectrum	Erythemal weighted irradiance
280	0.00	1.00000	0.000
285	0.00	1.00000	0.000
290	0.00	1.00000	0.000
295	0.00	1.00000	0.000
300	0.00	0.64863	0.001
305	0.02	0.21979	0.004
310	0.08	0.07450	0.006
315	0.34	0.02520	0.008
320	0.95	0.00855	0.008
325	2.27	0.00290	0.007
330	4.52	0.00136	0.006
335	7.31	0.00115	0.008

## Opinion on Biological effects of ultraviolet radiation relevant to health with particular reference to sun beds for cosmetic purposes

---

340	10.04	0.00097	0.010
345	12.25	0.00081	0.010
350	14.12	0.00068	0.010
355	14.48	0.00058	0.008
360	13.45	0.00048	0.007
365	16.71	0.00041	0.007
370	9.26	0.00034	0.003
375	6.80	0.00029	0.002
380	4.57	0.00024	0.001
385	2.94	0.00020	0.001
390	2.11	0.00017	0.000
395	1.33	0.00015	0.000
400	0.84	0.00012	0.000
		<b>Sum</b>	<b>0.108</b>