



Scientific Committee on Emerging and Newly Identified Health Risks

SCENIHR

Possible effects of Electromagnetic Fields (EMF) on Human Health



The SCENIHR adopted this opinion at the 16th plenary of 21 March 2007 after public consultation

Possible effects of Electromagnetic Fields (EMF) on Human Health

About the Scientific Committees

Three independent non-food Scientific Committees provide the Commission with the scientific advice it needs when preparing policy and proposals relating to consumer safety, public health and the environment. The Committees also draw the Commission's attention to the new or emerging problems which may pose an actual or potential threat.

They are: the Scientific Committee on Consumer Products (SCCP), the Scientific Committee on Health and Environmental Risks (SCHER) and the Scientific Committee on Emerging and Newly Identified Health Risks (SCENIHR) and are made up of external experts.

In addition, the Commission relies upon the work of the European Food Safety Authority (EFSA), the European Medicines Evaluation Agency (EMA), the European Centre for Disease prevention and Control (ECDC) and the European Chemicals Agency (ECHA).

SCENIHR

Questions concerning emerging or newly-identified risks and on broad, complex or multi-disciplinary issues requiring a comprehensive assessment of risks to consumer safety or public health and related issues not covered by other Community risk- assessment bodies.

In particular, the Committee addresses questions related to potential risks associated with interaction of risk factors, synergic effects, cumulative effects, antimicrobial resistance, new technologies such as nanotechnologies, medical devices, tissue engineering, blood products, fertility reduction, cancer of endocrine organs, physical hazards such as noise and electromagnetic fields and methodologies for assessing new risks.

Scientific Committee members

Anders Ahlbom, James Bridges, Wim De Jong, Jana Hajslová, Philippe Hartemann, Thomas Jung, Mats-Olof Mattsson, Jean-Marie Pagès, Konrad Rydzynski, Dorothea Stahl, Mogens Thomsen, David Williams

Contact:

European Commission
Health & Consumer Protection DG
Directorate C: Public Health and Risk Assessment
Unit C7 - Risk Assessment
Office: B232 B-1049 Brussels

Sanco-Sc1-Secretariat@ec.europa.eu

© European Commission 2007

The opinions of the Scientific Committees present the views of the independent scientists who are members of the committees. They do not necessarily reflect the views of the European Commission. The opinions are published by the European Commission in their original language only.

http://ec.europa.eu/health/ph_risk/risk_en.htm

Possible effects of Electromagnetic Fields (EMF) on Human Health

ACKNOWLEDGMENTS

Members of the working group are acknowledged for their valuable contribution to this opinion. The members of the working group are:

SCENIHR members:

Prof. Anders Ahlbom (*Chair and Rapporteur*)

Prof. James Bridges

Prof. Mats-Olof Mattsson

External experts:

Dr. René de Seze, INERIS (National Institute for Environment and Industrial Risk), France

Lena Hillert, MD, PhD, Department of Public Health Sciences, Division of Occupational Medicine, Karolinska Institute and Department of Occupational and Environmental Health, Stockholm Centre for Public Health, Sweden

Prof. Jukka Juutilainen, Department of Environmental Sciences, University of Kuopio, Finland

Dr. Georg Neubauer¹, Business Unit Mobile Communications Safety, Austrian Research Centers GmbH - ARC, Austria

Dr. Joachim Schüz, Institute of Cancer Epidemiology, The Danish Cancer Society, Denmark

PD Dr. Myrtil Simko, Institut für Zellbiologie und Biosystemtechnik, Universität Rostock, Germany

¹ Declared Interest (see minutes of the SCENIHR plenary meeting of 28-29 September 2005: http://ec.europa.eu/health/ph_risk/committees/04_scenihr/docs/scenihr_mi_007.pdf).

Possible effects of Electromagnetic Fields (EMF) on Human Health

ABSTRACT

The Scientific Committee on Emerging and Newly Identified Health Risks (SCENIHR) has updated the previous opinion on "Possible effects of Electromagnetic Fields (EMF), Radio Frequency Fields (RF) and Microwave Radiation on human health" by the Scientific Committee on Toxicity, Ecotoxicity and the Environment (CSTEE) from 2001, with respect to whether or not exposure to electromagnetic fields (EMF) is a cause of disease or other health effects. The opinion is primarily based on scientific articles, published in English language peer-reviewed scientific journals. Only studies that are considered relevant for the task are cited and commented upon in the opinion. The opinion is divided into frequency (f) bands, namely: radio frequency (RF) ($100 \text{ kHz} < f \leq 300 \text{ GHz}$), intermediate frequency (IF) ($300 \text{ Hz} < f \leq 100 \text{ kHz}$), extremely low frequency (ELF) ($0 < f \leq 300 \text{ Hz}$), and static (0 Hz) (only static magnetic fields are considered in this opinion). There is a separate section for environmental effects.

Radio Frequency Fields (RF fields)

Since the adoption of the 2001 opinion extensive research has been conducted regarding possible health effects of exposure to low intensity RF fields, including epidemiologic, in vivo, and in vitro research. In conclusion, no health effect has been consistently demonstrated at exposure levels below the limits of ICNIRP (International Committee on Non Ionising Radiation Protection) established in 1998. However, the data base for evaluation remains limited especially for long-term low-level exposure.

Intermediate Frequency Fields (IF fields)

Experimental and epidemiological data from the IF range are very sparse. Therefore, assessment of acute health risks in the IF range is currently based on known hazards at lower frequencies and higher frequencies. Proper evaluation and assessment of possible health effects from long-term exposure to IF fields are important because human exposure to such fields is increasing due to new and emerging technologies.

Extremely low frequency fields (ELF fields)

The previous conclusion that ELF magnetic fields are possibly carcinogenic, chiefly based on occurrence of childhood leukaemia, is still valid. For breast cancer and cardiovascular disease, recent research has indicated that an association is unlikely. For neurodegenerative diseases and brain tumours, the link to ELF fields remains uncertain. No consistent relationship between ELF fields and self-reported symptoms (sometimes referred to as electrical hypersensitivity) has been demonstrated.

Static Fields

Adequate data for proper risk assessment of static magnetic fields are very sparse. Developments of technologies involving static magnetic fields, e.g. with MRI (Magnetic Resonance Imaging) equipment require risk assessments to be made in relation to occupational exposure.

Environmental Effects

There are insufficient data to identify whether a single exposure standard is appropriate to protect all environmental species from EMF. Similarly the data are inadequate to judge whether the environmental standards should be the same or significantly different from those appropriate to protect human health.

Keywords:

EMF, electromagnetic fields, radiofrequency fields, intermediate frequency fields, extremely low frequency fields, static fields, health effects, human health, environmental effects, SCENIHR, Scientific Committee on Emerging and Newly Identified Health Risks

Opinion to be cited as:

SCENIHR (Scientific Committee on Emerging and Newly Identified Health Risks). Possible effects of Electromagnetic Fields (EMF) on Human Health. 21 March 2007

Possible effects of Electromagnetic Fields (EMF) on Human Health

TABLE OF CONTENTS

ACKNOWLEDGMENTS	3
ABSTRACT	4
EXECUTIVE SUMMARY	6
1. BACKGROUND	8
2. TERMS OF REFERENCE	10
3. SCIENTIFIC RATIONALE	11
3.1. Introduction	11
3.2. Methods.....	12
3.3. Radio Frequency Fields (RF fields)	13
3.3.1. Sources and distribution of exposure in the population.....	13
3.3.2. Cancer	15
3.3.2.1. Epidemiology.....	15
3.3.2.2. In vivo	20
3.3.2.3. In vitro.....	21
3.3.3. Symptoms.....	23
3.3.4. Nervous system effects	25
3.3.5. Miscellaneous human	27
3.3.6. Reproduction and development.....	27
3.3.7. Sensitivity of children.....	27
3.3.8. Conclusions about RF fields	28
3.4. Intermediate Frequency Fields (IF fields)	29
3.4.1. Sources and distribution of exposure in the population.....	29
3.4.2. Health Effects	29
3.4.3. Conclusions about IF fields.....	30
3.5. Extremely low frequency fields (ELF fields)	30
3.5.1. Sources and distribution of exposure in the population.....	30
3.5.2. Cancer	31
3.5.2.1. Epidemiology.....	31
3.5.2.2. In vivo	32
3.5.2.3. In vitro.....	33
3.5.3. Symptoms.....	35
3.5.4. Other Health Effects.....	36
3.5.4.1. Epidemiology.....	36
3.5.4.2. In vivo	36
3.5.4.3. In vitro.....	37
3.5.5. Conclusions about ELF fields.....	37
3.6. Static fields	38
3.6.1. Sources and distribution of exposure in population	38
3.6.2. Health effects.....	38
3.6.3. Conclusions about static fields	38
3.7. Environmental Effects	38
4. OPINION	41
5. COMMENTS RECEIVED DURING THE PUBLIC CONSULTATION	45
6. MINORITY OPINION	47
7. LIST OF ABBREVIATIONS	48
8. REFERENCES	50
9. GLOSSARY	63

Possible effects of Electromagnetic Fields (EMF) on Human Health

EXECUTIVE SUMMARY

The Scientific Committee on Toxicity, Ecotoxicity and the Environment (CSTEE) presented an opinion on "Possible effects of Electromagnetic Fields (EMF), Radio Frequency Fields (RF) and Microwave Radiation on human health" in 2001. The SCENIHR has been asked to update this opinion and also to continuously monitor new information that may influence the assessment of risks to human health. In preparation for this update, scientific data, published since the previous opinion, have been reviewed and their impact on the conclusions of the previous opinion has been assessed. The main focus of the opinion is whether health effects might occur at exposure levels below those of established biological mechanisms and, in particular, in relation to long term exposure at such low levels. The present opinion is divided according to frequency band. A separate section discusses environmental effects.

Radio Frequency Fields (RF fields)

Since the adoption of the 2001 opinion extensive research has been conducted regarding possible health effects of exposure to low intensity RF fields, including epidemiologic, in vivo, and in vitro research.

The balance of epidemiologic evidence indicates that mobile phone use of less than 10 years does not pose any increased risk of brain tumour or acoustic neuroma. For long-term use, data are sparse, and the following conclusions are therefore uncertain and tentative. However, from the available data it does appear that there is no increased risk for brain tumours in long-term users, with the exception of acoustic neuroma for which there is some evidence of an association. For diseases other than cancer, very little epidemiologic data are available.

A particular consideration is mobile phone use by children. While no specific evidence exists, children or adolescents may be more sensitive to RF field exposure than adults. Children of today will also experience a much higher cumulative exposure than previous generations. To date no epidemiologic studies on children are available.

RF field exposure has not convincingly been shown to have an effect on self-reported symptoms or well-being.

Studies on neurological effects and reproductive effects have not indicated any health risks at exposure levels below the ICNIRP-limits established in 1998.

Animal studies have not provided evidence that RF fields could induce cancer, enhance the effects of known carcinogens, or accelerate the development of transplanted tumours. The open questions include adequacy of the experimental models used and scarcity of data at high exposure levels.

There is no consistent indication from in vitro research that RF fields affect cells at the nonthermal exposure level.

The technical development is very fast and sources of RF field exposure become increasingly common. Yet, there is a lack of information on individual RF field exposure and the relative contribution of different sources to the overall exposure.

In conclusion, no health effect has been consistently demonstrated at exposure levels below the ICNIRP-limits established in 1998. However, the data base for this evaluation is limited especially for long-term low-level exposure.

Intermediate Frequency Fields (IF fields)

Experimental and epidemiological data from the IF range are very sparse. Therefore, assessment of acute health risks in the IF range is currently based on known hazards at

Possible effects of Electromagnetic Fields (EMF) on Human Health

lower frequencies and higher frequencies. Proper evaluation and assessment of possible health effects from long term exposure to IF fields are important because human exposure to such fields is increasing due to new and emerging technologies.

Extremely low frequency fields (ELF fields)

The previous conclusion that ELF magnetic fields are possibly carcinogenic, chiefly based on childhood leukaemia results, is still valid. There is no generally accepted mechanism to explain how ELF magnetic field exposure may cause leukaemia.

For breast cancer and cardiovascular disease, recent research has indicated that an association is unlikely. For neurodegenerative diseases and brain tumours, the link to ELF fields remains uncertain. A relation between ELF fields and symptoms (sometimes referred to as electromagnetic hypersensitivity) has not been demonstrated.

Static Fields

Adequate data for proper risk assessment of static magnetic fields are very sparse. Developments of technologies involving static magnetic fields, e.g. with MRI equipment require risk assessments to be made in relation to the exposure of personnel.

Environmental Effects

The continued lack of good quality data in relevant species means that there are insufficient data to identify whether a single exposure standard is appropriate to protect all environmental species from EMF. Similarly the data are inadequate to judge whether the environmental standards should be the same or significantly different from those appropriate to protect human health.

Research Recommendations

Important research needs were identified within all frequency bands.

Possible effects of Electromagnetic Fields (EMF) on Human Health

1. BACKGROUND

For the general public, Council Recommendation of 12 July 1999² on the limitation of exposure of the general public to electromagnetic fields (0 Hz to 300 GHz) fixes basic restrictions and reference levels to electromagnetic fields (EMFs). These restrictions and reference levels are based on the guidelines published by the International Commission on Non Ionising Radiation Protection (ICNIRP)³. The ICNIRP guidelines had been endorsed by the Scientific Steering Committee (SSC)⁴ in its opinion on health effects of EMFs of 25–26 June 1998⁵.

For workers, the Council and the Parliament have adopted Directive 2004/40/EC of 29 April 2004⁶ on the minimum health and safety requirements regarding the exposure of workers to the risks arising from physical agents (EMFs).

In a questionnaire sent to Member States in 2000, all have notified the Commission that they have implemented the provisions of Council Recommendation on the limitation of exposure of the general public to EMFs. The position of the new member states has not yet been ascertained.

The Commission has announced that it intends to prepare a report to the Council on the implementation of the Recommendation, taking account of an earlier report of 2002 on implementation by the then member states.⁷

The Scientific Committee on Toxicity, Ecotoxicity and the Environment (CSTEE) was requested to prepare an update of the Scientific Steering Committee's opinion. The request derived from the increasing exposure to EMF consequent to the further growth in the use of electricity, from the continuous development of the telecommunications industry, and to a rapid increase in the installation of transmitter masts used as radiotelephone base stations. In addition to domestic, industrial and medical electrical appliances and devices, the high voltage overhead transmission lines (and to a lesser extent underground cables) are major sources of exposure to Extremely Low Frequencies (ELF) in the environment. The CSTEE opinion "on Possible effects of Electromagnetic Fields (EMF), Radio Frequency Fields (RF) and Microwave Radiation on human health"⁸, of 30 October 2001, concluded that the information that had become available since the SSC opinion of June 1999 did not justify revision of the exposure limits recommended by the Council⁹.

A substantial number of scientific publications and reviews on the possible health effects of EMF (focusing mostly on mobile telephones) have become available since the CSTEE

² <http://europa.eu.int/eur-lex/lex/LexUriServ/LexUriServ.do?uri=CELEX:31999H0519:EN:HTML>, OJ L 199/59, 30.7.1999

³ <http://www.icnirp.de/>

⁴ http://ec.europa.eu/food/fs/sc/ssc/index_en.html

⁵ http://ec.europa.eu/food/fs/sc/ssc/out19_en.html

⁶ [http://europa.eu.int/eur-lex/lex/LexUriServ/LexUriServ.do?uri=CELEX:32004L0040R\(01\):EN:HTML](http://europa.eu.int/eur-lex/lex/LexUriServ/LexUriServ.do?uri=CELEX:32004L0040R(01):EN:HTML), OJ L 184/1, 24.5.2004

⁷ http://ec.europa.eu/health/ph_determinants/environment/EMF/Implement_rep_en.pdf

⁸ http://ec.europa.eu/health/ph_risk/committees/sct/documents/out128_en.pdf

⁹ The main frequencies in the ELF frequency range are 50 Hz in Europe and 60 Hz in North America. The RF and lower microwave frequencies are of particular interest for broadcasting, mobile telephony. The 2.45 GHz frequency is mainly used in domestic and industrial microwave ovens.

Possible effects of Electromagnetic Fields (EMF) on Human Health

opinion of 2001, for example the 2002 Dutch report¹⁰, the 2003 AGNIR report¹¹ and the 2004 British National Radiological Protection Board (NRPB) report on "Mobile phones and health"¹², which is the most recent of them. The NRPB provided a detailed review of the recent literature and useful contribution to the discussions on whether there are health effects related to the use of mobile phones. The report concluded that there is no hard evidence at present that the health of the public is being adversely affected by mobile phone technologies but uncertainties remain and a continued precautionary approach is recommended until the situation is clarified.

Additional results are expected shortly from Community funded research and development (R&D) activities, from national programmes, and from work within the International EMF Project of the World Health Organisation (WHO).

Community funded R&D comprises direct support to the Joint Research Centre and indirect support to competitive projects under the 5th Framework Programme¹³ and the 6th Framework Programme (FP6)¹⁴ for Research and Technological Development. Under FP6, the EMF-NET Coordination Action¹⁵ brings together European and national EMF programmes. EMF-NET will start publishing its first interpretation reports at the end of 2005. In total, this project will run another three years.

As part of its mission to protect public health and in response to public concern over health effects of EMF exposure, WHO established the International EMF Project¹⁶ in 1996 to assess the scientific evidence of possible health effects of EMF in the frequency range from 0 to 300 GHz. The EMF Project encourages focused research to fill important gaps in knowledge and to facilitate the development of internationally acceptable standards limiting EMF exposure.

In view of the amount of scientific evidence that has become available since the publication of the CSTEE of 30 October 2001 and of the Commission's intention to prepare a report to the Council and the frequency of new scientific publications on the health effects on EMF which require rapid assessment, the SCENIHR is asked to both update the previous opinion of the CSTEE and to continuously monitor new information that may influence the assessment of risks to human health in this area.

¹⁰ Mobile telephones – evaluation of health effects, Report of the Health Council of the Netherlands, 28.1.2002, <http://www.gr.nl>

¹¹ AGNIR (2003). Health effects from radiofrequency electromagnetic fields. Report of an Advisory Group on Non- ionising Radiation. Doc NRPB, 14(29), 1-177. Available at <http://www.nrpb.org>

¹² National Radiological Protection Board (NRPB) (2004). Mobile Phones and Health 2004, 15(5). Available at http://www.hpa.org.uk/web/HPAwebFile/HPAweb_C/1194947333240

¹³ <http://ec.europa.eu/research/fp5.html> and <http://cordis.europa.eu/fp5/home.html>.

¹⁴ http://ec.europa.eu/research/fp6/index_en.cfm and <http://cordis.europa.eu/fp6/dc/index.cfm?fuseaction=UserSite.FP6HomePage> .

¹⁵ <http://www.jrc.ec.europa.eu/eis-emf/emfnet.cfm>

¹⁶ <http://www.who.int/peh-emf/project/en/>

2. TERMS OF REFERENCE

The Committee is requested:

- (1) to update the CSTEE opinion of 30 October 2001 by summer 2006 in the light of newly available information;
- (2) to monitor the scientific literature concerning the health effects of EMF;
- (3) to draw the Commission's attention to significant new scientific findings;
- (4) to provide the Commission with an annual review of the opinion in the light of significant new evidence;
- (5) to take full advantage of the periodic reviews undertaken under the auspices of EMF-NET.

In reviewing and evaluating the studies on the potential health effects of EMF, the Committee is asked to pay particular attention to

- the nature of EMF studies, i.e., epidemiology, laboratory biology (in vivo vs. in vitro), clinical examinations (heart function, sleep electrophysiology, immune system, blood chemistry, hormones including melatonin, etc.), and theory;
- the methodology of EMF studies, in particular, epidemiology (e.g., "background health condition", "odds ratio", and the problem of ubiquitous technologies), measurement (cf., spot measurement, time-weighted average, personal monitor, calculated historical fields, laboratory measurement, etc.), and combined exposures (from different EMF sources as well as from simultaneous exposure to EMF and other factors such as chemicals, noise, stress, etc.);
- the characterization of risks, in particular, nature and magnitude of damage, likelihood of occurrence (expressed preferably in terms of natural frequencies rather than probabilities), uncertainty, geographical distribution, persistence over time, reversibility, delay, possible violation of equity, potential for public mobilization etc.; and
- the identification and physical characterization of existing and foreseeable sources of exposure to EMF, e.g., electromagnetic vs. magnetic including magnetic resonance imagery (MRI), from AC vs. DC current, new frequency ranges, higher transmission power, etc.

3. SCIENTIFIC RATIONALE

3.1. Introduction

The purpose of the opinion is to update the CSTE opinion of 2001 with respect to whether or not exposure to electromagnetic fields (EMF) is a cause of disease or other health effects and the purpose is not to provide a general review on electromagnetic fields and health. Recommendations regarding exposure guidelines or other risk management tools, including application of the precautionary principle are beyond the scope of the opinion. The methods that were used for the preparation of the opinion are explained below.

The objective of this section is to establish the scientific rationale that is necessary in order to provide an opinion in response to the request to the Committee, in particular to update the CSTE opinion of 30 October 2001. This section therefore summarizes what was known at the time of the 2001 Opinion, reviews the scientific data that have been published after 2001, and assesses to what extent these new data affect previous conclusions. Following the Committee's general principles, only studies published in peer reviewed journals have been considered.

The section is divided into four sub-sections according to frequency (f) range: radio frequency (RF) ($100 \text{ kHz} < f \leq 300 \text{ GHz}$), intermediate frequency (IF) ($300 \text{ Hz} < f \leq 100 \text{ kHz}$), extremely low frequency (ELF) ($0 < f \leq 300 \text{ Hz}$), and static (0 Hz) (only static magnetic fields are considered in this opinion). These frequency ranges are discussed in order of decreasing frequency, RF, IF, ELF, and static. For each frequency range the review begins with a description of sources and exposure to the population. This is followed, for each frequency range, by a discussion that is organized according to outcome. For each outcome relevant human, in vivo, and in vitro data are covered.

It is well recognized that there are established biophysical mechanisms that can lead to health effects as a consequence of exposure to sufficiently strong fields. For frequencies up to, say, 100 kHz the mechanism is stimulation of nerve and muscle cells due to induced currents and, for higher frequencies, tissue heating is the main mechanism. These mechanisms lead to acute effects. Existing exposure guidelines, such as those issued by ICNIRP, protect against these effects. The current issue is the possibility that health effects occur at exposure levels below those where the established mechanisms play a role and in particular as effects of long term exposure at low level. No further consideration is given to thermal effects.

Table 1 below illustrates some typical artificial sources of electromagnetic fields with frequency and intensity. Natural sources like the magnetic field of the earth are not included. Note, however, that big variations occur. For an explanation of some of the terminology used please be referred to the next chapter.

Possible effects of Electromagnetic Fields (EMF) on Human Health

Table 1. Typical sources of electromagnetic fields

Frequency range	Frequencies	Some examples of exposure sources
Static	0 Hz	VDU (video displays); MRI and other diagnostic / scientific instrumentation; Industrial electrolysis; Welding devices
ELF	0-300 Hz	Powerlines; Domestic distribution lines, Domestic appliances; Electric engines in cars, train and tramway; Welding devices
IF	300 Hz – 100 kHz	VDU; anti theft devices in shops, hands free access control systems, card readers and metal detectors; MRI; Welding devices
RF	100 kHz – 300 GHz	Mobile telephony; Broadcasting and TV; Microwave oven; Radar, portable and stationary radio transceivers, personal mobile radio; MRI

The Committee has been made aware of the military use of certain radiofrequency devices. Further consideration of this is outside the scope of this opinion.

3.2. Methods

As a general rule, scientific reports that are published in English language peer-reviewed scientific journals are considered primarily. This does not imply that all published articles are considered to be equally valid and relevant for health risk assessment. On the contrary, a main task is to evaluate and assess the articles and the scientific weight that is to be given to each of them. Only studies that are considered relevant for the task are commented upon in the opinion. Many more reports were considered than are cited in the reference list. However, only articles that contribute significantly to the update of the opinion are cited and commented upon. In some areas where the literature is particularly scarce it has been considered important to explain why the results of certain studies do not add useful information to the data base. The focus is on articles presented after the year 2000.

Relevant research for EMF health risk assessment can be divided into broad sectors such as epidemiologic studies, experimental studies in humans, experimental studies in animals, and cell culture studies. Also studies on biophysical mechanisms, dosimetry, and exposure assessment are considered. In a report of this nature it is not possible to consider the experiences of individuals. Nonetheless, such information often triggers a scientific study.

A health risk assessment evaluates the evidence within each of these sectors and then weighs together the evidence across the sectors to a combined assessment. This combined assessment should address the question of whether or not a hazard exists i.e., if there exists a causal relation between exposure and some adverse health effect. The answer to this question is not necessarily a definitive yes or no, but may express the weight of the evidence for the existence of a hazard. If such a hazard is judged to be present, the risk assessment should also address the magnitude of the effect and the shape of the dose-response function, i.e., the magnitude of the risk for various exposure levels and exposure patterns. A full risk assessment also includes exposure characterization in the population and estimates of the impact of exposure on burden of disease.

Epidemiological and experimental studies are subject to similar treatment in the evaluation process. It is of equal importance to evaluate positive and negative studies, i.e., studies indicating that EMF has an effect and studies not indicating the existence of

Possible effects of Electromagnetic Fields (EMF) on Human Health

such an effect. In the case of positive studies the evaluation focuses on alternatives to causation as explanation to the positive result: With what degree of certainty can one rule out the possibility that the observed positive result is produced by bias, e.g. confounding or selection bias, or chance. In the case of negative studies one assesses the certainty with which it can be ruled out that the lack of an observed effect is the result of (masking) bias, e.g., because of too small exposure contrasts or too crude exposure measurements; one also has to evaluate the possibility that the lack of an observed effect is the result of chance, a possibility that is a particular problem in small studies with low statistical power. Obviously, the presence or absence of statistical significance is only one factor in this evaluation. Rather, the evaluation considers a number of characteristics of the study. Some of these characteristics are rather general, such as study size, assessment of participation rate, level of exposure, and quality of exposure assessment. Particularly important aspects are the observed strength of association and the internal consistency of the results including aspects such as dose response relation. Other characteristics are specific to the study in question and may involve dosimetry, method for assessment of biological or health endpoint, the relevance of any experimental biological model used etc. Regarding experimental studies, additional important characteristics that are taken into consideration are the types of controls that have been used and to what degree replication studies have been performed. For a further discussion of aspects of study quality, refer for example to the Preamble to the IARC Monograph Series (IARC 2006). It is worth noting that the result of this process is not an assessment that a specific study is unequivocally negative or positive or whether it is accepted or rejected. Rather, the assessment will result in a weight that is given to the findings of a study

The step that follows the evaluation of the individual studies within a sector of research is the assessment of the overall evidence from that sector with respect to a given outcome. This implies integrating the results from all relevant individual studies into a total assessment. This is based on the evaluations of the individual studies and takes into account, for each study, both the observed magnitude of the effect and the quality of the study. Note again, that for this process to be valid, all studies must be considered equally irrespective of their outcome.

In the final overall evaluation phase, the available evidence is integrated over various sectors of research. This phase involves combining the existing relevant pieces of evidence on a particular end-point from studies in humans, from animal models, in vitro studies, and from other relevant areas. The integration of the separate lines of evidence should take place as the last, overall evaluation stage, after the critical assessment of all (relevant) available studies for particular end-points. In the first phase, epidemiological studies should be critically evaluated for quality irrespective of the putative mechanisms of biological action of a given exposure. In the final integrative stage of evaluation, however, the plausibility of the observed or hypothetical mechanism(s) of action and the evidence for that mechanism(s) is a factor to be considered. The overall result of the integrative phase of evaluation, combining the degree of evidence from across epidemiology, animal studies, in vitro and other data depends on how much weight is given on each line of evidence from different categories.

3.3. Radio Frequency Fields (RF fields)

3.3.1. Sources and distribution of exposure in the population

Nowadays the use of RF sources is widespread in our society. Prominent examples are mobile communication, broadcasting or medical and industrial applications. Information on emissions arising from RF sources is often available and can be used for compliance assessment or similar applications such as in-situ measurements. It has to be taken into account that information on the exposure of individual persons is scarce; such

Possible effects of Electromagnetic Fields (EMF) on Human Health

information is mainly needed for epidemiological studies, there is therefore a need to optimize methodology to assess individual exposure, e.g. by using and further developing existing dosimeters. The existing RF sources are operated in different frequency bands and can be subdivided in several categories.

Sources operated close to the human body

Many devices of this type are mobile RF transmitters. One of the examples is mobile phones; more than 2 billion people are using mobile phones worldwide. The most common mobile communication technologies in Europe are the digital technologies GSM 900, GSM 1800 and UMTS, analogue technologies are nowadays almost not in use any longer in Europe. Mobile phone use is common in Europe and the proportion of users can reach values of 80 % or more. Before mobile phones can be brought into the European market they have to show compliance with the requirements of European directives, i.e., it has to be shown that the limits for the amount of power absorbed in the human body are not exceeded. Standardized methods specified by the European Committee for Electrical Standardisation (CENELEC) are used to test mobile phones in Europe. The limit for mobile phone use is the specific absorption rate (SAR) of 2 W/kg for the human head. Mobile phones are tested under worst case conditions, i.e. at the highest power level, e.g., 2 W peak power corresponding to 250 mW maximum time averaged transmitted power for GSM at 900 MHz. Maximum local SAR values averaged over 10 gram of tissue range typically between 0.2 and 1.5 W/kg, depending on the type of mobile phone. It has to be taken into account that the emitted power is often orders of magnitude lower than the maximum power leading to much lower exposure due to power control and discontinuous transmission mode for GSM and UMTS phones. The power control of a GSM phone automatically reduces the emitted power by up to a factor of 1,000 for GSM and about 100.000.000 for UMTS if the intensity is not needed for stable transmission. No exposure occurs from a mobile phone being switched off. Phones operated in the standby mode cause typically much lower exposure compared to mobile phones operated with maximum power, but an accurate figure for this lower exposure depends on the exact details of the transmission path to base stations and on the traffic requested by the communication protocol and by incoming / outgoing SMS.

In addition to mobile phones, other wireless applications like cordless phones, e.g. DECT, or WLAN systems are very common. Due to the fact that they are usually operated with lower output power compared to mobile phones the exposure is typically below the level of mobile phones. The maximum time averaged power level of a DECT base station is 250 mW (worst case for a professional application handling communication with 25 handsets in parallel, a typical household application communicating with one handset has a time averaged power of 10 mW), for a DECT handset 10mW. The peak value of a WLAN terminal is 200 mW, however the averaged power depends on the traffic and is usually considerable lower. The exposure from such systems is therefore typically below that of mobile phones, however under certain circumstances, e.g. closeness to WLAN access points, exposure due to WLAN or DECT systems can become superior compared to the exposure from GSM or UMTS mobile phones. For example, close to a WLAN system exposure is typically below 0.5 mW/m². Anti-theft devices have become more and more common during recent years. They are typically operated at the exits of shops or similar areas to prevent theft of goods. Some of the existing systems are operated in the RF range; the exposure depends on the type of system and is, as long as the systems are operated according to the manufacturer's requirements, below the exposure limits. Several industrial appliances are operated in the RF and microwave range, for example for heating (e.g. RF sealers) or maintenance of broadcasting stations. The exposure of the worker operating such systems can reach values close or even above the limits of the Directive 2004/40/EC.

Sources operated far away from the human body

Such sources are typically fixed installed RF transmitters. An example is base stations that are an essential part of mobile communication networks necessary to establish the

Possible effects of Electromagnetic Fields (EMF) on Human Health

link between the mobile telephone and the rest of the network. In most European countries, base stations have become ubiquitous to guarantee connectivity in large areas of the respective countries; e.g., 18,000 base stations are operated in Austria. The so called reference level for the exposure of the general population at 900 MHz, an important frequency for mobile communication given in the European Council Recommendation 1999/519/EC is 4.5 W/m^2 . This reference level is given as an example, it has to be taken into account that the reference levels are frequency dependent and that other limits have to be applied at other frequencies. The range of exposure of the general population due to GSM signals is typically between some hundred nW/m^2 and some tens of mW/m^2 . The reasons for this large variation are both technical and environmental factors including distance. For UMTS, the available measurements are limited and so far the traffic is rather low compared to GSM. Values slightly over 1 mW/m^2 have been measured in a few cases, while minimum levels are a few hundred nW/m^2 . Other important RF sources are broadcasting systems (AM and FM). The maximum values measured in areas accessible for the public are typically below 10 mW/m^2 . Close to the fences of very powerful transmitters, exposure of about 300 mW/m^2 can be expected in some cases. Looking at the new digital TV technology (DVB-T), exposures between around 40 mW/m^2 and 0.003 mW/m^2 were registered in an Austrian study. The range of exposure is similar compared to analogue TV systems. However, the digital systems require more transmitters than the older analogue systems; therefore somewhat higher average exposure levels can be expected. In some countries digital audio broadcasting systems are already in operation. Other examples of sources relevant for far field exposure of the general population are civil and military radar systems, private mobile radio systems, or new technologies like WiMAX.

Medical applications

Several medical applications use electromagnetic fields in the RF range. Therapeutic applications such as soft tissue healing appliances, hyperthermia for cancer treatment, or diathermy expose the patient well above the recommended limit values to achieve the intended biological effects. These include heating of tissue (analgetic applications) or burning cells (to kill cancer cells). In these cases exposure of therapists or other medical personnel needs to be controlled to avoid that their exposure exceeds the exposure limit values foreseen by Directive 2004/40/EC for occupational exposure. Diagnostic applications, like magnetic resonance imaging (MRI), are allowed to exceed the basic restrictions of Council Recommendation 1999/519/EC as there is a benefit for the patient. Usual frequencies are those allowed for industrial, scientific, and medical applications similar to most industrial sources: 27 MHz, 433 MHz and 2.45 GHz. Magnetic resonance imaging devices in medical diagnostics use RF fields in addition to static and variable fields. Most actual clinical MRI devices work at 63 MHz.

3.3.2. Cancer

Studies on cancer in relation to mobile telephony have focused on intracranial tumours because deposition of energy from RF fields from a mobile phone is mainly within a small area of the skull near the handset. When whole body exposure is considered, as in some occupational and environmental studies, also other forms of cancer have been investigated.

3.3.2.1. Epidemiology

What was already known on this subject?

At the time of the previous CSTE opinion of 2001, most epidemiological studies on exposure to RF fields had examined exposures at the workplace. The overall evidence did not suggest consistent cancer excesses. With regard to mobile phones, only few studies were available at the time of the previous opinion and the short exposure period in these

Possible effects of Electromagnetic Fields (EMF) on Human Health

studies did not allow any firm conclusions. The few studies on residential exposure to RF fields from transmitters had serious methodological limitations.

What has been achieved since then?

In total, about 30 papers of original studies on mobile phone use and cancer were published in the last five years. Results are summarized in Table 2 for brain tumours and in Table 3 for acoustic neuroma. All but one study were case-control studies, mostly on brain tumours, some on salivary gland tumours or uveal melanoma. One was a large cohort study of all Danish mobile phone subscribers between 1982 and 1995 who were followed up for a variety of cancers; no increased risk for any cancer was observed but follow up time was short (Johansen et al. 2001). A recent update of the cohort study with an average follow up time of 8.5 years yielded 14,249 cancer cases observed in the cohort versus 15,001 expected cases based on cancer rates observed in the rest of the Danish adult population (Schüz et al. 2006b). The deficit was mainly attributable to smoking-related cancers, suggesting a healthy cohort effect. The overall relative risk estimates for brain tumours and leukaemia were close to one, however, only 28 brain tumour cases occurred in subscribers of a mobile phone of 10 years or more, whereas 42.5 cases were expected.

The Interphone study is a multinational case-control study coordinated by the International Agency for Research on Cancer (IARC). It is a population-based study with prospective ascertainment of incident cases and face-to-face interviews for exposure assessment. With regard to brain tumours, results from the first four components of the Interphone study suggest no risk increase for meningioma or glioma. This is consistently so among subjects with less than 10 years of use. For regular mobile phone users of 10 years or more, no indications of risk increases were seen in three out of four components, namely in Sweden (Lönn et al. 2005), Denmark (Christensen et al. 2005) and the UK (Hepworth et al. 2006), but the German component does reveal a somewhat raised relative risk estimate for glioma (Schüz et al. 2006a). This increase, however, is based on small numbers and due to the wide confidence interval the result is not in contradiction with the other Interphone components.

In contrast, a Swedish group not participating in the Interphone-study, conducting several case-control studies using self-administered questionnaires for exposure assessment, has repeatedly observed increased relative risk estimates for brain tumours. In 2006, the group revisited their previously published studies and reported statistically significant risk increases for both analogue and digital mobile phones as well as cordless phones already after one year of use (Hardell et al. 2006). After ten years of use they observed about a doubling of the relative risk estimates, with the strongest increase for high grade glioma.

Acoustic neuromas, benign tumours that develop very slowly, arise from the Schwann cells, which enfold the vestibulocochlear nerve (VIII. cranial nerve). They are of particular interest because of their location. The Hardell-group from Sweden has in several studies reported raised relative risk estimates for acoustic neuroma, also with very short induction periods (Hardell et al. 2005b). Three of the Interphone components, Denmark, Sweden, and Japan, have reported their country specific acoustic neuroma results (Christensen et al. 2004, Lönn et al. 2004, Takebayashi et al. 2006). Lönn et al. (2004) reported a doubling of the relative risk estimate after ten years of regular mobile phone use compared to subjects who never used a mobile phone regularly. This association became stronger when the analysis was restricted to preferred phone use at the same side as the tumour. Christensen's and Takebayashi's results did not support this, but they were based on fewer long-term users. Five of thirteen countries of the Interphone study (including Denmark, Finland, Norway, Sweden, and the UK) were pooled for a joint analysis to examine the association between mobile phone use and risk of acoustic neuroma (Schoemaker et al. 2005). While no overall association was seen among all long-term users (see Table 3), the data suggest that there may be an

Possible effects of Electromagnetic Fields (EMF) on Human Health

increased risk when the preferred side of the head of use is considered in the analysis. For 10+ years of use of mobile phones, the relative risk for acoustic neuroma at the preferred side of use was 1.8 (95%-CI 1.1-3.1). Because of methodological inter-study differences it would have been of considerable interest to compare the results across the six studies, but small numbers in most of the centres preclude that analysis.

All those studies are facing limitations in their exposure assessment, which was either a list of subscribers from the operators or self-reported mobile phone use. While the first method is an objective measure, it has limitations because subscription predicts use of a mobile phone only to some extent. Recent validation studies in volunteers comparing current self-reported use with traffic records from network operators show a moderate agreement, but it cannot be excluded that agreement is worse with respect to past mobile phone use or among patients with brain tumours (Vrijheid et al. 2006). Especially patients with high stage glioma showed some memory performance problems in the Danish Interphone study (Christensen et al. 2005). What seems to be reassuring despite these shortcomings is, that once the amount of mobile phone use is estimated with some validity, this is a satisfactory proxy for RF field exposure from these devices, as was shown in studies recording output power of mobile phones during operation (Berg et al. 2005). Laterality (side) of use is not easy to obtain in a retrospective study, as early symptoms may affect the side of use. Although some results are now available for long-term users, their numbers are still small and the results of the whole Interphone dataset should be awaited before drawing conclusions.

No striking new results appeared for studies on occupational and residential RF fields exposures since the previous opinion. While some positive associations have been reported from occupational studies, the overall picture is far from clear (Ahlbom et al. 2004). Many studies lack individual exposure assessment and only job titles or branches were used as exposure proxies. Studies on exposure from transmitters are limited by crude exposure measures and small numbers of exposed subjects, and the ecological nature of most studies.

Discussion

Mobile phones in relation to health are now being studied with great effort and in comprehensive studies, particularly in the Interphone Study. The results of the Interphone Study will soon become available. It has to be doubted, however, that the results will be entirely conclusive, as the first results from published national components of this study already raise a number of questions with respect to the potential of bias. Another limitation is that also in the current studies, long-term mobile phone users have had hardly more than 10 years of regular use of mobile phones, which still may be a relatively short latency period, particularly for slowly growing benign tumours. Among those long-term users, most were initially users of analogue mobile phone and thus, the number of long-term users of the digital technology is even smaller. Prospective long term follow up studies overcome both the limitations of retrospective exposure assessment and the latency problem and are recommended as a powerful long-term surveillance system for a variety of potential endpoints, including cancer, to fill current gaps in knowledge.

Possible effects of Electromagnetic Fields (EMF) on Human Health

Table 2. Results of epidemiological studies on mobile phone use and brain tumours¹⁷.

Study	Brain tumours		Brain tumours short latency		Brain tumours longer latency	
	Number exposed cases	RR ¹⁸ estimate (95% CI ¹⁹)	Number exposed cases	RR estimate (95% CI)	Number exposed cases	RR estimate (95% CI)
Hardell et al. 1999	78	1.0 (0.7-1.4)	78	1.0 (0.7-1.4) >1 yr	34	0.8 (0.5-1.4) >5 yr
					16	1.2 (0.6-2.6) >10 yr
Muscat et al. 2000	66	0.8 (0.6-1.2)	28	1.1 (0.6-2.0) 2-3 yr	17	0.7 (0.4-1.4) ≥4 yr
Inskip et al. 2001	139	0.8 (0.6-1.1)	51	1.0 (0.6-1.6) 0.5-3 yr	54	1.0 (0.6-1.6) ≥ 3 yr
					22	0.7 (0.4-1.4) ≥5 yr
Johansen et al. 2001	154	1.0 (0.8-1.1)	87	1.1 (0.9-1.3) 1-4 yr	24	1.0 (0.7-1.6) >5 yr
Auvinen et al. 2002	40 analogue, 16 digital	1.3 (0.9-1.8)	15 analogue, 11 digital	1.2 (0.7-2.0) 1-2 yr	17 analogue, 1 digital	1.5 (0.9-2.5) >2 yr
Hardell et al. 2002	188* analogue	1.3 (1.0-1.6)	188* analogue	1.3 (1.0-1.6) >1 yr	46* analogue	1.3 (0.8-2.3) >10 yr
	224* digital	1.0 (0.8-1.2)	224* digital	1.0 (0.8-1.2) >1 yr	33* digital	0.9 (0.6-1.5) >5 yr
Christensen et al. 2005	47 low-grade glioma	1.1 (0.6-2.0)	19	0.9 (0.4-1.8) 1-4 yr	6	1.6 (0.4-6.1) ≥10 yr
	59 high-grade glioma	0.6 (0.4-0.9)	24	0.6 (0.3-1.0) 1-4 yr	8	0.5 (0.2-1.3) ≥10 yr
	67 meningioma	0.8 (0.5-1.3)	35	0.8 (0.5-1.3) 1-4 yr	6	1.0 (0.3-3.2) ≥10 yr
Lönn et al. 2005	214 glioma	0.8 (0.6-1.0)	112	0.8 (0.6-1.1) 1-4 yr	25	0.9 (0.5-1.5) ≥10 yr
	118 meningioma	0.7 (0.5-0.9)	64	0.6 (0.4-0.9) 1-4 yr	12	0.9 (0.4-1.9) ≥10 yr
Hardell et al. 2005a, Hardell et al. 2005b	68 malignant, analogue	2.6 (1.5-4.3)	20 analogue	1.8 (0.9-3.5) 6-10 yr†	48 analogue	3.5 (2.0-6.4) >10 yr
	198 malignant, digital	1.9 (1.3-2.7)	100 digital	1.6 (1.1-2.4) 1-5 yr	19 digital	3.6 (1.7-7.5) >10 yr
	35 meningioma, analogue	1.7 (1.0-3.0)	1 analogue	1.2 (0.1-12) 1-5 yr	20 analogue	2.1 (1.1-4.3) >10 yr
	151 meningioma, digital	1.3 (0.9-1.9)	96 digital	1.2 (0.8-1.8) 1-5 yr	8 digital	1.5 (0.6-3.9) >10 yr
Hepworth et al. 2006	508 glioma	0.9 (0.8-1.1)	271 glioma	0.9 (0.7-1.1)	170 glioma	1.0 (0.8-1.3) 5-9 yr
				1.5-4yr	66 glioma	0.9 (0.6-1.3) ≥10yr

¹⁷ The table is modified from the report to the Swedish Radiation Protection board (SSI's Independent Expert Group on Electromagnetic Fields 2005).

¹⁸ RR – Relative Risk

¹⁹ CI – Confidence Interval

Possible effects of Electromagnetic Fields (EMF) on Human Health

Study	Brain tumours		Brain tumours short latency		Brain tumours longer latency	
	Number exposed cases	RR ¹⁸ estimate (95% CI ¹⁹)	Number exposed cases	RR estimate (95% CI)	Number exposed cases	RR estimate (95% CI)
Schüz et al. 2006a	138 glioma	1.0 (0.7-1.3)	82 glioma	0.9 (0.6-1.2) 1-4 yr	51 glioma	1.1 (0.8-1.7) ≥5yr
	104 meningioma	0.8 (0.6-1.1)	73 meningioma	0.9 (0.6-1.2) 1-4 yr	12 glioma 23 meningioma 5 meningioma	2.2 (0.9-5.1) ≥10yr 0.9 (0.5-1.5) ≥5yr 1.1 (0.4-3.4) ≥10yr
Schüz et al. 2006b	580	1.0 (0.9-1.1)	266	1.0 (0.9-1.2) 1-4 yr	28	0.7 (0.4-1.0) ≥10yr
			235	1.0 (0.8-1.1) 5-9 yr		

* Discordant pairs

† No cases had shorter than 6 years latency

Possible effects of Electromagnetic Fields (EMF) on Human Health

Table 3. Results of epidemiological studies on mobile phone use and acoustic neuroma.²⁰

Study	Acoustic neuroma		Acoustic neuroma short latency		Acoustic neuroma longer latency	
	Number exposed cases	RR ²¹ (95% CI ²²)	Number exposed cases	RR (95% CI)	Number exposed cases	RR (95% CI)
Hardell et al. 1999	5	0.8 (0.1-4.2)				
Inskip et al. 2001	22	1.0 (0.5-1.9)	8	1.8 (0.7-4.5) 0.5-2 yr	5	1.9 (0.6-5.9)
Johansen et al. 2001	7	0.6 (0.3-1.3)				
Muscat et al. 2002			7	0.5 (0.2-1.3) 1-2 yr	11	1.7 (0.5-5.1) 3-6 yr
Hardell et al. 2002	38* analogue	3.5 (1.8-6.8)	12* analogue	3.0 (1.0-9.3) 1-5 yr	7* analogue	3.5 (0.7-16.8) >10 yr
	23* digital	1.2 (0.7-2.2)	21* digital	1.2 (0.6-2.2) 1-5 yr	2* digital	2.0 (0.2-22.1) >5 yr
Lönn et al. 2004	89	1.0 (0.6-1.5)	44	0.8 (0.5-1.3) 1-4 yr	14	1.9 (0.9-4.1) ≥10 yr
Christensen et al. 2004	45	0.9 (0.5-1.6)	23	0.9 (0.5-1.6) 1-4 yr	2	0.2 (0.0-1.1) ≥10 yr
Hardell et al. 2005a	20 analogue	4.2 (1.8-10)	2 analogue	9.9 (1.4-69) 1-5 yr	11 analogue	5.1 (1.9-14) 5-10 yr
	53 digital	2.0 (1.0-3.8)	29 digital	1.7 (0.9-3.5) 1-5 yr	7 analogue 23 digital	2.6 (0.9-8.0) >10 yr 2.7 (1.3-5.7) 5-10 yr
Schoemaker et al. 2005†	360	0.9 (0.7-1.1)	174	0.8 (0.7-1.0) 1.5-4 yr	139	0.9 (0.7-1.2) 5-9 yr
					47	1.0 (0.7-1.5) ≥10 yr
Schüz et al. 2006b	32	0.7 (0.5-1.0)				
Takebayashi et al. 2006	97	0.7 (0.4-1.2)	21	0.8 (0.4-1.5) 4-7 yr	4	0.8 (0.2-2.7) ≥8yr

* Discordant pairs

† Partly overlapping with Lönn et al. (2004) and Christensen et al. (2004)

3.3.2.2. In vivo

What was already known on this subject?

The possible carcinogenicity of RF field exposure had been investigated in a number of experimental systems. Results had been essentially negative. An interesting exception is that of Repacholi et al. (1997), who had induced a two-fold increase in lymphoma incidence in a strain of lymphoma-prone transgenic mice (Eμ-Pim1) following exposure (2x30 min daily for up to 18 months) to 900 MHz RF fields with a signal similar to the GSM modulation (pulse repetition frequency of 217 Hz and a pulse width of 0.6 ms). No

²⁰ The table is modified from the report to the Swedish Radiation Protection board (SSI's Independent Expert Group on Electromagnetic Fields 2005).

²¹ RR – Relative Risk

²² CI – Confidence Interval

Possible effects of Electromagnetic Fields (EMF) on Human Health

attempt to replicate this finding had been published at the time of publication of the previous opinion.

What has been achieved since then?

Utteridge et al. (2002) failed to confirm the results of the Repacholi et al. (1997) study. Utteridge and co-workers found that exposure to RF fields (898 MHz; GSM modulation; 0.25/1.0/2.0/4.0 W/kg; 1 hour/day, 5 days/week for 104 weeks) had no statistically significant effect (95%-CI) on the incidence of lymphoma. Utteridge et al. (2002) used the same strain of mouse as the earlier study and they were obtained from same supplier; the investigators also fed the same food to the mice. The later study had some refinements in experimental design: four SAR levels were used instead of one in the original study, animals were restrained during the exposure for better control of variations in exposure level, and full necropsy was performed on all mice at the end of the study. Other differences from the Repacholi et al study were that animals were exposed once per day instead of during two episodes of 30 minutes 5 days per week.

Several other recent studies have evaluated carcinogenicity of RF fields in a variety of experimental models. Several studies have tested whether RF fields alone induce any type of cancer in normal or genetically predisposed animals (Zook and Simmens 2001, La Regina et al. 2003, Anderson et al. 2004, Sommer et al. 2004b), and several other studies investigated whether exposure to RF fields could enhance the development of tumours induced by chemical carcinogens, X-rays or UV radiation (Zook and Simmens 2001, Anane et al. 2003a, Bartsch et al. 2002, Imaida et al. 2001, Huang et al. 2005, Shirai et al. 2005, Heikkinen et al. 2001, Heikkinen et al. 2003, Heikkinen et al. 2006). No statistically significant increase of tumour incidence has been reported in any of these studies.

Most of the recent and earlier co-carcinogenicity studies on RF fields have used initiation-promotion protocols, which, however, may not be sufficient to test all aspects of co-carcinogenicity (Juutilainen et al. 2000). In addition, most of the carcinogenicity studies have used only one, relatively low, RF field exposure level.

3.3.2.3. In vitro

What was already known on this subject?

Various biological endpoints have been investigated after RF field exposure in vitro. Much of the work had focused on genotoxic effects, although there was no prior indication that non-thermal RF fields induce DNA damage. However, since some reports indicated genotoxic effects from RF fields, the earlier CSTE opinion recommended the confirmation of these findings.

What has been achieved since then?

Genotoxic effects

The photon energy of radiation from mobile phones is much lower than the energy necessary to break chemical bonds. It is therefore generally accepted that RF fields do not directly damage DNA. However, it is possible that certain cellular constituents altered by exposure to EMF, such as free radicals, indirectly affect DNA. In most studies, the genotoxic effects have been investigated after short-term exposure (for review see Moulder et al. 1999, Vijayalaxmi and Obe 2004).

The REFLEX study performed by twelve research groups in seven European countries, investigated basic mechanisms induced by EMF using toxicological and molecular biological technologies at cellular and sub-cellular levels in vitro. One of the REFLEX investigators (Diem et al. 2005) reported DNA strand breaks (measured by both the neutral and alkaline versions of the "comet" assay) in human diploid fibroblasts and cultured rat granulosa cells after RF field exposure (1800 MHz; SAR 1.2 or 2 W/kg;

Possible effects of Electromagnetic Fields (EMF) on Human Health

different modulations; during 4, 16 and 24h; intermittent 5 min on/10 min off or continuous wave), whereas it is not clear if continuous exposure of non-modulated or modulated 1800 MHz was used. Statistically significant increases in micronucleus formation and in chromosomal aberrations were observed in fibroblasts as well. In a recent replication study, (Speit et al. 2007) continuous wave with intermittent exposure (1800 MHz; SAR 2 W/kg) was applied using the same cell system and clearly negative results were obtained. Nikolova et al. (2005) reported after a 6-h but not after a 48-h RF field exposure a low and transient increase of DNA strand breaks in embryonic stem cell-derived neural progenitor cells.

Non-genotoxic effects

Several studies investigated the influence of RF fields on cell cycle kinetics, but in the majority of the investigations no effects were detected (Vijayalaxmi et al. 2001, Higashikubo et al. 2001, Zeni et al. 2003, Miyakoshi et al. 2005, Lantow et al. 2006c). Alteration in cell proliferation was described only in a few reports (Pacini et al. 2002, Capri et al. 2004b).

Programmed cell death which is also called apoptosis is a physiological mode of cell death occurring in development and cell differentiation and in response to mild damaging stimuli. It is an important protection mechanism against cancer, as it removes potential tumour cells. Several reports have investigated whether RF fields can induce apoptosis in human peripheral blood mononuclear cells (Capri et al. 2004a), lymphoblastoid cells (Marinelli et al. 2004), epidermis cancer cells (Caraglia et al. 2005), human Mono Mac 6 cells (Lantow et al. 2006c) and in Molt4 cells (Hook et al. 2004). No difference in apoptosis induction was detected between sham-exposed and RF field exposed cells. On the other hand, Marinelli et al. reported better survival rate of T lymphoblastoid leukaemia cells exposed to 900 MHz non-modulated RF fields and Caraglia et al. (2005) found apoptosis induction in human epidermoid cancer cells after exposure to 1.95 GHz RF fields.

Participants of the REFLEX-study reported no effects of RF fields on cell cycle, cell proliferation, cell differentiation, apoptosis induction, DNA synthesis, and immune cell functionality. The authors described some findings after RF fields exposure on the transcript level of genes related to apoptosis and cell cycle control; however, these responses were not associated with detectable changes of cell physiology (Nikolova et al. 2005). Analysis on whole-genome cDNA arrays showed alterations in gene expression after various RF exposure conditions using different cell types, but no consistent RF-signature such as stress response could be identified (Remondini et al. 2006)

Heat-shock proteins (HSP) are an important group of cell response proteins. They act primarily as molecular chaperones to eliminate unfolded or miss-folded proteins, which can also appear from cellular stress. This stress response can be induced by many different external factors, including temperature, chemicals, oxidative stress, heavy metals, ionizing and non-ionizing radiation and ultrafine carbon black particles. Hsp70 has been shown to interfere with post-mitochondrial events to prevent free radical mediated apoptosis (Gotoh et al. 2001). An increased expression level of Hsp70 can thus confer protection against cellular stress. On the other hand, it is discussed that heat-shock proteins are also involved in oncogenic processes (Jolly et al. 2000, Inoue et al. 1999, French et al. 2001). Some investigators have described increased heat-shock protein level after RF field exposure (Leszczynski et al. 2002, Kwee et al. 2001, de Pomerai et al. 2000). However, these results are controversial, because there are other negative findings (for a review see Cotgreave (2005)). Interestingly, de Pomerai and his co-workers could not confirm their earlier findings, and the new data indicate that small temperature differences may have contributed to the earlier results (Dawe et al. 2006).

Nikolova et al. (2005), authors of the REFLEX-study, described modulation in gene regulation after RF fields exposure at a SAR of 1.5 W/kg in p53-deficient embryonic stem

Possible effects of Electromagnetic Fields (EMF) on Human Health

cells. Proteomic analyses of human endothelial cell lines showed RF fields induced changes in the expression and phosphorylation state of numerous proteins including the heat shock protein hsp27.

Free radicals are able to interact with DNA or other cellular components and are involved in many cell regulatory processes.

In leukocytes, physiological activation is associated with the onset of phagocytosis and leads to increased formation of reactive oxygen species (ROS). These cells exert a wide variety of functions including the regulation of the immune response (pro and anti inflammatory processes), scavenging of senescent cells, phagocytosis of infected or malignant cells, wound healing, repair, and detoxification, but also the generation of free radicals to kill invading micro-organisms. Each type and source of free radicals enhances important physiological processes, e.g., signal transduction of various membrane receptors and further immunological functions. An imbalance between excessive formation of reactive oxygen species and the limited antioxidant defense, known as oxidative burst (Sies and Cadenas 1985), can cause damage to nucleic acids, membranes, proteins, lipids and polysaccharides (Beckman and Ames 1998). During healthy conditions free radicals are neutralized by an elaborate defense system. Only a few publications are available describing the capacity of RF fields to affect free radical dependent processes in cells. In recent studies (Lantow et al. 2006a, Lantow et al. 2006b, Simkó et al. 2006) no increased free radical level was detected.

Influences on immune system cells were investigated in a few studies. No significant effects were observed on intracellular production of interleukin-2 (IL-2) and interferon (INF) gamma in lymphocytes, IL-1 and tumour necrosis factor (TNF)-alpha in monocytes, on immune-relevant genes (IL 1-alpha and beta, IL-2, IL-2-receptor, IL-4, macrophage colony stimulating factor (MCSF)-receptor, TNF-alpha, TNF-alpha-receptor) (Tuschl et al. 2005, Black and Heynick 2003).

Discussion

Effects of RF fields on different biological systems have been investigated. Although the majority of studies have found no evidence of genotoxic effects, there are a few positive findings that should be followed up. Some in vitro studies provide evidence that gene expression is affected at RF exposure close to the guidelines. If these studies are confirmed they will be important for a mechanistic understanding of the interaction of RF fields with cellular tissue. Overall, there is little evidence of any health-relevant in vitro effects of RF electromagnetic fields below guidelines. While it seems appropriate to perform experimental studies using pure experimental RF fields, it may be needed to emulate the complex modulation patterns and intensity variations typical to real mobile phone use in future studies. This way data can be obtained which are better suited for comparison to epidemiologic studies.

3.3.3. Symptoms

What was already known on this subject?

In the 2001 opinion it was concluded that the knowledge was insufficient for the implementation of measures aimed at the identification and protection of a highly sensitive sub-group of the population. With regard to reports of subjective symptoms from individuals (possibly "hypersensitive"), the limited number of studies on volunteers had found no connection between reported symptoms and exposure to electromagnetic fields. There was a lack of information on the role of conditions of exposure (frequency, concentration duration etc) and possible biological mechanism. While epidemiological studies had not shown any consistent evidence of effects on humans, it was pointed out

Possible effects of Electromagnetic Fields (EMF) on Human Health

that this could not be taken to mean that RF field exposure does not pose any hazard to human health.

What has been achieved since then?

A variety of non-specific self-reported symptoms (for example headache, fatigue, dizziness and concentration difficulties) have been suggested to be triggered by exposure to RF fields. These possible health effects have been discussed and studied mainly from two different aspects: 1) a possible increase in symptoms in populations living close to mobile communication base stations and 2) reports from individuals that exposure to RF from mobile phones (and sometimes also base stations) triggers symptoms. In the latter case, some individuals attribute their health problems to an increased sensitivity (hypersensitivity) to electromagnetic fields. The term "electromagnetic hypersensitivity" (EHS) has been used to describe such cases of non-specific medically unexplained health problems attributed by the afflicted individuals to electromagnetic fields (in many cases including ELF fields) or to being in the vicinity of electrical equipment (see also the section on ELF fields 3.5.3 Symptoms).

There have only been a few attempts to study symptom prevalence and symptom severity in relation to exposure to RF fields from base stations. The methodological limitations of these cross-sectional epidemiological studies preclude conclusions regarding a possible causal relationship between increase in symptoms and exposure to RF. One study by Hutter et al. (2006) performed measurements of RF fields in the bedrooms of the participants. In this Austrian study, inhabitants of homes in the vicinity (distance 20-600m) of ten selected base stations (five in Vienna and five in a rural area, Carinthia) were invited to participate in a study on environment and health. Mobile communication base stations were not mentioned in the invitation. Spot measurements were done after the questionnaires and computer based tests were completed. The participants were classified into three exposure groups based on calculations of the theoretical maximal power density from the selected base stations (i.e. when the base station is using 100% of its capacity). The mean power densities were 0.04, 0.23 and 1.3 mW/m² in the respective groups. Three out of 14 self-reported symptoms (headache, cold hands or feet and difficulties to concentrate) were significantly more commonly reported in the highest exposure group. The results were adjusted for age, sex, region, regular use of mobile phone and fear of adverse effects of the base station, but not for socio-demographic factors. Exposure from other sources (e.g. other base stations) was not reported. The results should be replicated before any conclusions on a causal relationship between RF exposure and the reported effect on some symptoms may be drawn. Provocation studies on symptoms should also be considered. Compared to epidemiological studies provocation studies may have the advantage of better controlled exposure conditions and randomization. The focus is however somewhat different in provocation studies, which will assess only acute effects while epidemiological studies may include long term effects on well-being.

A relationship between RF and symptoms in healthy volunteers was investigated in one provocation study (Koivisto et al. 2001). No increase in symptoms was observed during RF exposure as compared to sham exposure. The limited number of studies on detection of RF at exposure levels relevant to mobile communication systems under blind conditions has not provided any consistent proof of ability to detect the fields, neither in healthy individuals nor in subjects who report EHS.

Health complaints described as EHS and reported to be triggered by mobile phones have also been studied in a limited number of provocation studies. A WHO Workshop on Electrical Hypersensitivity (WHO 2005) and recent reviews of the literature on subjective health complaints associated with electromagnetic fields of mobile phone communication (Seitz et al. 2005) and provocation studies including subjects reporting EHS (Rubin et al. 2005) have presented similar conclusions. The main conclusion is that although symptoms described as EHS are real and may be severe and disabling, a relationship

Possible effects of Electromagnetic Fields (EMF) on Human Health

between symptoms and RF field exposure has not been proven. Most likely, the health problems described as EHS are not related to the physical presence of EMF and more research is needed to learn more about the conditions inducing EHS.

The studies published after the WHO workshop and the reviews mentioned above have not provided any information that would contradict these conclusions. The influence of a base station-like exposure of 2140 MHz RF fields on well-being that was reported already in 2003 (Zwamborn et al. 2003) was not confirmed in a follow-up study (Regel et al. 2006). In a study published by the TNO-Institute²³ (the so called TNO study), only available as grey cover report Zwamborn et al. (2003) found a decrease in well-being (based on an index of 23 items from subscales of anxiety, somatic symptoms, inadequacy, depression and hostility) during exposure to 2140 MHz RF fields (UMTS). The decrease in well-being was shown in the group of subjects who were recruited based on self-reported sensitivity to RF fields as well as in the non-symptomatic control group. No effect on symptoms was observed in any of the groups for 945 MHz or 1840 MHz RF fields (GSM). The study, while reporting an effect of UMTS exposure, thus failed to confirm the reported belief in the group with self-reported sensitivity that GSM exposure triggered symptoms. The TNO study was widely discussed since a possible health effect of UMTS might have large implications on the introduction and use of the third generation of mobile networks (UMTS). Follow-up studies were initiated in several countries and in 2006 the first results were published (Regel et al. 2006). The experimental protocol was improved as compared to the TNO-study, e.g. a larger study group, better dosimetry and longer intervals between the exposure sessions. Two exposure levels of RF fields (E-fields strength of 1 V/m –as in the TNO study- and 10 V/m) were applied, in order to assess any possible dose-response relationship. Peak spatial SARs in the brain (averaged over 10 g) were 45 and 4500 $\mu\text{W}/\text{kg}$. Well-being was assessed in two standard questionnaires (including the one used in the TNO study). No effect on symptoms was observed, neither in the group of subjects with self-reported RF-related symptoms nor in the non-symptomatic control group. There was no association between perceived field strengths and actual RF exposure.

Discussion

Scientific studies have failed to provide support for a relationship between RF exposure and self-reported symptoms sometimes referred to as EHS. Present knowledge suggests that symptoms are not correlated to RF field exposure, but few studies have addressed this issue directly. The exposure levels from base stations are very low compared to the exposure during the use of a mobile phone. Research regarding health effects from base stations where exposure is significantly lower than for mobile phone users is mainly driven by concern in the general population

The symptoms attributed to ELF and RF fields are similar and in many cases the afflicted subjects report both ELF and RF fields to trigger symptoms. There are more studies on self-reported symptoms and exposure to ELF fields, but also in this case the scientific studies have failed to confirm a causal relationship (see also the section on ELF fields 3.5.3 Symptoms).

3.3.4. Nervous system effects

What was already known on this subject?

Due to the proximity of mobile phones to the head, public concerns were raised regarding a potentially toxic effect of RF on the central nervous system. Five aspects are usually considered in toxicology regarding the nervous system: morphology, brain

²³ Located in The Hague, The Netherlands

Possible effects of Electromagnetic Fields (EMF) on Human Health

function, electrophysiology, behaviour and development (which is addressed in a later paragraph).

Several studies had been published concerning the potential neurotoxic effects of radiofrequencies emitted by the mobile phones. Transient minor effects were observed on the electroencephalogram (EEG), sleep structure, and on cognitive processes in human subjects (Mann and Röschke 1996, Preece et al. 1999, Huber et al. 2000, Koivisto et al. 2000a, Koivisto et al. 2000b, Krause et al. 2000). Some of the observations could not be replicated (Wagner et al. 1998, Wagner et al. 2000), and studies with negative outcomes were also published (Röschke and Mann 1997).

In animals, some previous studies did show disturbance of work memory in rats exposed to RF (Lai et al. 1994, Wang and Lai 2000). However, the most surprising effect was that very low SAR values (mW/kg) caused increased permeability of the blood-brain-barrier (BBB) in rats (Salford et al. 1994, Persson et al. 1997). Alterations of the BBB had also been found in another study (Neubauer et al. 1990), but not by Tsurita et al. (2000). In rats exposed to 2 W/kg Fritze and co-workers demonstrated effects on the BBB only at SAR levels above 7.5 W/kg (Fritze et al. 1997). The BBB isolates the CNS from the rest of the organism, controls molecule fluxes, and protects the brain (Purves et al. 2001). Increased permeability of the BBB can allow unwanted substances to reach the CNS, with possible pathological consequences (inflammation, neurone death).

What has been achieved since then?

Human studies

In humans, transitory minor effects (both positive and negative) have been observed on EEG patterns, sleep structure, and cognitive processes (D'Costa et al. 2003, Cook et al. 2002, Hossmann and Hermann 2003, Sienkiewicz et al. 2005). Also studies where no effects were documented have been published, even after a repeated exposure (Besset et al. 2005).

Since the ear is very close to the exposure source, some studies have checked the auditory system under or after exposure, and even after repeated cumulative exposure. No effect has been observed (Ozturan et al. 2002, Arai et al. 2003, Bak et al. 2003, Parazzini et al. 2005, Uloziene et al. 2005).

Animal studies

Slight changes in EEG activity and neurotransmitters have been observed in animals at low SARs (reviewed by Sienkiewicz et al. 2005). Regarding cognitive functions, a recent report showed that a disturbance of learning and memory in rats exposed at 2.45 GHz CW could be inhibited by a magnetic field (incoherent noise) (Lai 2004a). Results from earlier studies on learning and memory at non-thermal RF levels have not been corroborated (Dubreuil et al. 2003, Yamaguchi et al. 2003, Cobb et al. 2004, Cassel et al. 2004). No morphological effects have been observed below thermal thresholds (D'Andrea et al. 2003).

Salford and co-workers published another work showing changes in BBB permeability at low SAR (Salford et al. 2003), whereas others did not find any such alteration (Finnie et al. 2001), even with repeated exposures up to 2 years (Finnie et al. 2002).

No effects have been seen on auditory system function (Aran et al. 2004) or on development of multiple sclerosis in rats (Anane et al. 2003b).

What are the overall conclusions?

Overall analyses do not show any clear neurotoxic effect, at any level studied. Slight changes in electrical activity or neurotransmitter biochemistry have been observed. Those changes do not act on cognitive processes, behaviour or memory and do not suggest pathological hazards. Furthermore, no clear role of modulation has appeared.

Possible effects of Electromagnetic Fields (EMF) on Human Health

Although extrapolation from animals to humans raises some difficulty, the rat or the mouse are common models to look for toxicity and the few studies showing significant alterations are to be considered carefully. For all cognitive experiments in animals, stress effects due to restraint must be clearly identified and prevented when looking at effects of RF fields.

3.3.5. Miscellaneous human

Initial observations of a blood pressure decrease after mobile phone exposure have not been replicated (Braune et al. 1998, Braune et al. 2002). The only effects on cardiovascular functions that have been replicated are increased blood-flow in the external ear (Monfrecola et al. 2003, Roelandts 2003). Local temperature increases during exposure have been reported (Paredi et al. 2001, Curcio et al. 2004), possibly related to vasodilation caused by heating of mobile phone electronics and battery.

3.3.6. Reproduction and development

Epidemiological studies of adverse pregnancy outcomes following exposure to RF fields have been reviewed by Verschaeve and Maes (1998), Heynick and Merrit (2003) and Feychting (2005a). The evidence on possible effects of RF fields on pregnancy outcomes is virtually limited to occupational exposures among physiotherapists. The endpoints studied include spontaneous abortions, birth weight, gender ratio, and congenital malformations. Although some positive findings have been reported, no specific type of malformation or other adverse outcome has been consistently reported. Several of the studies have limited statistical power, especially for rare outcomes such as malformation, and there is a potential for recall bias. The available results do not allow any definite conclusions.

Numerous studies have evaluated developmental effects of RF fields on mammals, birds, and other non-mammalian species. These studies, reviewed recently by Heynick and Merritt (2003) and Juutilainen (2005), have clearly shown that RF fields are teratogenic at exposure levels that are sufficiently high to cause significant increase of temperature and exceed reference levels from exposure guidelines. There is no consistent evidence of effects at nonthermal exposure levels. However, only a few studies have evaluated possible effects on postnatal development using sensitive endpoints, such as behavioural effects.

3.3.7. Sensitivity of children

Concerns about the potential vulnerability of children to RF fields have been raised because of the potentially greater susceptibility of their developing nervous system; in addition, their brain tissue is more conductive than that of adults since it has a higher water content and ion concentration, RF penetration is greater relative to head size, and they have a greater absorption of RF energy in the tissues of the head at mobile telephone frequencies. Finally, they will have a longer lifetime exposure.

Few relevant epidemiological or laboratory studies have addressed the possible effects of RF field exposure on children. Owing to widespread use of mobile phones among children and adolescents and relatively high exposures to the brain, investigation of the potential effect of RF fields in the development of childhood brain tumour is warranted. The characteristics of mobile phone use among children, their potential biological vulnerability and longer lifetime exposure make extrapolation from adult studies problematic.

There is an ongoing debate on possible differences in RF absorption between children and adults during mobile phone usage, e.g. due to differences in anatomy (Wiert et al. 2005,

Possible effects of Electromagnetic Fields (EMF) on Human Health

Christ and Kuster 2005). Several scientific questions like possible differences of the dielectric tissue parameters remain open.

The anatomical development of the nervous system is finished around 2 years of age, when children do not yet use mobile phones although baby phones have recently been introduced.

Functional development, however, continues up to adult age and could be disturbed by RF fields.

3.3.8. Conclusions about RF fields

Since the adoption of the 2001 opinion, extensive research has been conducted regarding possible health effects of exposure to low intensity RF fields. This research has investigated a variety of possible effects and has included epidemiologic, in vivo, and in vitro research. The overall epidemiologic evidence suggests that mobile phone use of less than 10 years does not pose any increased risk of brain tumour or acoustic neuroma. For longer use, data are sparse, since only some recent studies have reasonably large numbers of long-term users. Any conclusion therefore is uncertain and tentative. From the available data, however, it does appear that there is no increased risk for brain tumours in long-term users, with the exception of acoustic neuroma for which there is limited evidence of a weak association. Results of the so-called Interphone study will provide more insight, but it cannot be ruled out that some questions will remain open.

Scientific studies have failed to provide support for a relation between RF exposure, lower than the reference values in the present ICNIRP guidelines and self-reported symptoms (sometimes referred to as electromagnetic hypersensitivity). Available studies suggest that self-reported symptoms are not correlated to an acute exposure to RF fields, but the limited number of studies does not allow any firm conclusion.

Currently available studies on neurological effects and reproductive effects have not indicated any health risks at exposure levels below guidelines.

Animal cancer studies have not provided evidence that RF radiation could induce cancer, enhance the effects of known carcinogens, or accelerate the development of transplanted tumours. The open questions include adequacy of the experimental models used and scarcity of data at high exposure levels. These questions are addressed by the still ongoing and planned carcinogenicity studies.

There is no reliable indication from in vitro research that RF fields affect cells at nonthermal exposure. However, recent results suggesting genotoxic effects need to be better understood.

Thus, no health effect has been consistently demonstrated at exposure levels below existing exposure guidelines for the general public. However, data on long term exposure and intracranial tumours are still sparse and in particular for acoustic neuroma some data indicate that an association with RF fields from mobile telephony is possible. For diseases other than cancer, very little epidemiologic data are available. A particular consideration is mobile phone use by children. While no specific evidence exists, there is a general concern that children or adolescents may be more sensitive to RF field exposure than adults. Children, as adults, will probably have a higher cumulative exposure compared to today's adults. To date no epidemiologic studies on children are available.

The technical development is very fast and sources of RF exposure become increasingly common. Yet, there is a profound lack of mechanistic understanding of effects below the guidelines and of information on individual RF exposure and the relative contribution of different sources to the overall exposure.

Possible effects of Electromagnetic Fields (EMF) on Human Health

3.4. Intermediate Frequency Fields (IF fields)

Intermediate frequencies are, in the frame of this report, defined as frequencies between 300 Hz and 100 kHz. They involve two different mechanisms, namely induced currents and dielectric absorption. The frequency limit when one predominates over the other is not precisely defined. Existence of effects depends upon two superimposed phenomena: absorption of the external field in the organism at the macroscopic level and the stimulation of biological effects by the penetrating fields. Those two phenomena depend on the kind of field, electric or magnetic, and on the frequency. Well-known biological effects are nerve stimulation at low frequencies and heating at high frequencies.

3.4.1. Sources and distribution of exposure in the population

The number of applications in this frequency range has been increasing in recent years. Examples are anti theft devices operated, e.g., at the exits of shops. Depending on the type of system, they are operated at very different frequencies ranging from some tens of Hz to a few GHz. The majority of these applications are operated in the intermediate frequency range. Close to some systems the so called reference levels can be exceeded under worst case conditions, but for most of the systems the exposure is well below the recommended limits. Other applications are induction hobs and hotplates typically operated at frequencies between 20 to 50 kHz, electric engines, and badge readers (typical frequency about 100 kHz). Information on the exposure due to such applications is scarce. Still common sources are visual display units containing cathode ray tubes which are causing emissions in the ELF range and the IF range, in the order of 1 nT up to 50 nT. Radio transmitters operated in the long wave range (30 kHz to 300 kHz) can cause exposure in the intermediate frequencies with levels above the recommended limits. Therefore, safety precautions need to be implemented both for the general public and workers. Some industrial applications like induction heating and welding need to be mentioned. Welding devices can cause considerable exposure up to a few hundred kHz. Induction heaters are operated in a frequency band from typically some tens of Hz to some tens of kHz, the exposure levels can reach values of about 100 μ T or more. Welding is a complex process that can cause emissions up to a few 100 kHz. The sparse information on IF field exposure due to welding devices available so far indicates that safety measures need to be implemented in some cases.

Some medical applications exist in the IF range. One example is electrosurgery used very commonly in hospitals. These systems are operated at some hundred kHz. In addition, the IF fields of typically up to 10 kHz arising from MRI applications need to be mentioned.

3.4.2. Health Effects

Epidemiological studies on IF fields were reviewed at a workshop organized by the WHO and by ICNIRP in 1999 (Hietanen 1999). It seems that very few useful data are available for health risk assessment. The few studies that do exist are relatively old occupational studies using occupations or job titles but no actual exposure estimation. Groups that have been studied include VDU (video display units) users and radio and telegraph operators. One study looked at radio amateurs. The studied outcomes include ocular effects, cardiovascular effects, cancer, and reproductive effects.

The available in vivo and in vitro evidence was reviewed in articles published in the proceedings of the WHO/ICNIRP seminar on IF fields (Juutilainen and Eskelinen 1999, Glaser 1999, Litvak and Repacholi 1999). In contrast to the active experimental and epidemiological research on ELF and RF fields, only a very limited number of studies have addressed the biological effects of IF fields. While there is limited evidence for effects on reproduction and development (Juutilainen 2005, Huuskonen et al. 1998),

Possible effects of Electromagnetic Fields (EMF) on Human Health

studies on other effects (such as carcinogenicity, genotoxicity, nervous system effects and general toxicity) are almost totally lacking.

3.4.3. Conclusions about IF fields

It is considered that the well established hazard mechanisms in the IF range are associated with a limited number of phenomena and apply to acute exposures. However, the extension to long term effects is based on weak grounds and on possibly unjustified assumptions about frequency dependence of effects (Litvak et al. 2002). In addition to established mechanisms, comprehensive risk assessment should consider also other information, such as well-conducted epidemiological and laboratory studies. Studies on possible effects associated with chronic exposure at low exposure levels (below exposure limits) are particularly relevant for assessing risks to human health and for confirming adequacy of current exposure limits.

Proper evaluation and assessment of possible health effects from exposure to IF fields is essential because human exposure to such fields increases due to new and emerging technologies.

3.5. Extremely low frequency fields (ELF fields)

3.5.1. Sources and distribution of exposure in the population

The exposure due to electric fields and magnetic flux densities in the ELF range arises from a wide variety of sources (IARC 2002). The most prominent frequencies are 50 and 60 Hz and their harmonics, often called power frequencies. For residential exposure, the major sources are household appliances, nearby power and high voltages transmission lines, and domestic installations. In some cases trains have to be considered, too. Looking at occupational exposure, installations of the electric power industry, welding, induction heaters and electrified transporting systems are important examples of ELF exposure sources. The highest electric field strengths typically occur close to high voltage transmission lines and can reach 5 kV/m and in a few cases more. The highest magnetic flux densities can be found close to induction furnaces and welding machines. Levels of a few mT are possible.

It needs to be mentioned that the maximum possible exposure next to a specific source often differs by some orders of magnitude from the average individual exposure of a person (to specify time weighted average exposure in many cases the arithmetic mean or the geometric mean or the median value are applied). To evaluate the distribution of the exposure in the population, meters are used. For assessment of compliance with exposure limits, the maximum possible exposure next to devices must be measured. An example might be a lineman: the average exposure due to magnetic flux density could be about 4 μ T (IARC 2002), but the maximum exposure close to a transmission line can reach 40 μ T or more. For the general population even larger variations between maximum and average exposure can be expected. Information on ELF exposure is mainly based on data from the United States and Western Europe.

Exposure of the general population

Several fixed installed sources are operated in our environment. Prominent examples are high voltage transmission lines operated between 110 and 400 kV at 50 or 60 Hz. The exposure of bypassing people can typically reach values of 2 to 5 kV/m for the electric field strength. The exposure due to magnetic flux density depends on the actual current on the line; fields up to 40 μ T are possible but are usually lower. It is important to notice that such exposure levels occur only directly below the lines; exposure decreases with the square of distance to the lines. In addition, intermediate voltage transmission lines

Possible effects of Electromagnetic Fields (EMF) on Human Health

(10 kV to 30 kV) and distribution lines (400 V) have to be considered; exposure levels are in such cases much lower. Typically values of 100 to 400 V/m and 0.5 to 3 μT can be reached, the exposure is usually instantaneous. Another approach to establish power supply is the use of underground buried cables. Electric field strength exposure can be neglected in this case; the distribution of the magnetic flux density differs compared to power lines. Substations and power plants are usually not accessible to the general public. Railway power supply installations are often operated at 16 2/3 Hz. The exposure decreases linearly with the distance. The exposure levels in areas accessible for the general public are below the limits. The highest magnetic flux densities can be found close to several domestic appliances that incorporate motors, transformers, and heaters. Such exposure levels are very local and decrease rapidly with the distance, exposure is instantaneous. An example is a vacuum cleaner: at a distance of 5 cm magnetic flux densities of about 40 μT can occur, but at 1 m the exposure will be around 0.2 μT . Looking at the individual exposure of persons, a few percent of the European population are in their homes exposed above a median magnetic flux density above 0.2 μT .

Exposure of workers

In a few locations in installations of the electric power industry the exposure limits given in the directive 2004/40/EC for occupational exposure can be reached or exceeded. Safety measures for such areas have to be implemented. An example is a peak electric field strength of more than 20 kV/m that was measured in a power station. Other examples of industrial applications in the ELF range are induction and light arc ovens or welding devices. The frequencies of such applications fall both in the ELF and in the intermediate frequency range. Exposure of workers has to be controlled for such devices. Next to welding devices maximum flux densities of several hundred μT are possible, depending on the welding current and the type of application.

Medical applications

Bone growth stimulation is used as a therapeutic application in the ELF range. In this case coils are applied where the fracture is located to stimulate the healing process. Other applications include Transcranial Magnetic Stimulation, wound healing, or pain treatment. A diagnostic application is the bioimpedance measurement for cancer detection.

3.5.2. Cancer

3.5.2.1. Epidemiology

What was already known on this subject?

In 2002, the International Agency on Research on Cancer (IARC) published a monograph on the evaluation of carcinogenic risks of static and extremely low-frequency (ELF) electric and magnetic fields to humans (IARC, 2002). ELF magnetic fields were classified into group "2B" ("possibly carcinogenic to humans"). While the outcome of this evaluation was already known at the time of the last opinion, the IARC reasons for this decision were not yet published. The justification states limited evidence in humans based on consistent results from sound epidemiological studies showing an association with an increased leukaemia risk in children at average field strengths above 0.3/0.4 μT (Ahlbom et al. 2000, Greenland et al. 2000), but bias could explain some of the raised risk. The findings from observational studies are not supported by studies in experimental animals, which provide inadequate evidence of carcinogenicity.

Furthermore, the IARC monograph concluded, there was no evidence for an association of ELF magnetic fields with any other type of cancer. ELF electric fields were grouped into "3" ("is not classifiable as to its carcinogenicity to humans").

Possible effects of Electromagnetic Fields (EMF) on Human Health

What has been achieved since then?

Only a few studies on childhood leukaemia have been conducted since the adoption of the previous opinion, and they did not add anything substantial to the previous studies. At a workshop of WHO in 2004, possible explanations for the childhood leukaemia finding have been put forward (summarized in Kheifets et al. (2005)). None of them reaches a level beyond hypothesis. One recent study has observed a decreased survival in children with leukaemia being exposed to average ELF magnetic fields above 0.3 μ T (Foliart et al. 2006). This finding, however, is based on small numbers and no mechanism has been proposed, so confirmation studies have to be awaited before conclusions should be drawn. Most new ELF studies have been looking into breast cancer or brain tumour risk. Breast cancer caught particular interest because of experimental results suggesting that melatonin synthesis was related to ELF field exposure and because melatonin might play a role in the development of breast cancer. Several studies also reported an increased breast cancer risk among subjects with elevated ELF exposure. However, later big and well controlled studies have been entirely negative and the hypothesis of a link between ELF field exposure and breast cancer risk is essentially written off (Forsen et al. 2005). While some new data on brain tumours have appeared since the previous opinion, firm conclusions can still not be drawn.

Discussion

Little data that have an impact on the evaluation have appeared since the previous opinion. Therefore, the previous assessments stay the same. The fact that the epidemiologic results for childhood leukaemia have little support from known mechanisms or experimental studies is intriguing and it is of high priority to reconcile these data.

3.5.2.2. In vivo

What was already known on this subject?

The previous opinion did not evaluate evidence of carcinogenicity from animal studies. However, such data were included in the monograph by IARC that classified ELF magnetic fields into group 2B, "possibly carcinogenic to humans", based on epidemiological studies showing an association between residential ELF magnetic fields and childhood leukaemia (IARC 2002). The long-term animal carcinogenicity studies reviewed by IARC provided very little evidence that exposure to ELF magnetic fields alone could induce any type of cancer, including hemopoietic, mammary, brain and skin tumours. Negative results were also obtained from studies that evaluated the effects of ELF magnetic fields on growth of transplanted tumour cells. Animal studies that combined magnetic fields with known carcinogenic agents produced more equivocal results, although also these co-carcinogenicity studies were mostly negative. Among the few positive findings are enhanced development of UV-induced mouse skin tumours in one study (Kumlin et al. 1998) and accelerated development of rat mammary tumours induced by 7,12-dimethylbenz(a)anthracene (DMBA) in several experiments by a German research group (Löscher et al. 1993, Baum et al. 1995, Mevissen et al. 1996, Mevissen et al. 1998, Thun-Battersby et al. 1999). The latter findings were not substantiated in independent replication studies (Anderson et al. 1999, Boorman et al. 1999), but there are differences in experimental details that could potentially explain the differences in results (Anderson et al. 2000, Löscher 2001). Based on the available experimental studies, IARC concluded that there is *inadequate evidence* for carcinogenicity of ELF magnetic fields in experimental animals.

What has been achieved since then?

Motivated by the epidemiological findings of increased leukaemia risk in children, Sommer and Lerchl (2004a) investigated the influence of 50 Hz (1 or 100 μ T) magnetic fields in the AKR/J mouse strain genetically predisposed to thymic lymphoblastic lymphoma. There was no effect of magnetic field exposure on survival, and the time to

Possible effects of Electromagnetic Fields (EMF) on Human Health

lymphoma development did not differ between exposed and sham-exposed animals. The results do not support the hypothesis that chronic exposure to 50 Hz magnetic fields increases the risk of hemopoietic malignancy in this experimental model. However, the relevance of the model to human childhood leukaemia is limited.

New results have been published by German researchers who have reported accelerated development of DMBA-induced rat mammary tumours. In their most recent study (Fedrowitz et al. 2004) they tested the hypothesis that use of different sub strains of SD rats explains the difference between their previous results and those of the replication studies. The results were consistent with the hypothesis: exposure to a 100 μ T, 50 Hz magnetic field enhanced mammary tumour development in one sub strain of SD rats, but not in another sub strain obtained from the same breeder. The tumour data were supported by the finding that exposure to MF increased cell proliferation in the mammary gland of the MF-sensitive strain, but no such effect was seen in the insensitive sub strain. The finding is potentially important for explaining the inconsistent results, if the sub strain-specific effect of MF exposure is confirmed in further independent experiments.

Although short-term animal studies are considered less relevant for cancer risk assessment than long-term carcinogenicity and co-carcinogenicity studies, they can provide important contributions to understanding the mechanisms of carcinogenic effects. *Genotoxicity* of ELF magnetic fields was studied by Lai and Singh (2004b), who reported significantly increased DNA damage after exposure to a 60 Hz, 10 μ T magnetic field for 24 or 48 hours. Although the effect was relatively small, it was seen in several independent experiments. The effects were blocked by treatment with a radical scavenger, a nitric oxide synthase inhibitor and an iron chelator, suggesting involvement of free radicals and iron in the effects of magnetic fields. The same authors have previously reported similar effects after short (2 hour) exposure to higher magnetic flux densities of 0.1-0.25 mT. Environmental agents can promote the development of cancer also through *non-genotoxic* mechanisms such as stimulation of cell proliferation and inhibition of apoptosis. In support of their previous results suggesting co-carcinogenic effects of ELF magnetic fields (described above), two research groups have reported increase in cell proliferation markers in rat mammary gland (Fedrowitz et al. 2002) and inhibition of UV radiation-induced apoptosis in mouse skin (Kumlin et al. 2002) after short-term exposure to magnetic fields at 100 μ T. The results of the short-term animal studies are interesting and, if confirmed in further independent experiments, potentially important for understanding possible cancer-related effects of magnetic fields.

Discussion

Overall there is no evidence from animal studies that ELF magnetic field exposure alone causes tumours or that it enhances the growth of implanted tumours. There is some inconsistent evidence that ELF magnetic fields of about 100 μ T may enhance the development of tumours induced by known carcinogens, but the majority of studies evaluating such co-carcinogenic effects have been negative. Results from recent studies are potentially helpful for explaining mechanisms and inconsistencies of previous findings, but they lack confirmation in independent experiments, and are not sufficient to challenge IARC's evaluation that the experimental evidence for carcinogenicity of ELF magnetic fields is *inadequate*.

3.5.2.3. In vitro

What was already known on this subject?

There are many observations of cellular responses induced by ELF magnetic fields in vitro. A large number of cellular components, cellular processes, and cellular systems can conceivably be affected by EMF exposure. However, because evidence from theoretical and experimental studies suggest that ELF fields are unlikely to induce DNA damage directly, most studies have been conducted to examine effects on the cell membrane, general and specific gene expression, and signal transduction pathways. In addition, a

Possible effects of Electromagnetic Fields (EMF) on Human Health

large number of studies have been performed to investigate effects on processes such as cell proliferation, cell cycle regulation, cell differentiation, metabolism, and various physiological characteristics of cells.

Summaries of in vitro studies are found in Portier and Wolfe (1998) and IARC (2002). In particular, studies focusing on cell cycle kinetics, proliferation, differentiation, gene expression, DNA damage, signal transduction pathways, apoptosis and membrane characteristics have received attention and are useful in carcinogen evaluation.

What has been achieved since then?

It is generally accepted that ELF fields do not transfer energy to cells in sufficient amounts to cause direct DNA damage and subsequent genotoxic effects. However, it is possible that certain cellular processes, such as DNA repair, are altered by exposure to EMF, which could indirectly affect the structure of DNA causing strand breaks and other chromosomal aberrations, including sister chromatid exchange, or micronucleus formation.

A recent review of genotoxic effects after ELF field exposure (Vijayalaxmi and Obe 2005) analysed studies published 1990-2003 and found a very mixed picture. Overall, studies with positive or negative, or inconclusive, findings were more or less equal in frequency.

By analyzing studies using combinations of ELF and other factors (chemical as well as physical) with known carcinogenic or mutagenic effects, a recent review suggests that effects of these co-exposures are far more frequently appearing in the literature than effects of pure ELF exposure (Juutilainen et al. 2006). This finding suggests a possible interaction of ELF magnetic fields with other agents. Furthermore, this review suggests that since effects frequently appear from 0.10 mT and higher, the radical pair mechanism (Brocklehurst and McLaughlan 1996) could explain the presence of positive findings at such flux densities.

Regarding more recent experimental findings, studies on genotoxic effects performed as part of the REFLEX project have received considerable attention. Different types of human and other mammalian cells (including human fibroblasts and lymphocytes) were exposed to a range of frequencies, flux densities and exposure regimes (Ivancsits et al. 2003a, Ivancsits et al. 2003b, Ivancsits et al. 2005, Winker et al. 2005). Chromosomal damage (DNA strand breaks, micronucleus formation) due to exposure was found in some, but not all cell types (e.g. lymphocytes not affected), after intermittent but not after continuous exposure. Flux density, frequency, and exposure time were important for observed effects, as well as age of cell donors. Similar studies have been performed to ascertain the replicability of the results. The outcome of these studies are at present not completely available and do thus not allow for final interpretation of the data, although at least one study could not confirm the initial findings (Scarfi et al. 2005). Other recent studies using human cells have also shown inconsistent results regarding DNA damage after ELF exposure (alone or in combination with chemical or other physical agents). These studies vary considerable both in exposure conditions and in techniques employed to test for clastogenic effects, making it difficult to draw firm conclusions at present. However, Mairs et al. (2007) recently showed that by using the very sensitive microsatellite sequence analysis, 50 Hz EMF at 1 mT could alone increase mutation rate in human glioma cells, as well as increase the mutagenic capacity of ionizing radiation. Also a study by Wahab et al. (2006) has recently indicated genotoxic actions of exposure to 50 Hz EMF. In this study it was seen that frequencies of sister chromatid exchanges were elevated in EMF exposed human lymphocytes. Any mechanism responsible for these possible genotoxic effects is not shown.

During the last years, there has been increased attention towards effects by ELF fields on free radical homeostasis as an indirect mechanism for several biological responses (Simkó and Mattsson 2004). Experiments with several cellular systems have shown that

Possible effects of Electromagnetic Fields (EMF) on Human Health

exposure leads to increased radical levels (e.g. Simkó et al. 2001, Rollwitz et al. 2004, Lupke et al. 2004). Interestingly, DNA damage in human cells (Wolf et al. 2005) exposed to ELF magnetic fields was counteracted by addition of antioxidants, suggesting that ELF magnetic fields can indirectly, possibly via changes in radical homeostasis, affect integrity of DNA.

Finally, based on data obtained with modern high-throughput screening methods and real-time PCR, Lupke et al. (2006) have suggested a comprehensive pathway by which ELF fields could influence cells of the immune system. The suggested pathway includes that membrane-associated events are affected by the fields, causing changes in radical homeostasis, and leading to down-stream events that include changes in gene expression, which could be of importance for regulation of proliferation.

Other biological endpoints relevant for carcinogenesis (e.g. cell cycle regulation, proliferation, apoptosis, gene expression) have been investigated in a number of studies. There are mixtures of positive and negative findings, not allowing for a general conclusion to be made regarding the overall potency for ELF fields to participate in the carcinogenic process. However, an interesting exception is three replication studies of an older study showing that low intensity 60 Hz MF can inhibit the antiproliferative effect of tamoxifen on a specific subclone of human MCF-7 breast cancer cells (Blackman et al. 2001, Ishido et al. 2001, Girgert et al. 2005). These are among the few EMF studies that have yielded reproducible results in several independent laboratories.

Discussion

The value of in vitro studies is in providing information on mechanisms of damage to cells and tissues. Published in vitro studies cannot explain epidemiological findings, but do not contradict them either. There is a need for independent replication of certain studies suggesting genotoxic effects and for better understanding of combined effects of ELF magnetic fields with other agents, their effects on free radical homeostasis, as well as of the possible implications of ELF field inhibition of tamoxifen effects. Studies with improved design are also needed.

3.5.3. Symptoms

What was already known on this subject?

A variety of symptoms (dermatological symptoms such as redness, tingling and burning sensations as well as for example fatigue, headache, concentration difficulties, nausea, heart palpitation) have been suggested to be caused by ELF field exposure. The term "electromagnetic hypersensitivity" (EHS) has come into common usage based on the reported experience by the afflicted individuals that electric and/or magnetic fields, or vicinity to activated electrical equipment trigger the symptoms.

In the CSTE opinion of 2001, the possibility of hypersensitivity in some individuals was said to require confirmation and the reports of such health problems did not provide a basis for changes in exposure limits.

What has been achieved since then?

Since the CSTE opinion of 2001 only few new provocation studies have been published on symptoms and ELF fields (for EHS and RF fields see Chapter 3.3.3). As stated in the WHO Fact sheet on electromagnetic hypersensitivity, well controlled and conducted double-blind studies have not shown any correlation between symptoms and EMF (WHO 2005). Rubin et al. (2005) reviewed 31 provocation studies (using different frequencies and EMF sources) testing more than 700 individuals reporting EHS (Rubin et al. 2005). The results in 24 of these studies did not support a relationship between the health problems and EMF. In seven of the other studies some supporting evidence was found,

Possible effects of Electromagnetic Fields (EMF) on Human Health

but in two cases the same research group failed to replicate their own findings. For another three studies Rubin and co-authors suspected that the results were statistical artefacts and in the final two studies the results were mutually incompatible.

Discussion

A relationship between ELF field exposure and symptoms has not been shown in scientific studies. From these results it seems clear that ELF field exposure is neither a necessary nor a sufficient factor to trigger health complaints in individuals reporting symptoms. Whether ELF fields may be a contributing factor under some conditions remains to be determined.

3.5.4. Other Health Effects

3.5.4.1. Epidemiology

Following the initial epidemiological study on childhood cancer a great number of other diseases have also been studied in relation to ELF fields. These diseases include cardiovascular disease, neurodegenerative disease and psychiatric disorders. An effect of heart rate variability seen in laboratory studies was the basis for a hypothesis that ELF exposure might affect the risk of cardiovascular disease and some initial epidemiologic results supported this. However, later well controlled studies have dismissed this hypothesis. For several of the other outcomes the support was never strong. Nevertheless, several neurodegenerative diseases are still considered worthy of study in this respect, and this refers particularly to ALS (amyotrophic lateral sclerosis) and Alzheimer disease (Ahlbom et al. 2001).

3.5.4.2. In vivo

What was already known on this subject?

The previous opinion did not evaluate evidence of health effects from animal studies. However, such data have been reviewed by IARC (2002) and ICNIRP (Bernhardt et al. 2003).

Nervous system and behaviour. While strong ELF fields are known to affect nerve and muscle cells and can be perceived, little evidence was found for effects on the nervous system or behaviour at environmental exposure levels. Effects of ELF magnetic fields on the EEG, cognition, behaviour and neurotransmitter levels have been described in a few studies, but there are inconsistencies in these data.

Reproduction and development. Several independent studies have suggested effects of ELF magnetic fields on the embryonic development of birds and other non-mammalian species, but the results are inconsistent. The evidence in mammalian species is restricted to minor skeletal anomalies seen in some studies with rats and mice. No consistent effects have been seen in any other reproductive or developmental endpoints in mammals. Minor skeletal variations are relatively common findings in teratological studies on rodents and often considered biologically insignificant.

Endocrine system. There is limited evidence of effects on melatonin production in experimental animals exposed to ELF magnetic fields, but such effects are not supported by other animal studies, and no statistically significant effects²⁴ have been seen on human volunteers under controlled laboratory conditions.

Other effects. No consistent evidence has been found for cardiovascular or immune system effects of ELF fields.

²⁴ 5%-significance level

Possible effects of Electromagnetic Fields (EMF) on Human Health

What has been achieved since then?

Two recent animal studies have provided evidence that ELF magnetic field exposure may affect melatonin production by modifying the response of dairy cows to the length of photoperiod (Rodriguez et al. 2004) and by affecting the sensitivity of mice to circadian light variations (Kumlin et al. 2005). The results of two new studies are interesting biological observations suggesting EMF interactions with the effects of light (photoperiod) on melatonin production. These observations may help to explain the inconsistencies of earlier research on EMFs and melatonin. However, the results of both studies are only suggestive and should be confirmed in further experiments. The suggested EMF effects on melatonin are subtle and apparently observable only in specific conditions. For these reasons, these results are not helpful for human health risk assessment.

Discussion

Although some studies have described ELF magnetic field effects on the nervous system, animal development and melatonin production, the evidence for such effects is weak and ambiguous. No conclusions concerning possible human health risks can be drawn from these data.

3.5.4.3. In vitro

What was already known on this subject?

Very few in vitro studies have been directed at answering the question if ELF fields are involved in the onset of other diseases than cancer (Portier and Wolfe 1998). Naturally, many basic cell and molecular studies were performed, mostly to understand more about fundamental interaction mechanisms, but also to understand how certain ELF fields can be used for therapeutic purposes (bone and wound healing especially).

What has been achieved since then and discussion

Few studies are available that directly address any specific disease or group of disease. This is partly due to that few diseases are characterised in such a way that specific disease models exist on the cell level, but also due to that ELF fields have not been convincingly shown to be involved in specific non-cancerous diseases. However, continuously there are reports showing that ELF fields during certain circumstances can give rise to cellular responses that are relevant for diseases of the nervous system, the immune system, endocrine organs, the skeleto-muscular apparatus, etc. Such studies do not at the present time allow extrapolation from the in vitro finding to any specific health state.

3.5.5. Conclusions about ELF fields

The previous opinion came to a similar conclusion regarding carcinogenicity of ELF fields as IARC's evaluation, namely that ELF magnetic fields are possibly carcinogenic. This conclusion was mainly based on epidemiologic results indicating that exposure to ELF fields might be a cause of childhood leukaemia. This assessment is still valid. The fact that the epidemiological results for childhood leukaemia have little support from known mechanisms or experimental studies is intriguing and it is a high research priority to reconcile these data.

For some other diseases, notably breast cancer and cardiovascular diseases, later research has indicated that an association is unlikely. For yet some other diseases, such as neurodegenerative disease and brain cancer, the issue of an association to ELF fields remains open and more research is called for. A relation between ELF fields and symptoms has not been demonstrated.

Of current interest is to arrive at a better understanding of recently published genotoxicity results including those from the REFLEX study.

3.6. Static fields

3.6.1. Sources and distribution of exposure in population

The number of artificial sources of static magnetic fields is small but there is a rapid development of technologies using static magnetic fields. The number of people with implants that can be affected by static magnetic fields is also growing. Static magnetic fields up to some mT are found in certain occupational settings, e.g., in the aluminium and chloralkali industries, in welding processes, and certain railway and underground systems. A very prominent application is MRI: different types of tissue in the human body can be identified and located by using static magnetic fields, magnetic gradients and RF fields. Close to the device a few hundred mT can be reached. Recently developed devices, currently only used by some research and specialised teams for specific applications, can use up to 10 T.

3.6.2. Health effects

There are only a few epidemiological studies available and the majority of these have focused on cancer risks. There are some reports on reproductive outcomes, and sporadic studies of other outcomes. Overall, few occupational studies have focused specifically on effects of static magnetic fields and exposure assessment has been poor. In summary, the available evidence from epidemiological studies is not sufficient to draw any conclusions about potential health effects of static magnetic field exposure (Feychting 2005b).

A large number of biological studies have been carried out in an effort to detect biological effects of static magnetic fields. The studies include in vitro and in vivo laboratory studies as well as studies on human volunteers. This research has been reviewed comprehensively in UNEP/WHO (2006). Known effects of magnetic fields are orientation of forces applied on biological molecules with magnetic properties: haemoglobin, rhodopsin (visual pigment), free radicals, nitric oxide; these effects are detectable at field levels of about 1 T, without known health consequences.

3.6.3. Conclusions about static fields

Adequate data for proper risk assessment of static magnetic fields are almost totally lacking. The advent of new technology, and in particular MRI equipment, makes it a priority for research.

3.7. Environmental Effects

What was already known on this subject?

The CSTEE opinion did not consider possible environmental impacts of EMF. It is noted that the majority of the relatively few published studies on environmental effects at the time of the CSTEE opinion were laboratory based using short exposure periods, in a single species. In addition some field investigations were reported around intense point sources of EMF, in particular overhead power cables.

Certain species have been recognised as likely to be particularly sensitive to EMF namely:

- species that are strongly dependent on magnetic fields for orientation/migration (migratory birds, certain fish and insects, bats etc) and/or possess electric sense organs (e.g. sharks and rays).

Possible effects of Electromagnetic Fields (EMF) on Human Health

- species with a high vulnerability to stress due to poorly developed or impaired defence mechanisms. For example animals with poor thermoregulation may be more vulnerable to the effects of high frequency EMF.

Nonetheless data to characterise this vulnerability and its implications have been very limited. Foster and Repacholi (2000) in their important review of the published data concluded that: 'attempts at environmental analysis of the effects of environmental EMF, with few exceptions have been scattered in focus, sporadic in publication and uneven in quality'.

The available data thus provided a seriously inadequate basis to assess the risk of EMF to environmental species. However, apart from some local minor effects no significant effects of EMF on environmental species were identified.

What has been achieved since then?

Despite the obvious need for some definitive studies there has been no significant increase in the volume or general quality of research activity in this area since. The majority of these studies have focussed on ELF fields.

There has however been a substantial shift in the form of the studies, in particular in the nature of the endpoints examined. Thus the majority of studies published before 2000 used visible endpoints that are obviously associated with an adverse effect. These had the advantage that their interpretation is quite straightforward. However such endpoints in many cases lack sensitivity. In the last few years an increasing number of studies on the effects of EMF have concentrated on the measurement of more sensitive biomarkers.

These have included:

- antioxidant status/ antioxidant enzyme measurements
- stress markers e.g. alanine (plants) and heat shock proteins (animals)
- changes in cell growth (e.g. meristems in plants)
- DNA changes (e.g. using the comet assay).

The majority of the few publications on the impact of EMF on environmental species have been in plants. The paper by Monselise et al. (2003) illustrates the use of new markers of cell/tissue change. These authors found that in duck weed, exposed in the laboratory to low intensity sinusoidally varying magnetic fields at 60 and 100Hz, an accumulation of alanine occurred. Alanine accumulation is found as a stress signal following many other kinds of stress. (NB This effect may have parallels with the formation of heat shock proteins in the mammalian kidney in response to various stressors). The authors postulate that this effect arose from free radical generation by the EMF.

Regoli et al. (2005) have reported the effect in snails of low frequency 50Hz EMF fields both in the laboratory and under overhead power cables. A range of biological markers was employed. They demonstrated that the EMF had particular effects on markers of oxidative stress such as catalase and glutathione reductase both in the laboratory and in the field situations. The time to an effect was shown to be dose dependant with effects in the field occurring even at low levels (after 40 days at 0.75 μ T). The authors attribute the effects to the generation of free radicals by the low frequency electromagnetic fields. The authors also observed a reduction in lysosomal stability and of DNA integrity (at 2.88 μ T under field conditions). However, no physical damage to the snails was reported.

These biomarkers do appear to be detecting changes at low, much more environmentally relevant field strengths. However, their interpretation in terms of species and ecosystem health is more challenging. Unfortunately these techniques have not focussed particularly on species that would be expected to be among the most sensitive to EMF.

Possible effects of Electromagnetic Fields (EMF) on Human Health

Using more classical endpoints Zaidi and Khatoon (2003) have studied the impact on pollen production of plants growing under overhead power cables using plants grown nearby as a control group. They found that plants growing under the high tension lines at higher voltages (132000 and 220000 volts) had some decrease in pollen fertility and that the pollen had a higher percentage of diads and diploid pollen grains which is an indicator of genetic change. This finding needs to be examined further.

Several studies have examined the impact of co-exposure to EMF and other stressors in plants. Thus Tafforeau et al. (2004) describe the impact of exposure to EMF combined with calcium deprivation, from either a GSM telephone or a single 2h exposure to 105GHz (from a Gunn oscillator) on meristem production in flax seedlings (i.e. increase in actively dividing cells in the hypocotyls of the growing seedling). An increase in meristem production was observed from each of these EMF sources. It should be noted however that no visible damage to the seedlings was observed in these studies and that other environmental stressors can also produce an increase in meristem production.

Yao et al. (2005) have examined the impact of EMF (0.2 and 0.45T) together with UV-B radiation on cucumber seedling growth. EMF alone produced an increase in seedling germination, seedling growth in parallel with an increase in lipid peroxidation. However in combination with UV-B seedling growth and development were significantly decreased.

These studies raise the question as to whether the impact of EMF may be additive with other significant environmental stressors in the field situation and if so, what are the practical consequences of this for individual plants and ecosystems. The data presently available are inadequate to assess this.

Discussion

The continued lack of good quality data in relevant species means that there are insufficient data to identify whether a single exposure standard is appropriate to protect all environmental species from EMF. Similarly the data are totally inadequate to judge whether the environmental standard(s) should be the same or significantly different from those appropriate to protect human health.

The demonstration that the impact of EMF may be additive with some other environmental stressors at least in plants needs further examination to gauge its practical significance.

At present it is not possible to draw any conclusions regarding human health from this data base. Nonetheless, long-term monitoring of the viability of carefully selected species and/or ecosystems may be valuable to gauge the potential of EMF to influence human health.

4. OPINION

Radio Frequency Fields (RF fields)

In its opinion from 2001 the CSTEE concluded regarding radiofrequency (RF) electromagnetic fields:

"The additional information which has become available on carcinogenic and other nonthermal effects of radiofrequency and microwave radiation frequencies in the last years does not justify a revision of exposure limits set by the Commission on the basis of the conclusions of the 1998 opinion of the Steering Scientific Committee. In particular, in humans, no evidence of carcinogenicity in either children or adults has resulted from epidemiological studies (the size of some of which was very large, although the period of observation was not long enough for a definitive statement). A relatively large series of laboratory studies has not provided evidence of genotoxicity. Subjective symptoms affecting some individuals possibly exist, but not enough information is available on: the levels of exposure producing such effect, on the features underlying individual susceptibility, on the possible biological mechanisms *or* the prevalence of susceptible individuals in different populations. Thus, current knowledge is insufficient for the implementation of measures aimed at the identification and protection of a highly sensitive sub-group of the population."

Based on the scientific rationale presented above the SCENIHR has updated the CSTEE opinion and concludes the following in regard to non-thermal effects:

The balance of epidemiologic evidence indicates that mobile phone use of less than 10 years does not pose any increased risk of brain tumour or acoustic neuroma. For longer use, data are sparse and any conclusions therefore are uncertain. From the available data, however, it does appear that there is no increased risk for brain tumours in long-term users, with the exception of acoustic neuroma for which there are some indications of an association.

For diseases other than cancer, very little epidemiologic data are available.

A particular consideration is mobile phone use by children. While no specific evidence exists, children or adolescents may be more sensitive to RF field exposure than adults in view of their continuing development. Children of today may also experience a much higher cumulative exposure than previous generations. To date no epidemiologic studies on children are available.

RF exposure has not consistently been shown to have an effect on self-reported symptoms (e.g. headache, fatigue, dizziness and concentration difficulties) or well-being.

Studies on neurological effects and reproductive effects have not indicated any health risks at exposure levels below the ICNIRP-limits established in 1998.

Animal studies have not provided evidence that RF fields could induce cancer, enhance the effects of known carcinogens, or accelerate the development of transplanted tumours. The open questions include adequacy of the experimental models used and scarcity of data at high exposure levels.

There is no consistent indication from in vitro research that RF fields affect cells at the nonthermal exposure level.

In conclusion, no health effect has been consistently demonstrated at exposure levels below the ICNIRP-limits established in 1998. However, the data base for this evaluation is limited especially for long-term low-level exposure.

Possible effects of Electromagnetic Fields (EMF) on Human Health

Intermediate Frequency Fields (IF fields)

In its opinion from 2001 the CSTEE did not comment specifically on intermediate frequencies (IF).

Based on the scientific rationale presented above the SCENIHR, however, updates the 2001 opinion with the following statement regarding intermediate frequencies:

Experimental and epidemiological data from the IF range are very sparse. Therefore, assessment of acute health risks in the IF range is currently based on known hazards at lower frequencies and at higher frequencies. Proper evaluation and assessment of possible health effects from long term exposure to IF fields are important because human exposure to such fields is increasing due to new and emerging technologies.

Extremely low frequency fields (ELF fields)

In its 2001 opinion the CSTEE reached the following conclusions regarding extremely low frequency (ELF) fields:

- "Combined analyses of the epidemiological studies on the association between exposure to ELF and childhood leukaemia have strengthened the evidence of an association. However, given some inconsistencies in exposure measurements and the absence of other criteria commonly used in assessing causality (particularly a plausible explanation of underlying biological mechanisms, see above), the association does not meet adequate criteria for being considered causal. Thus the overall evidence for 50/60 Hz magnetic fields to produce childhood leukaemia must be regarded as being limited.
- The effect, if any, seems to be limited to exposures above 0.4 μT . In European countries, the proportion of children exposed to such levels is less than 1%. Assuming that the risk is doubled among the exposed, in the general population this would roughly correspond to an excess incidence of less than 1% childhood leukaemia. To put this in context, in European countries, the incidence of leukaemia is around 45 per million children (age 0-14) per year.
- Whether changes of recommended exposure limits to 50/60 Hz magnetic fields (12) ought to be recommended on this basis is a problem for risk managers, falling beyond the remit of the CSTEE.
- There is no convincing suggestion of any other carcinogenic effect of ELF on either children or adults. Current information on this respect does not provide clues for reconsidering exposure limits.
- Reports on possibly hypersensitive individuals require confirmation and do not provide a basis for proposing changes in the exposure limits."

Based on the scientific rationale presented above the SCENIHR updates the previous opinion and concludes the following:

The previous conclusion that ELF magnetic fields are a possible carcinogen, chiefly based on childhood leukaemia results, is still valid. There is no generally accepted mechanism to explain how ELF magnetic field exposure may cause leukaemia. Animal studies have not provided adequate evidence for a causal relationship.

No consistent relationship between ELF fields and self-reported symptoms (sometimes referred to as electrical hypersensitivity) has been demonstrated.

Possible effects of Electromagnetic Fields (EMF) on Human Health

In addition, for breast cancer and cardiovascular disease, recent research has indicated that an association is unlikely. For neurodegenerative diseases and brain tumours, the link to ELF fields remains uncertain.

Static fields

In its opinion from 2001 the CSTE did not comment specifically on static magnetic fields.

Based on the scientific rationale presented above the SCENIHR, however, updates the 2001 opinion with the following statement regarding static magnetic fields:

Adequate data for proper risk assessment of static magnetic fields are very sparse. Developments of technologies involving static magnetic fields, e.g. with MRI equipment require risk assessments to be made in relation to the exposure of personnel.

Environmental Effects

The CSTE did not consider environmental effects in its opinion of 2001.

The continued lack of good quality studies in relevant species means that there are insufficient data to identify whether a single exposure standard is appropriate to protect all environmental species from EMF. Similarly the data are inadequate to judge whether the environmental standards should be the same or significantly different from those appropriate to protect human health.

Overall conclusion

The Committee is mindful of the mandate that requested particular attention to be paid to a wide variety of issues. In most cases the data available are very limited. Some of these issues will be addressed in further opinions as more data become available.

Research Recommendations

In view of the identified important gaps in knowledge the following research recommendations are being made.

RF fields

- A long term prospective cohort study. Such a study would overcome problems that were discussed in relation to existing epidemiological studies, including the Interphone study. These problems include recall bias and other aspects of exposure assessment, selection bias due to high proportions of non-responders, too short induction period, and restriction to intracranial tumours.
- Health effects of RF exposure in children. To date no study on children exists. This issue can also be addressed by studies on immature animals. This research has to take into consideration that dosimetry in children may differ from that in adults.
- Exposure distribution in the population. The advent of personal dosimeters has made it possible to describe individual exposure in the population and to assess the relative contribution of different sources to the total exposure. Such a project would require that groups of people with different characteristics are selected and that they wear dosimeters for a defined period of time.

Possible effects of Electromagnetic Fields (EMF) on Human Health

There are several experimental studies that need to be replicated. Examples are studies on genotoxicity and cognition involving sleep quality parameters. For studies on biomarkers it is essential that the impact on human health is considered. Valid exposure assessment including all relevant sources of exposure is essential. A general comment is that all studies must use high quality dosimetry.

IF fields

- Data on health effects from IF fields are sparse. This issue should be addressed both through epidemiologic and experimental studies.

ELF fields

- Epidemiological results indicate an increased risk of leukaemia in children exposed to high levels of ELF magnetic fields, however, this is not supported by animal data. The mechanisms responsible for the childhood leukaemia and the reasons for the discrepancy are unknown and require a better understanding and clarification.

Static fields

- A cohort study on personnel dealing with equipment that generates strong magnetic fields is required. The start of this would have to be a thorough feasibility study.
- Relevant experimental studies such as studies on carcinogenicity, genotoxicity as well as developmental and neurobehavioural effects would have to be conducted as well.

Additional considerations

- Studies including exposure to combinations of frequencies as well as combinations of electromagnetic fields and other agents need to be considered.

5. COMMENTS RECEIVED DURING THE PUBLIC CONSULTATION

Information about the public consultation has been circulated on a broad level to national authorities, international organizations, and other stakeholders. Many thoughtful and detailed responses to the opinion have been received. However, many of the submissions, while interesting, addressed aspects that were beyond the task given to the SCENIHR.

In evaluating the responses from the consultation, submitted material has only been considered for revision of the opinion if

- 1) it is directly referring to the content of the report and relating to the issues that the report addresses,
- 2) it contains specific comments and suggestions on the scientific basis of the opinion,
- 3) it is directly relevant to the task of updating the CSTEE opinion of 2001,
- 4) it refers to the peer-reviewed published literature from 2000 onwards and published in English, the pre-agreed working language of the SCENIHR and the working group,
- 5) it has the potential to add to the preliminary opinion of SCENIHR.

Information on individual cases or any other material not included in published reports was not considered.

Each submission which meets these criteria has been carefully considered by the Working Group. The opinion has been revised based on these comments.

None of the comments that were received during the public consultation process led to changes in the overall conclusions or in the opinion proper.

Many of the received comments were based on a misunderstanding of the purpose of the opinion and were based on the assumption that the opinion was in fact a comprehensive scientific review paper. Based on this assumption it was thought that every single scientific article of possible relevance to the area had to be cited. Other comments expected the opinion to comment upon exposure guidelines or risk management strategies such as the precautionary principle. This, however, was outside the scope of the opinion. The task given to the SCENIHR and the purpose of the opinion has now been stated more explicitly in the Introduction. Many other comments were based on misconceptions regarding the scientific method that is used for research in the context of risk evaluation. In order to help the readers to overcome this obstacle a methods section has been added.

Since about half a year has passed during the public consultation process, the literature has been updated with relevant publications up to the end of 2006 and this has not led to modifications in the conclusions of the opinion.

The version that was sent out for public consultation included a table that aimed to describe sources and levels of exposure to various types of fields. This table received quite a number of comments and has been revised as it was felt that it could be misinterpreted and not equally representing exposure scenarios in different European countries. However, this illustrates the fact that systematic comprehensive and systematic information about exposure is still scarce.

Quite a number of comments disagreed with the conclusions regarding whether or not exposure to electromagnetic fields is a cause of self-reported symptoms. The Committee does not doubt the existence of these symptoms. However, their association to EMF-

Possible effects of Electromagnetic Fields (EMF) on Human Health

exposure has been discussed thoroughly by the working group and no changes were made on the conclusions with this respect.

In addition, the comments showed that some sentences or paragraphs could be misinterpreted and that some wordings did not appropriately reflect the viewpoint of the working group or the SCENIHR. In those instances the text has been revised accordingly.

6. MINORITY OPINION

None

7. LIST OF ABBREVIATIONS

μT	microtesla
μW	microwatt
AC	Alternating current
AGNIR	Advisory Group on Non Ionising Radiation
ALS	Amyotrophic Lateral Sclerosis
AM	Amplitude modulation
B	Magnetic flux density
BBB	Blood-Brain-Barrier
cDNA	complementary DNA
CENELEC	European Committee for Electrotechnical Standardization
CI	Confidence Interval
CNS	Central Nervous System
CSTEE	Scientific Committee on Toxicity, Ecotoxicity and the Environment
CW	Continuous wave
DC	Direct current
DECT	Digital Enhanced Cordless Telephone
DMBA	7,12-dimethylbenz[a]anthracene
DNA	Deoxyribonucleic acid
DVB-T	Digital Terrestrial Television
EEG	Electroencephalogram
EHS	Electromagnetic hypersensitivity
ELF	Extremely low frequency
EMF	Electromagnetic field
EMF-NET	Coordination Action (funded under FP6) to provide a framework for coordination of results of research activities related to the biological effects of EMF, including occupational exposure.
f	Frequency
FP6	6 th Framework Programme
FM	Frequency Modulation
GSM	Global System for Mobile Communication
H	Magnetic field strength
HSP	Heat-shock proteins
Hz	Frequency in Hertz
IARC	International Agency for Research on Cancer
ICNIRP	International Committee on Non Ionising Radiation Protection
IF	Intermediate frequencies
IL	Interleukin
kg	Kilogram
kHz	Kilohertz
kV	Kilovolt
MCSF	Macrophage Colony-Stimulating Factor

Possible effects of Electromagnetic Fields (EMF) on Human Health

MF	Magnetic field
MHz	Megahertz
MRI	Magnetic Resonance Imaging
m²	square meter
ms	milliseconds
mT	millitesla
mW	milliwatt
NB	<i>nota bene</i> (note well)
NRPB	National Radiological Protection Board, United Kingdom
nT	nanotesla
nW	nanowatt
OR	Odds Ratio
PCR	Polymerase Chain Reaction
REFLEX-study	European Research Project of the 5th Framework Programme regarding the Risk Evaluation of Potential Environmental Hazards from Low Energy Electromagnetic Field (EMF) Exposure using sensitive in vitro methods
RF	Radio Frequency
ROS	Reactive Oxygen Species
RR	Relative Risk
SAR	Specific Absorption Rate
SCENIHR	Scientific Committee on Emerging and Newly Identified Health Risks
SD	Sprague-Dawley
SMS	Short Message Service
SSC	Scientific Steering Committee
SSI	Statens Strålskydds Institut (Swedish Radiation Protection Agency)
T	Tesla
TNF	Tumor Necrosis Factor
TNO	Nederlandse Organisatie voor Toegepast-Natuurwetenschappelijk Onderzoek (Netherlands Organisation for Applied Scientific Research)
UMTS	Universal Mobile Telephony System
UNEP	United Nations Environmental Programme
UV	Ultraviolet
UV-B	Ultraviolet-B
V	Volt
VDT	Video Display Terminals
VDU	Video display units (for computers, videos, TV and some measurement devices using cathode ray tubes)
W	Watt
WHO	World Health Organisation
WiMAX	Worldwide Interoperability for Microwave Access
WLAN	Wireless Local Area Network
yr	years

8. REFERENCES

- Ahlbom A, Day N, Feychting M, Roman E, Skinner J, Dockerty J, et al. A pooled analysis of magnetic fields and childhood leukaemia. *Br J Cancer* 2000; 83:692-698.
- Ahlbom A. Neurodegenerative diseases, suicide and depressive symptoms in relation to EMF. *Bioelectromagnetics* 2001; Suppl 5:S132-43.
- Ahlbom A, Green A, Kheifets L, Savitz D, Swerdlow A (ICNIRP Standing Committee on Epidemiology). Epidemiology of health effects of radiofrequency exposure. *Environ Health Perspect* 2004; 112:1741-54.
- Anane R, Dulou PE, Taxile M, Geffard M, Crespeau FL, Veyret B. Effects of GSM-900 microwaves on DMBA-induced mammary gland tumors in female Sprague-Dawley rats. *Radiat Res* 2003a; 160:492-7.
- Anane R, Geffard M, Taxile M, Bodet D, Billaudel B, Dulou PE, et al. Effects of GSM-900 microwaves on the experimental allergic encephalomyelitis (EAE) rat model of multiple sclerosis. *Bioelectromagnetics* 2003b; 24:211-3.
- Anderson LE, Boorman GA, Morris JE, Sasser LB, Mann PC, Grumbein SL, et al. Effect of 13 week magnetic field exposures on DMBA-initiated mammary gland carcinomas in female Sprague-Dawley rats. *Carcinogenesis* 1999; 20:1615-20.
- Anderson LE, Morris JE, Sasser LB, Löscher W. Effects of 50- or 60-hertz, 100 microT magnetic field exposure in the DMBA mammary cancer model in Sprague-Dawley rats: possible explanations for different results from two laboratories. *Environ Health Perspect* 2000; 108:797-802.
- Anderson LE, Sheen DM, Wilson BW, Grumbein SL, Creim JA, Sasser LB. Two-year chronic bioassay study of rats exposed to a 1.6 GHz radiofrequency signal. *Radiat Res* 2004; 162:201-10.
- Arai N, Enomoto H, Okabe S, Yuasa K, Kamimura Y, Ugawa Y. Thirty minutes mobile phone use has no short term adverse effects on central auditory pathway. *Clin Neurophysiol* 2003; 114:1390-4.
- Aran JM, Carrere N, Chalan Y, Dulou PE, Larrieu S, Letenneur L, et al. Effects of exposure of the ear to GSM microwaves: in vivo and in vitro experimental studies. *Int J Audiology* 2004; 43:245-54.
- Auvinen A, Hietanen M, Luukkonen R, Koskela RS. Brain tumours and salivary gland cancers among cellular telephone users. *Epidemiology* 2002; 13:356-9.
- Bak M, Sliwinska-Kowalska M, Zmyslony M, Dudarewicz A. No effects of acute exposure to the electromagnetic field emitted by mobile phones on brainstem auditory potentials in young volunteers. *Int J Occup Med Environ Health* 2003; 16:201-8.
- Bartsch H, Bartsch C, Seebald E, Deerberg F, Dietz K, Vollrath L, et al. Chronic exposure to a GSM-like signal (mobile phone) does not stimulate the development of DMBA-induced mammary tumors in rats: results of three consecutive studies. *Radiat Res* 2002; 157:183-90.
- Baum A, Mevissen M, Kamino K, Mohr U, Löscher W. A histopathological study on alterations in DMBA-induced mammary carcinogenesis in rats with 50 Hz, 100 μ T magnetic field exposure. *Carcinogenesis* 1995; 16:119-25.

Possible effects of Electromagnetic Fields (EMF) on Human Health

Beckman KB, Ames BN. The free radical theory of aging matures. *Physiol Rev* 1998; 78:547-81.

Berg G, Schüz J, Samkange-Zeeb F, Blettner M. Assessment of radiofrequency exposure from cellular telephone daily use in an epidemiological study: German Validation study of the international case-control study of cancers of the brain-INTERPHONE-Study. *J Expo Anal Environ Epidemiol* 2005; 15:217-24.

Bernhardt JH, Matthes R, Repacholi MH, editors. Non-Thermal effects of RF electromagnetic fields. Proceedings of the International Seminar on Biological Effects of Non-Thermal Pulsed and Amplitude Modulated RF Electromagnetic Fields and Related Health Risks; 1996 Nov 20-21; Munich, Germany. Munich: International Commission on Non-Ionizing Radiation Protection; 1997.

Bernhardt JH, Matthes R, McKinlay A, Vecchia P, Veyret B, editors. Exposure to Static and Low Frequency Electromagnetic Fields, Biological Effects and Health Consequences (0-100 kHz) - Review of the Scientific Evidence and Health Consequences. Munich: International Commission on Non-Ionizing Radiation Protection; 2003.

Besset A, Espa F, Dauvilliers Y, Billiard M, de Seze R. No effect on cognitive function from daily mobile phone use. *Bioelectromagnetics* 2005; 26:102-8.

Black DR, Heynick LN. Radiofrequency (RF) effects on blood cells, cardiac, endocrine, and immunological functions. *Bioelectromagnetics* 2003; Suppl 6:S187-95.

Blackman CF, Benane SG, House DE. The influence of 1.2 microT, 60 Hz magnetic fields on melatonin- and tamoxifen-induced inhibition of MCF-7 cell growth. *Bioelectromagnetics* 2001; 22:122-8.

Boorman GA, Anderson LE, Morris JE, Sasser LB, Mann PC, Grumbein SL, et al. Effect of 26 week magnetic field exposures in a DMBA initiation-promotion mammary gland model in Sprague-Dawley rats. *Carcinogenesis* 1999; 20:899-904.

Braune S, Wrocklage C, Raczek J, Gailus T, Lucking CH. Resting blood pressure increase during exposure to a radio-frequency electromagnetic field. *Lancet* 1998; 351:1857-8.

Braune S, Riedel A, Schulte-Monting J, Raczek J. Influence of a radiofrequency electromagnetic field on cardiovascular and hormonal parameters of the autonomic nervous system in healthy individuals. *Radiation Research* 2002; 158:352-6.

Brocklehurst B, McLauchlan KA. Free radical mechanism for the effects of environmental electromagnetic fields on biological systems. *Int J Radiat Biol* 1996; 69:3-24.

Capri M, Scarcella E, Bianchi E, Fumelli C, Mesirca P, Agostini C, et al. 1800 MHz radiofrequency (mobile phones, different Global System for Mobile communication modulations) does not affect apoptosis and heat shock protein 70 level in peripheral blood mononuclear cells from young and old donors. *Int J Radiat Biol* 2004a; 80:389-97.

Capri M, Scarcella E, Fumelli C, Bianchi E, Salvioli S, Mesirca P et al. In vitro exposure of human lymphocytes to 900 MHz CW and GSM modulated radiofrequency: studies of proliferation, apoptosis and mitochondrial membrane potential. *Radiat Res* 2004b; 162:211-8.

Caraglia M, Marra M, Mancinelli F, D'Ambrosio G, Massa R, Giordano A et al. Electromagnetic fields at mobile phone frequency induce apoptosis and inactivation of the

Possible effects of Electromagnetic Fields (EMF) on Human Health

multi-chaperone complex in human epidermoid cancer cells. *J Cell Physiol* 2005; 204:539-48.

Cassel JC, Cosquer B, Galani R, Kuster N. Whole-body exposure to 2.45 GHz electromagnetic fields does not alter radial-maze performance in rats. *Behav Brain Res* 2004; 155:37-43.

Christ A, Kuster N. Differences in RF energy absorption in the heads of adults and children. *Bioelectromagnetics* 2005; Suppl 7:S31-44.

Christensen HC, Schüz J, Kosteljanetz M, Poulsen HS, Thomsen J, Johansen C. Cellular telephone use and risk of acoustic neuroma. *Am J Epidemiol* 2004; 159:277-83.

Christensen HC, Schüz J, Kosteljanetz M, Poulsen HS, Boice JD Jr., McLaughlin JK, et al. Cellular telephones and risk for brain tumours: a population-based, incident case-control study. *Neurology* 2005; 64:1189-95.

Cobb B, Jauchem J, Adair E. Radial arm maze performance of rats following repeated low level microwave radiation exposure. *Bioelectromagnetics*, 2004; 25:49-57.

Cook CM, Thomas AW, Prato FS. Human electrophysiological and cognitive effects of exposure to ELF magnetic and ELF modulated RF and microwave fields: A review of recent studies. *Bioelectromagnetics* 2002; 23:144-57.

Cotgreave IA. Biological stress responses to radio frequency electromagnetic radiation: are mobile phones really so (heat) shocking? *Arch Biochem Biophys* 2005; 435:227-40.

Curcio G, Ferrara M, De Gennaro L, Cristiani R, D'Inzeo G, Bertini M. Time-course of electromagnetic field effects on human performance and tympanic temperature. *Neuroreport* 2004; 15:161-4.

D'Andrea JA, Chou CK, Johnston SA, Adair ER. Microwave effects on the nervous system. *Bioelectromagnetics* 2003; Suppl 6:S107-47.

Dawe AS, Smith B, Thomas DW, Greedy S, Vasic N, Gregory A et al. A small temperature rise may contribute towards the apparent induction by microwaves of heat-shock gene expression in the nematode *Caenorhabditis Elegans*. *Bioelectromagnetics* 2006; 27:88-97.

D'Costa H, Trueman G, Tang L, Abdel-Rahman U, Abdel-Rahman W, Ong K, et al. Human brain wave activity during exposure to radiofrequency field emissions from mobile phones. *Australas Phys Eng Sci Med* 2003; 26:162-7.

de Pomerai D, Daniells C, David H, Allan J, Duce I, Mutwakil M, Thomas D, et al. Non-thermal heat-shock response to microwaves. *Nature* 2000; 405:417-8.

Diem E, Schwarz C, Adlkofer F, Jahn O, Rudiger H. Non-thermal DNA breakage by mobile-phone radiation (1800 MHz) in human fibroblasts and in transformed GFSH-R17 rat granulosa cells in vitro. *Mutat Res* 2005; 583:178-83.

Dubreuil D, Jay T, Edeline JM. Head only exposure to GSM 900 MHz EMF does not alter rats memory in spatial and non spatial tasks. *Behav Brain* 2003; 145:51-61.

Fedrowitz M, Westermann J, Löscher W. Magnetic field exposure increases cell proliferation but does not affect melatonin levels in the mammary gland of female Sprague Dawley rats. *Cancer Res* 2002; 62:1356-63.

Possible effects of Electromagnetic Fields (EMF) on Human Health

Fedrowitz M, Kamino K, Löscher W. Significant differences in the effects of magnetic field exposure on 7,12-dimethylbenz(a)anthracene-induced mammary carcinogenesis in two substrains of Sprague-Dawley rats. *Cancer Res* 2004; 64:243-51.

Feychting M. Non-cancer EMF effects related to children. *Bioelectromagnetics* 2005a; Suppl 7:S69-74.

Feychting M. Health effects of static magnetic fields – a review of the epidemiological evidence. *Prog Biophys Mol Biol* 2005b; 87:241-6.

Finnie JW, Blumbergs PC, Manavis J, Utteridge TD, GebSKI V, Swift JG, et al. Effect of global system for mobile communication (GSM) like radiofrequency fields on vascular permeability in mouse brain. *Pathology* 2001; 33:338-40.

Finnie JW, Blumbergs PC, Manavis J, Utteridge TD, GebSKI V, Davies RA, et al. Effect of long-term mobile communication microwave exposure on vascular permeability in mouse brain. *Pathology* 2002; 34:344-7.

Foliart DE, Pollock BH, Mezei G, Iriye R, Silva JM, Ebi KL, et al. Magnetic field exposure and long-term survival among children with leukaemia. *Br J Cancer* 2006; 94:161-4.

Forsen UM, Rutqvist LE, Ahlbom A, Feychting M. Occupational magnetic fields and female breast cancer: a case-control study using Swedish population registers and new exposure data. *Am J Epidemiol* 2005; 161:250-9.

Foster KR, Repacholi MH. Environmental impacts of electromagnetic fields from major electrical technologies. Geneva: WHO; 2000. http://www.who.int/peh-emf/publications/reports/en/env_impact_emf_from_major_elect_tech_foster_repacholi.pdf (accessed 24 April 2007).

French PW, Penny R, Laurence JA, McKenzie DR. Mobile phones, heat shock proteins and cancer. *Differentiation* 2001; 67:93-7.

Fritze K, Sommer C, Schmitz B, Mies G, Hossmann KA, Kiessling M, et al. Effect of global system for mobile communication (GSM) microwave exposure on blood-brain barrier permeability in rat. *Acta Neuropathol (Berl)* 1997; 94:465-70.

Girgert R, Schimming H, Korner W, Grundker C, Hanf V. Induction of tamoxifen resistance in breast cancer cells by ELF electromagnetic fields. *Biochem Biophys Res Commun* 2005; 336:1144-9.

Glaser R. In-vitro studies of electromagnetic exposure between 300 Hz and 10 MHz. In: Matthes R, van Rongen E, Repacholi MH, editors. *Health Effects of Electromagnetic Fields in the Frequency Range 300 Hz to 10 MHz*. Proceedings of the International Seminar on Health Effects of Electromagnetic Fields in the Frequency Range 300 Hz to 10 MHz; 1999 June 7-8; Maastricht, The Netherlands. Munich: International Commission on Non-Ionizing Radiation Protection; 1999. p.105-21.

Greenland S, Sheppard AR, Kaune WT, Poole C, Kelsh MA. A pooled analysis of magnetic fields, wire codes, and childhood leukemia. *Childhood Leukemia-EMF Study Group. Epidemiology* 2000; 11:624-34.

Gotoh T, Terada K, Mori M. hsp70-DnaJ chaperone pairs prevent nitric oxide-mediated apoptosis in RAW 264.7 macrophages. *Cell Death Differ* 2001; 8:357-66.

Hardell L, Nasman A, Pahlson A, Hallquist A, Hansson Mild K. Use of cellular telephones and the risk for brain tumours: A case-control study. *Int J Oncol* 1999; 15:113-6.

Possible effects of Electromagnetic Fields (EMF) on Human Health

Hardell L, Hallquist A, Mild KH, Carlberg M, Pahlson A, Lilja A. Cellular and cordless telephones and the risk for brain tumours. *Eur J Cancer Prev* 2002; 11:377-86.

Hardell L, Carlberg M, Hansson Mild K. Case-control study on cellular and cordless telephones and the risk for acoustic neuroma or meningioma in patients diagnosed 2000-2003. *Neuroepidemiology* 2005a; 25:120-8.

Hardell L, Carlberg M, Mild KH. Case-control study of the association between the use of cellular and cordless telephones and malignant brain tumours diagnosed during 2000-2003. *Environ Res* 2005b; 100:232-41.

Hardell L, Mild KH, Carlberg M, Söderqvist F. Tumour risk associated with use of cellular telephones or cordless desktop telephones. *World J Surg Oncol* 2006; 4:74.

Heikkinen P, Kosma VM, Hongisto T, Huuskonen H, Hyysalo P, Komulainen H, et al. Effects of mobile phone radiation on X-ray-induced tumorigenesis in mice. *Radiat Res* 2001; 156:775-85.

Heikkinen P, Kosma VM, Alhonen L, Huuskonen H, Komulainen H, Kumlin T, et al. Effects of mobile phone radiation on UV-induced skin tumourigenesis in ornithine decarboxylase transgenic and non-transgenic mice. *Int J Radiat Biol* 2003; 79:221-33.

Heikkinen P, Ernst H, Huuskonen H, Komulainen H, Kumlin T, Mäki-Paakkanen J, et al. Effects of radiofrequency (RF) radiation on 3-chloro-4-(dichloromethyl)-5-hydroxy-2(5H)-furanone (MX)-induced tumorigenesis in Wistar rats. *Radiat Res* 2006; in press.

Hepworth SJ, Schoemaker MJ, Muir KR, Swerdlow AJ, van Tongeren MJ, McKinney PA. Mobile phone use and risk of glioma in adults: case-control study. *BMJ* 2006; 332:883-7.

Heynick LN, Merritt JH. Radiofrequency fields and teratogenesis. *Bioelectromagnetics* 2003; Suppl 6:S174-86.

Hietanen M. Review of epidemiological studies at intermediate frequencies. In: Matthes R, van Rongen E, Repacholi MH (eds.). *Health Effects of Electromagnetic Fields in the Frequency Range 300 Hz to 10 MHz*. Proceedings of the International Seminar on Health Effects of Electromagnetic Fields in the Frequency Range 300 Hz to 10 MHz; 1999 June 7-8; Maastricht, The Netherlands. Munich: International Commission on Non-Ionizing Radiation Protection; 1999. p.147-54.

Higashikubo R, Ragouzis M, Moros EG, Straube WL, Roti JL. Radiofrequency electromagnetic fields do not alter the cell cycle progression of C3H 10T and U87MG cells. *Radiat Res* 2001; 156:786-95.

Hook GJ, Zhang P, Lagroye I, Li L, Higashikubo R, Moros EG, et al. Measurement of DNA damage and apoptosis in Molt-4 cells after in vitro exposure to radiofrequency radiation. *Radiat Res* 2004; 161:193-200.

Hossmann KA, Hermann DM. Effects of electromagnetic radiation of mobile phones on the central nervous system. *Bioelectromagnetics* 2003; 24:49-62.

Huang TQ, Lee JS, Kim TH, Pack JK, Jang JJ, Seo JS. Effect of radiofrequency radiation exposure on mouse skin tumorigenesis initiated by 7,12-dimethylbenz[alpha]anthracene. *Int J Radiat Biol* 2005; 81:861-7.

Possible effects of Electromagnetic Fields (EMF) on Human Health

Huber R, Graf T, Cote KA, Wittmann L, Gallmann E, Matter D, et al. Exposure to pulsed high-frequency electromagnetic field during waking affects human sleep EEG. *Neuroreport* 2000; 11:3321-5.

Hutter HP, Moshammer H, Wallner P, Kundi M. Subjective symptoms, sleeping problems, and cognitive performance in subjects living near mobile phone base stations. *Occup Environ Med* 2006; 63:307-16.

Huuskonen H, Lindbohm ML, Juutilainen J. Teratogenic and reproductive effects of low-frequency magnetic fields. *Mutat Res* 1998; 410:167-83.

IARC (International Agency for Research on Cancer). Non-Ionizing Radiation, Part 1: Static and extremely low-frequency (ELF) electric and magnetic fields. IARC Monographs on the Evaluation of carcinogenic Risks to Humans: Volume 80. Lyon: IARC Press; 2002.

IARC (International Agency for Research on Cancer). IARC Monographs on the Evaluation of Carcinogenic Risks to Humans – Preamble. 2006
<http://monographs.iarc.fr/ENG/Preamble/CurrentPreamble.pdf> (accessed 24.4.2007)

Imaida K, Kuzutani K, Wang J, Fujiwara O, Ogiso T, Kato K, et al. Lack of promotion of 7,12-dimethylbenz[a]anthracene-initiated mouse skin carcinogenesis by 1.5 GHz electromagnetic near fields. *Carcinogenesis* 2001; 22:1837-41.

Inoue Y, Sato Y, Nishimura M, Seguchi M, Zaitzu Y, Yamada K, et al. Heat-induced drug resistance is associated with increased expression of Bcl-2 in HL60. *Anticancer Res* 1999; 19:3989-92.

Inskip PD, Tarone RE, Hatch EE, Wilcosky TC, Shapiro WR, et al. Cellular-telephone use and brain tumours. *N Engl J Med* 2001; 344:79-86.

Ishido M, Nitta H, Kabuto M. Magnetic fields (MF) of 50 Hz at 1.2 microT as well as 100 microT cause uncoupling of inhibitory pathways of adenylyl cyclase mediated by melatonin 1a receptor in MF-sensitive MCF-7 cells. *Carcinogenesis* 2001; 22:1043-8.

Ivancsits S, Diem E, Jahn O, Rudiger HW. Age-related effects on induction of DNA strand breaks by intermittent exposure to electromagnetic fields. *Mech Ageing Dev* 2003a; 124:847-50.

Ivancsits S, Diem E, Jahn O, Rudiger HW. Intermittent extremely low frequency electromagnetic fields cause DNA damage in a dose-dependent way. *Int Arch Occup Environ Health* 2003b; 76:431-6.

Ivancsits S, Pilger A, Diem E, Jahn O, Rudiger HW. Cell type-specific genotoxic effects of intermittent extremely low-frequency electromagnetic fields. *Mutat Res* 2005; 583:184-8.

Johansen C, Boice J Jr., McLaughlin J, Olsen J. Cellular telephones and cancer-a nationwide cohort study in Denmark. *J Natl Cancer Inst* 2001; 93:203-7.

Jolly C, Morimoto RI. Role of the heat shock response and molecular chaperones in oncogenesis and cell death. *J Natl Cancer Inst* 2000; 92:1564-72.

Juutilainen J, Eskelinen T. In vivo studies on the health effects of electromagnetic fields in the frequency range 300 Hz to 10 MHz. In: Matthes R, van Rongen E, Repacholi MH (eds.). Health Effects of Electromagnetic Fields in the Frequency Range 300 Hz to 10 MHz. Proceedings of the International Seminar on Health Effects of Electromagnetic Fields in the Frequency Range 300 Hz to 10 MHz; 1999 June 7-8; Maastricht, The

Possible effects of Electromagnetic Fields (EMF) on Human Health

Netherlands. Munich: International Commission on Non-Ionizing Radiation Protection; 1999. p.123-30.

Juutilainen J, Lang S, Rytomaa T. Possible cocarcinogenic effects of ELF electromagnetic fields may require repeated long-term interaction with known carcinogenic factors. *Bioelectromagnetics* 2000; 21:122-8.

Juutilainen J. Developmental effects of electromagnetic fields. *Bioelectromagnetics* 2005; Suppl 7:S107-15.

Juutilainen J, Kumlin T, Naarala J. Do extremely low frequency magnetic fields enhance the effects of environmental carcinogens? A meta-analysis of experimental studies. *Int J Radiat Biol* 2006; 82:1-12.

Kheifets L, Repacholi M, Saunders R, van Deventer E. The sensitivity of children to electromagnetic fields. *Pediatrics* 2005; 116:e303-13.

Koivisto M, Krause CM, Revonsuo A, Laine M, Hamalainen H. The effects of electromagnetic field emitted by GSM phones on working memory. *Neuroreport* 2000a; 11:1641-3.

Koivisto M, Revonsuo A, Krause C, Haarala C, Sillanmaki L, Laine M, et al. Effects of 902 MHz electromagnetic field emitted by cellular telephones on response times in humans. *Neuroreport* 2000b; 11:413-5.

Koivisto M, Haarala C, Krause CM, Revonsuo A, Laine M, Hämäläinen H. GSM phone signals does not produce subjective symptoms. *Bioelectromagnetics* 2001; 22:212-5.

Krause CM, Sillanmaki L, Koivisto M, Haggqvist A, Saarela C, Revonsuo A, et al. Effects of electromagnetic field emitted by cellular phones on the EEG during a memory task. *Neuroreport* 2000; 11:761-4.

Kumlin T, Kosma VM, Alhonen L, Janne J, Komulainen H, Lang S, et al. Effects of 50 Hz magnetic fields on UV-induced skin tumorigenesis in ODC-transgenic and non-transgenic mice. *Int J Radiat Biol* 1998; 73:113-21.

Kumlin T, Heikkinen P, Kosma VM, Alhonen L, Janne J, Juutilainen J. p53-independent apoptosis in UV-irradiated mouse skin: possible inhibition by 50 Hz magnetic fields. *Radiat Environ Biophys* 2002; 41:155-8.

Kumlin T, Heikkinen P, Laitinen JT, Juutilainen J. Exposure to a 50-Hz Magnetic Field Induces a Circadian Rhythm in 6-hydroxymelatonin Sulfate Excretion in Mice. *J Radiat Res (Tokyo)* 2005; 46:313-8.

Kwee S, Raskmark P, Velizarov S. Changes in cellular proteins due to environmental non-ionizing radiation. 1. Heat shock proteins. *Electro- and Magnetobiology* 2001; 20:141-52.

La Regina M, Moros EG, Pickard WF, Straube WL, Baty J, Roti JL. The effect of chronic exposure to 835.62 MHz FDMA or 847.74 MHz CDMA radiofrequency radiation on the incidence of spontaneous tumors in rats. *Radiat Res* 2003; 160:143-51.

Lai H, Horita A, Guy AW. Microwave irradiation affects radial-arm maze performance in the rat. *Bioelectromagnetics* 1994; 15:95-104.

Lai H. Interactions of MW and temporally incoherent magnetic field on spatial learning in rat. *Physiology and Behavior* 2004a; 84:785-9.

Possible effects of Electromagnetic Fields (EMF) on Human Health

Lai H, Singh NP. Magnetic-field-induced DNA strand breaks in brain cells of the rat. *Environ Health Perspect* 2004b; 112:687-94.

Lantow M, Lupke M, Frahm J, Mattsson MO, Kuster N, Simkó M. ROS release and Hsp70 expression after exposure to 1800 MHz radiofrequency electromagnetic fields in primary human monocytes and lymphocytes. *Radiat Environ Biophys* 2006a; 45:55-62.

Lantow M, Schuderer J, Hartwig C, Simkó M. Free radical release and Hsp70 expression in two human immune relevant cell lines after exposure to 1800 MHz radio frequency radiation. *Radiat Res* 2006b; 165:88-94.

Lantow M, Viergutz T, Weiss DG, Simkó M. Comparative study of cell cycle kinetics and induction of apoptosis or necrosis after exposure to radiofrequency radiation in human Mono Mac 6 cells. *Radiat Res* 2006c; 166:539-43.

Leszczynski D, Joenvaara S, Reivinen J, Kuokka R. Non-thermal activation of the hsp27/p38MAPK stress pathway by mobile phone radiation in human endothelial cells: molecular mechanism for cancer and blood-brain barrier-related effects. *Differentiation* 2002; 70:120-9.

Litvak E, Repacholi MH. Gaps in knowledge about effects from exposure to EMF in the frequency range 300 Hz to 10 MHz. In: Matthes R, van Rongen E, Repacholi MH (eds.). *Health Effects of Electromagnetic Fields in the Frequency Range 300 Hz to 10 MHz. Proceedings of the International Seminar on Health Effects of Electromagnetic Fields in the Frequency Range 300 Hz to 10 MHz; 1999 June 7-8; Maastricht, The Netherlands. Munich: International Commission on Non-Ionizing Radiation Protection; 1999. p.169-175.*

Litvak E, Foster KR, Repacholi MH. Health and Safety Implications of Exposure to Electromagnetic Fields in the Frequency Range 300 Hz to 10 MHz. *Bioelectromagnetics* 2002; 23:68-82.

Lönn S, Ahlbom A, Hall P, Feychting M. Mobile phone use and the risk of acoustic neuroma. *Epidemiology* 2004; 15:653-9.

Lönn S, Ahlbom A, Hall P, Feychting M. Long-term mobile phone use and brain tumour risk. *Am J Epidemiol* 2005; 161:526-35.

Löscher W, Mevissen M, Lehmacher W, Stamm A. Tumor promotion in a breast cancer model by exposure to a weak alternating magnetic field. *Cancer Lett* 1993; 71:75-81.

Löscher W. Do cocarcinogenic effects of ELF electromagnetic fields require repeated long-term interaction with carcinogens? Characteristics of positive studies using the DMBA breast cancer model in rats. *Bioelectromagnetics* 2001; 22:603-14.

Lupke M, Rollwitz T, Simkó M. Cell activating capacity of 50 Hz magnetic fields to release reactive oxygen intermediates in human umbilical cord blood-derived monocytes and in Mono Mac 6 cells. *Free Radic Res* 2004; 38:985-93.

Lupke M, Frahm J, Lantow M, Maercker C, Remondini D, Bersani F, et al. Gene expression analysis of ELF-MF exposed human monocytes indicating the involvement of the alternative activation pathway. *Biochim Biophys Acta* 2006; 1763:402-12.

Mairs RJ, Hughes K, Fitzsimmons S, Prise KM, Livingstone A, Wilson L, et al. Microsatellite analysis for determination of the mutagenicity of extremely low-frequency electromagnetic fields and ionising radiation in vitro. *Mutat Res* 2007; 626:34-41.

Possible effects of Electromagnetic Fields (EMF) on Human Health

Mann K, Roschke J. Effects of pulsed high-frequency electromagnetic fields on human sleep. *Neuropsychobiology* 1996; 33:41-7.

Marinelli F, La Sala D, Ciccio G, Cattini L, Trimarchi C, Putti S, et al. Exposure to 900 MHz electromagnetic field induces an unbalance between pro-apoptotic and pro-survival signals in T-lymphoblastoid leukaemia CCRF-CEM cells. *J Cell Physiol* 2004; 198:324-32.

Mevissen M, Lerchl A, Szamel M, Löscher W. Exposure of DMBA-treated female rats in a 50-Hz, 50 microTesla magnetic field: effects on mammary tumor growth, melatonin levels, and T lymphocyte activation. *Carcinogenesis* 1996; 17:903-10.

Mevissen M, Haussler M, Lerchl A, Löscher W. Acceleration of mammary tumorigenesis by exposure of 7,12-dimethylbenz[a]anthracene-treated female rats in a 50-Hz, 100-microT magnetic field: replication study. *J Toxicol Environ Health A* 1998; 53:401-18.

Miyakoshi J, Takemasa K, Takashima Y, Ding GR, Hirose H, Koyama S. Effects of exposure to a 1950 MHz radio frequency field on expression of Hsp70 and Hsp27 in human glioma cells. *Bioelectromagnetics* 2005; 26:251-7.

Monfrecola G, Moffa G, Procaccini EM. Non-ionizing electromagnetic radiations, emitted by a cellular phone, modify cutaneous blood flow. *Dermatology* 2003; 207:10-4.

Monselise EB, Parola AH, Kost D. Low frequency electromagnetic fields induce a stress effect upon higher plants, as evident by the universal stress indicator, alanine. *Biochem Biophys Acta* 2003; 302:427-34.

Moulder JE, Erdreich LS, Malyapa RS, Merritt J, Pickard WF, Vijayalaxmi. Cell phones and cancer: what is the evidence for a connection? *Radiat Res* 1999; 151:513-31.

Muscat JE, Malkin MG, Thompson S, Shore RE, Stellman SD, McRee D, et al. Handheld cellular telephone use and risk of brain cancer. *JAMA* 2000; 284:3001-7.

Muscat JE, Malkin MG, Shore RE, Thompson S, Neugut AI, Stellman SD, et al. Handheld cellular telephones and risk of acoustic neuroma. *Neurology* 2002; 58:1304-6.

Neubauer C, Phelan AM, Kues H, Lange DG. Microwave irradiation of rats at 2.45 GHz activates pinocytotic-like uptake of tracer by capillary endothelial cells of cerebral cortex. *Bioelectromagnetics* 1990; 11:261-8.

Nikolova T, Czyz J, Rolletschek A, Blyszczuk P, Fuchs J, Jovtchev G, et al. Electromagnetic fields affect transcript levels of apoptosis-related genes in embryonic stem cell-derived neural progenitor cells. *FASEB J* 2005; 19:1686-8.

Ozturan O, Erdem T, Miman MC, Kalcioğlu MT, Oncel S. Effects of the electromagnetic field of mobile telephones on hearing. *Acta Otolaryngol* 2002; 122:289-93.

Pacini S, Ruggiero M, Sardi I, Aterini S, Gulisano F, Gulisano M. Exposure to global system for mobile communication (GSM) cellular phone radiofrequency alters gene expression, proliferation, and morphology of human skin fibroblasts. *Oncol Res* 2002; 13:19-24.

Parazzini M, Bell S, Thuroczy G, Molnar F, Tognola G, Lutman ME, et al. Influence on the mechanisms of generation of distortion product otoacoustic emissions of mobile phone exposure. *Hear Res* 2005; 208:68-78.

Paredi P, Kharitonov SA, Hanazawa T, Barnes PJ. Local vasodilator response to mobile phones. *Laryngoscope* 2001; 111:159-62.

Possible effects of Electromagnetic Fields (EMF) on Human Health

Persson BRR, Salford LG, Brun A. Blood-brain barrier permeability in rats exposed to electromagnetic fields used in wireless communication. *Wireless Networks* 1997; 3:455-61.

Portier CJ, Wolfe MS, editors. Assessment of health effects from exposure to power-line frequency electric and magnetic fields. Research Triangle Park, NC: National Institute of Environmental Health Sciences, National Institutes of Health; 1998. (Working Group report) <http://www.niehs.nih.gov/emfrapid/html/WGReport/emf.pdf> (accessed 24.4.2007).

Preece AW, Iwi G, Davies-Smith A, Wesnes K, Butler S, Lim E, et al. Effect of a 915-MHz simulated mobile phone signal on cognitive function in man. *Int J Radiat Biol* 1999; 75:447-56.

Purves D, Augustine GJ, Fitzpatrick D, Katz LC, LaMantia AS, McNamara JO, et al. *Neuroscience*. 2nd ed. Sunderland (MA): Sinauer Associates, Inc.; 2001.

Regel SJ, Negovetic S, Rööslü M, Berdinas V, Schuderer J, Huss A, et al. UMTS base station-like exposure, well being and cognitive performance. *Environ Health Perspect* 2006; 114:1270-5.

Regoli F, Gorbi S, Machella N, Tedesco S, Benedetti M, Bocchetti R, et al. Pro-oxidant effects of extremely low frequency electromagnetic fields in the land snail *helix aspersa*. *Free Radic Biol Med* 2005; 39:1620-8.

Remondini D, Nylund R, Reivinen J, Poullétier de Gannes F, Veyret B, Lagroye I,. Gene expression changes in human cells after exposure to mobile phone microwaves. *Proteomics* 2006; 6:4745-54.

Repacholi MH, Basten A, Gebiski V, Noonan D, Finnie J, Harris AW. Lymphomas in E mu-Pim1 transgenic mice exposed to pulsed 900 MHz electromagnetic fields. *Radiat Res* 1997; 147:631-40.

Rodriguez M, Petitclerc D, Burchard JF, Nguyen DH, Block E. Blood melatonin and prolactin concentrations in dairy cows exposed to 60 Hz electric and magnetic fields during 8 h photoperiods. *Bioelectromagnetics* 2004; 25:508-15.

Roelandts R. Cellular phones and the skin. *Dermatology* 2003; 207:3-5.

Rollwitz J, Lupke M, Simkó M. Fifty-hertz magnetic fields induce free radical formation in mouse bone marrow-derived promonocytes and macrophages. *Biochim Biophys Acta* 2004; 1674:231-8.

Röschke J, Mann K. No short-term effects of digital mobile radio telephone on the awake human electroencephalogram. *Bioelectromagnetics* 1997; 18:172-6.

Rubin GJ, Das Munshi J, Wessely S. Electromagnetic hypersensitivity: a systematic review of provocation studies. *Psychosom Med* 2005; 67:224-32.

Salford LG, Brun A, Stureson K, Eberhardt JL, Persson BR. Permeability of the blood-brain barrier induced by 915 MHz electromagnetic radiation, continuous wave and modulated at 8, 16, 50, and 200 Hz. *Microsc Res Tech* 1994; 27:535-42.

Salford LG, Brun AE, Eberhardt JL, Malmgren L, Persson BR. Nerve cell damage in mammalian brain after exposure to micro waves from GSM mobile phones. *Environ Health Perspec* 2003; 111:881-3.

Possible effects of Electromagnetic Fields (EMF) on Human Health

Scarfi MR, Sannino A, Perrotta A, Sarti M, Mesirca P, Bersani F. Evaluation of genotoxic effects in human fibroblasts after intermittent exposure to 50 Hz electromagnetic fields: a confirmatory study. *Radiat Res* 2005; 164:270-6.

Schoemaker MJ, Swerdlow AJ, Ahlbom A, Auvinen A, Blaasaas KG, Cardis E, et al. Mobile phone use and risk of acoustic neuroma: results of the Interphone case-control study in five North European countries. *Br J Cancer* 2005; 93:842-8.

Schüz J, Böhler E, Berg G, Schlehofer B, Hettinger I, Schlaefer K, et al. Cellular phones, cordless phones, and the risk of glioma and meningioma (Interphone study group, Germany). *Am J Epidemiol* 2006a; 163:512-20.

Schüz J, Jacobsen R, Olsen JH, Boice JD Jr, McLaughlin JK, Johansen C. Cellular telephone use and cancer risk: update of a nationwide Danish cohort. *J Natl Cancer Inst* 2006b; 98:1707-13.

Seitz H, Stinner D, Eikmann Th, Herr C, Röösl M. Electromagnetic hypersensitivity (EHS) and subjective health complaints associated with electromagnetic fields of mobile phone communication – a literature review published between 2000 and 2004. *Sci Total Envir* 2005; 349:45-55.

Shirai T, Kawabe M, Ichihara T, Fujiwara O, Taki M, Watanabe S, et al. Chronic exposure to a 1.439 GHz electromagnetic field used for cellular phones does not promote N-ethylnitrosourea induced central nervous system tumors in F344 rats. *Bioelectromagnetics* 2005; 26:59-68.

Sienkiewicz Z, Jones N, Bottomley A. Neurobehavioural effects of electromagnetic fields. *Bioelectromagnetics* 2005; Suppl 7:S116-26.

Sies H, Cadenas E. Oxidative stress: damage to intact cells and organs. *Philos Trans R Soc Lond B Biol Sci* 1985; 311:617-31.

Simkó M, Richard D, Kriehuber R, Weiss DG. Micronucleus induction in Syrian hamster embryo cells following exposure to 50 Hz magnetic fields, benzo(a)pyrene, and TPA in vitro. *Mutat Res* 2001; 495:43-50.

Simkó M, Mattsson MO. Extremely low frequency electromagnetic fields as effectors of cellular responses in vitro: Possible immune cell activation. *J Cell Biochem* 2004; 93:83-92.

Simkó M, Hartwig C, Lantow M, Lupke M, Mattsson MO, Rahman Q, et al. Hsp 70 expression and free radical release after exposure to non-thermal radio-frequency electromagnetic fields and ultrafine particles in human Mono Mac 6 cells. *Toxicol Lett* 2006; 161:73-82.

Sommer AM, Lerchl A. The risk of lymphoma in AKR/J mice does not rise with chronic exposure to 50 Hz magnetic fields (1 microT and 100 microT). *Radiat Res* 2004a; 162:194-200.

Sommer AM, Streckert J, Bitz AK, Hansen VW, Lerchl A. No effects of GSM-modulated 900 MHz electromagnetic fields on survival rate and spontaneous development of lymphoma in female AKR/J mice. *BMC Cancer* 2004b; 4:77.

Speit G, Schutz P, Hoffmann H. Genotoxic effects of exposure to radiofrequency electromagnetic fields (RF-EMF) in cultured mammalian cells are not independently reproducible. *Mutat Res* 2007; 10; 626:42-7.

Possible effects of Electromagnetic Fields (EMF) on Human Health

SSI's Independent Expert Group on Electromagnetic Fields. Recent Research on EMF and Health Risks. Third annual report from SSI's Independent Expert Group on Electromagnetic Fields. Stockholm: SSI; 2005. http://www.ssi.se/PdfUpload/SSI_EMF_2005.pdf (accessed 24 April 2007).

Tafforeau M, Verdus MC, Norris V, White GJ, Cole M, Demarty M, et al. Plant sensitivity to low intensity 105GHz electromagnetic radiation. *Bioelectromagnetics* 2004; 25:403-7.

Takebayashi T, Akiba S, Kikuchi Y, Taki M, Wake K, Watanabe S, et al. Mobile phone use and acoustic neuroma risk in Japan. *Occup Environ Med* 2006; 63:802-7.

Thun-Battersby S, Mevissen M, Löscher W. Exposure of Sprague-Dawley rats to a 50-Hertz, 100-microTesla magnetic field for 27 weeks facilitates mammary tumorigenesis in the 7,12-dimethylbenz[a]-anthracene model of breast cancer. *Cancer Res* 1999; 59:3627-33.

Tsurita G, Nagawa H, Ueno S, Watanabe S, Taki M. Biological and morphological effects on the brain after exposure of rats to a 1439 MHz TDMA field. *Bioelectromagnetics* 2000; 21:364-71.

Tuschl H, Novak W, Molla-Djafari H. In vitro effects of GSM modulated radiofrequency fields on human immune cells. *Bioelectromagnetics* 2005; 27:188-96.

Uloziene I, Uloza V, Gradauskiene E, Saferis V. Assessment of potential effects of the electromagnetic fields of mobile phones on hearing. *BMC Public Health* 2005; 5:39.

UNEP/WHO (United Nations Environmental Programme/World Health Organization). Static Fields. Environmental Health Criteria 232. Geneva: World Health Organization; 2006.

Utteridge TD, GebSKI V, Finnie JW, Vernon-Roberts B, Kuchel TR. Long-term exposure of E-mu-Pim1 transgenic mice to 898.4 MHz microwaves does not increase lymphoma incidence. *Radiat Res* 2002; 158:357-64.

Verschaeve L, Maes A. Genetic, carcinogenic and teratogenic effects of radiofrequency fields. *Mutat Res* 1998; 410:141-65.

Vijayalaxmi, Bisht KS, Pickard WF, Meltz ML, Roti JL, Moros EG. Chromosome damage and micronucleus formation in human blood lymphocytes exposed in vitro to radiofrequency radiation at a cellular telephone frequency (847.74 MHz, CDMA). *Radiat Res* 2001; 156:430-2.

Vijayalaxmi, Obe G. Controversial cytogenetic observations in mammalian somatic cells exposed to radiofrequency radiation. *Radiat Res* 2004; 162:481-96.

Vijayalaxmi, Obe G. Controversial cytogenetic observations in mammalian somatic cells exposed to extremely low frequency electromagnetic radiation: a review and future research recommendations. *Bioelectromagnetics* 2005; 26:412-30.

Vrijheid M, Cardis E, Armstrong BK, Auvinen A, Berg G, Blaasaas KG, et al. Validation of Short-Term Recall of Mobile Phone Use for the Interphone Study. *Occup Environ Med* 2006; 63:237-43.

Wagner P, Roschke J, Mann K, Hiller W, Frank C. Human sleep under the influence of pulsed radiofrequency electromagnetic fields: A polysomnographic study using standardized conditions. *Bioelectromagnetics* 1998; 19:199-202.

Possible effects of Electromagnetic Fields (EMF) on Human Health

Wagner P, Roschke J, Mann K, Fell J, Hiller W, Frank C, et al. Human sleep EEG under the influence of pulsed radio frequency electromagnetic fields - Results from polysomnographies using submaximal high power flux densities. *Neuropsychobiology* 2000; 42:207-12.

Wahab MA, Podd JV, Rapley BI, Rowland RE. Elevated sister chromatid exchange frequencies in dividing human peripheral blood lymphocytes exposed to 50 Hz magnetic fields. *Bioelectromagnetics* 2006; 28:281-8

Wang B, Lai H. Acute exposure to pulsed 2450-MHz microwaves affects water-maze performance of rats. *Bioelectromagnetics* 2000; 21:52-6.

WHO (World Health Organization). Electromagnetic fields and public health. Electromagnetic Hypersensitivity. WHO Fact sheet N°296. Geneva: World Health Organization; 2005.

Wiat J, Hadjem A, Gadi N, Bloch I, Wong MF, Pradier A, et al. Modeling of RF head exposure in children. *Bioelectromagnetics* 2005; Suppl 7:S19-30.

Winker R, Ivancsits S, Pilger A, Adlkofer F, Rudiger HW. Chromosomal damage in human diploid fibroblasts by intermittent exposure to extremely low-frequency electromagnetic fields. *Mutat Res* 2005; 585:43-9.

Wolf FI, Torsello A, Tedesco B, Fasanella S, Boninsegna A, D'Ascenzo M, et al. 50-Hz extremely low frequency electromagnetic fields enhance cell proliferation and DNA damage: possible involvement of a redox mechanism. *Biochim Biophys Acta* 2005, 1743:120-9.

Yamaguchi H, Tsurita G, Ueno S, Watanabe S, Wake K, Taki M, et al. 1439 MHz pulsed TDMA fields affect performances of rats in a T-maze task only when body temperature is elevated. *Bioelectromagnetics* 2003; 24:223-30.

Yao Y, Li Y, Yang Y, Li C. Effect of seed pre-treatment by magnetic field on the sensitivity of cucumber (*cucumis sativus*) to ultraviolet B radiation. *Environ Exp Bot* 2005; 54:286-94.

Zaidi S, Khatoon S. Effect of electromagnetic fields (created by high tension lines) on the indigenous floral diversity in the vicinity of Karachi – I: Studies on PMC meiosis, meiotic products and pollen fertility. *Pakistan J Botany* 2003; 35:743-755.

Zeni O, Chiavoni AS, Sannino A, Antolini A, Forigo D, Bersani F, et al. Lack of genotoxic effects (micronucleus induction) in human lymphocytes exposed in vitro to 900 MHz electromagnetic fields. *Radiat Res* 2003; 160:152-8.

Zook BC, Simmens SJ. The effects of 860 MHz radiofrequency radiation on the induction or promotion of brain tumors and other neoplasms in rats. *Radiat Res* 2001; 155:572-83.

Zwamborn APM, Vossen SHJA, van Leersum BJAM, Ouwens MA, Mäkel WN. Effects of global communication system radio-frequency fields on well being and cognitive functions of human subjects with and without subjective complaints. The Hague: TNO Physics and Electronics Laboratory; 2003 (TNO-report FEL-03-C148). Available from URL: http://home.scarlet.be/milieugezondheid/dossiers/gsm/TNO_rapport_Nederland_sept_20_03.pdf . See also www.gr.nl/pdf.php?ID=1042 . (accessed 24 April 2007).

9. GLOSSARY

This section includes technical terms and definitions used within the document. The definitions are given in alphabetical order.

Conductivity: A property of materials that determines the magnitude of the electric current density when an electric field is impressed on the material.

Dielectric properties: In the context of this document the properties of materials conductivity and permeability.

Electric field strength (E): The magnitude of a field vector at a point that represents the force (F) on a charge (q). E is defined as $E = F/q$ and is expressed in units of Volt per meter (V/m).

Electromagnetic field: Electromagnetic phenomena expressed in vector functions of space and time.

Electromagnetic radiation: The propagation of energy in the form of electromagnetic waves through space.

EMF: Electromagnetic field.

Exposure: Exposure occurs wherever a person is subjected to electric, magnetic or electromagnetic fields or contact currents other than those originating from physiological processes in the body.

Extremely low frequency (ELF): Extremely low frequency fields include, in this document, electromagnetic fields from 1 to 300 Hz.

Frequency modulation (FM): Frequency Modulation is a type of modulation representing information as variations in the frequency of a carrier wave. FM is often used at VHF frequencies (30 to 300 MHz) for broadcasting music and speech.

Frequency (Hz): The number of cycles of a repetitive waveform per second.

Intermediate frequencies (IF): Intermediate frequencies are, in the frame of this report, defined as frequencies between 300 Hz and 100 kHz.

Magnetic flux density (B): The magnitude of a field vector at a point that results in a force (F) on a charge (q) moving with the velocity (v). The force F is defined by $F = q*(v \times B)$ and is expressed in units of Tesla (T).

Magnetic field strength (H): The magnitude of a field vector that is equal to the magnetic flux density (B) divided by the permeability (μ) of the medium. H is defined as $H = B/\mu$ and is expressed in units of Ampere per metre (A/m).

Microwaves: Microwaves are defined in the frame of this expertise as electromagnetic waves with wavelengths of approximately 30 cm (1 GHz) to 1 mm (300 GHz).

Milliwatt (mW): A unit of power equal to 10^{-3} Watt.

Nanowatt (nW): A unit of power equal to 10^{-9} Watt.

Non – thermal effects (or athermal effects): An effect which can only be explained in terms of mechanisms other than increased molecular motion (i.e. heating), or occurs

Possible effects of Electromagnetic Fields (EMF) on Human Health

at absorbed power levels so low, that a thermal mechanism seems unlikely, or displays so unexpected a dependence upon some experimental variable that it is difficult to see how heating could be the cause (see also Bernhardt et al. (1997)).

Permeability (μ): A property of materials that indicates how much polarisation occurs when an electric field is applied.

Power density (S): Power per unit area normal to the direction of propagation, usually expressed in watt per square meter (W/m^2).

Radio frequency (RF): The frequencies between 100 kHz and 300 GHz of the electromagnetic spectrum.

Specific absorption rate (SAR): A measure of the rate of power absorbed by or dissipated in an incremental mass contained in a volume element of dielectric materials such as biological tissues. SAR is usually expressed in terms of watts per kilogram (W/kg).

Static electric field: Static fields produced by fixed potential differences.

Static magnetic fields: Static fields established by permanent magnets and by steady currents.

VDU: Video display units for computers, videos, TV and some measurement devices using cathode ray tubes.