

EIP-AGRI Focus Group Reducing livestock emissions from Cattle farming

Mini-paper – Precision Livestock Farming

Authors

Thomas Bartzanas, Barbara Amon, Salva Calvet, Marcello Mele, Diego Morgavi, Tomas Norton, David Yanez-Ruiz, and Claude Vandongen

Introduction

While livestock production forms one of the pillars of the EU food industry it faces many societal challenges, not least from the rising demand for meat protein, increasingly stringent environmental regulations, coupled with the falling numbers of young farmers entering the industry. Environmental degradation as a result of the releases of harmful substances in the atmosphere can ultimately lead to regional tensions and violence, affecting economic policies and farmers' income.

It is estimated that 7.1 Gigatonnes of CO₂-equiv are emitted yearly in the livestock sector, representing 14.5 percent of all anthropogenic GHG emissions. Cattle (raised for both beef and milk) are the animal species responsible for the most emissions, representing about 65% of the livestock sector's emissions (FAO, 2013). In terms of activities, feed production and processing (including land use change) and enteric fermentation from ruminants are the two main sources of emissions, representing 45 and 39 percent of total emissions, respectively. Manure storage and processing represent 10 percent of total GHG emissions (Gerber, 2013). The remainder is attributable to the processing and transportation of animal products. Emission intensities currently vary widely within and across geographic regions and production systems, by a factor of between two and four, especially for products from ruminant animals (meat and milk) but also for pork and poultry. Intensive animal production systems tend to have higher overall GHG emissions, but their emissions intensity is lower than in low-yield extensive systems.

For all livestock production systems, opportunities exist and are being developed to improve the efficiency of production and decrease emissions per unit of animal product. Some of these options require novel technological interventions whereas others are 'simple' principles that can be applied already in most production systems. As farms continue to increase in size, taking advantage of economies of scale, farmers continuously seek new tools and technologies to reduce their operational cost and for increasing their sustainability. Livestock operations today are characterized by narrower profit margins than in the past, whereas the competition in the market has gradually increased. Consequently, small changes in production efficiency can have a major impact on profitability. In addition, given that today's livestock production is influenced by the consumers demands for high quality and safer products, there is pressure for a more environmental friendly production, zero zoonotic disease transmission, and improved animal welfare. All these reflect a continuing change in the way in which livestock operations are managed and lead to new production model which has to be based around technology and innovation, exploiting the principles of Precision Livestock Farming (PLF) (Berckmans, 2004). In general terms,

precision farming has been previously considered in a specific EIP Agri focus group with the following aim: to organise data capture and processing to mainstream the application of Precision Farming for input and yield optimisation, while trying to identify the main reasons behind the current lack of adoption, and identifying the key barriers to the implementation of Precision Farming on European farms (EIP-Agri, 2015).

PLF technologies enable continuous, automatic monitoring of animal welfare, health, production and environmental impact in real-time (Berckmans, 2007). PLF technologies can help farmers to increase livestock production and quality of production in a sustainable manner. Although nowadays, there is an abundance of information and technologies available to livestock farmers and managers, it is not generally structured in a way that can be readily applied. One reason could be that PLF research has been primarily carried out using traditional scientific approach of dividing on-farm processes into discrete and manageable pieces to determine solutions and provide important process improvements. Furthermore, many producers perceive that adopting high productive management systems involves increased financial risk. Currently, PLF techniques are predominantly applied in monitoring the health and welfare of pigs, poultry and dairy cows. Even though these technologies have shown great promise in the areas of health, reproduction and quality control, it is the precision feeding seems to be the most promising for reducing emissions of GHG and ammonia (Gerber et al. 2013).

Precision livestock technologies

Precision Farming (PF) in Europe uses new technologies in the handling and management of farm information, and in managing of spatial and temporal variability found on all crop producing farms. The premise of PF is that this better use of this information improves economic returns and reduces environmental impact. More information for the precision farming with focus on data management, use of precision farming technologies for input and yield optimisation and the main barriers for the implementation of Precision Farming on European farms can be found in the recent final report of the EIP-Agri focus group for precision farming which published in November 2015. Similar to Precision Farming, Precision Livestock Farming research and development has, over many years, adapted and developed information and communication technologies for livestock farming systems. Sensors, cameras, microphones, satellite positioning systems, automated equipment control and robots, wireless communication tools, Internet connections and cloud storage are at the technical core of an information-driven or 'information intensive' livestock production.

Precision Livestock Farming is defined as "the use of information and communication technologies for improved control of fine-scale animal and physical resource variability to optimize economic, social, and environmental dairy farm performance" (Eastwood et al., 2012). Another more or less similar definition coming from the most recent EU funded project for precision livestock farming, the EU-PLF project. According to this, Precision Livestock Farming (PLF) can be defined as "intelligent management and care of (individual) animals in livestock farming by continuous automated monitoring/controlling of the production/reproduction, health and welfare of (individual) animals, thereby allowing quick corrections when deviations from normal are monitored" (<http://www.eu-plf.eu/>). Precision livestock farming can offer a management tool that enables a farmer to monitor animals automatically and to create added value by helping to secure improved health, welfare, yields and environmental impact. Precision livestock farming is an approach that seeks to cater for the individual animal's needs in bigger herds, by integrating health, genetics, feed, social behaviour and resource use and availability, through the support of sensor-based monitoring systems. Precision livestock farming thus can readily be extended to individual animal level approaches towards optimising feed quality and digestibility, and animal health and husbandry (Wathes et al., 2008).




Fig. 1 presents the basis of PLF. Although the concept is for biological process (animal physiology, welfare, behaviour) the same concept is also applied to physical process (Aerts et al. 1998, 2003)




Fig. 1. Schematic overview of the key components of PLF to control biological processes (after Aerts et al. 1998, 2003)

There are several PLF technologies but still, very few of them used in practise. Although there are not developed for emission control or reduction, their use can also lead to a better emission control.

Reference	Application
Frost et al. 2000	Sensor placement robot for pigs
Gates et al. 2001	Intelligent ventilation control in livestock buildings
Matz et al. 2003	Dairy management to maximise profit
Stewart et al. 2007	Cattle monitoring system
Shao et al. 2008	Improved thermal control for pigs via machine vision
Banhazi et al. 2009	Improving profitability via precision feeding for pigs
Maertens et al. 2011	GaitWise: real time cow gait tracking
Van Hertem et al., 2016	Automated lameness detection in cows

http://agrifarm.dk/en/technologies/air-cleaning-in-the-dairy-barn	Indoor climate control
http://agrifarm.dk/en/the-first-hybridventilated-farm	Indoor climate control

Although there are several PLF technologies aimed at increasing the sustainability of livestock farms, only the precision feeding is indicated by the Global Research Alliance (2013) as the most promising PLF technology for reducing ammonia and GHG emissions from livestock farms. However, taking into account the rapid development of monitoring and climate control technologies we should probably see in the near future the use of intelligent PLF systems for emissions reduction through optimised barn climate control and better air flow through smart ventilation systems.

Precision feeding

Precision feeding is about getting the right nutrient to the right animal at the right time. Although direct mitigation effects are uncertain and hard to predict, precision feeding will increase feed efficiency and productivity and consequently can improve farm profitability. This can be done by controlling:

- individual feed intake
- the amount and composition of manure produced and the associated emissions from manure
- the enteric CH₄ production.

Customised balanced feeding programmes in grazing dairy cattle systems have shown to increase productivity and reduce enteric methane emissions intensity (15-20%) and also N excretion (20-30%), which results in reduced emissions from manure.

According to Knapp et al. (2014), in dairy cattle the mitigation of methane emission by controlling nutrition approach may be obtained i) altering the rumen VFA production pattern; ii) altering rumen microbe populations and increasing the rumen passage rate of digesta; iii) optimising feeding in order to increase milk yield (dilution effect).

Feed efficiency explains large part of the variability in GHG emission intensity (Thoma et al., 2010). Moreover, feed efficiency was identified as one of the most important parameters controlling GHG-emissions intensity, acidification and eutrophication potential in both beef and dairy cattle (Cederberg et al., 2009; Guerci et al., 2013).

Monitoring feed intake is one of the main constraints both for precision feeding and for estimate methane emission. Dry matter intake, in fact, is the main factors affecting methane emission (Hristov et al., 2013). Some technologies are available for monitoring feed intake on free-stall cattle (Chizzotti et al., 2015; Bikker et al., 2014; Mendes et al., 2011), whereas more efforts should be devoted for obtaining similar results also on grazing cattle. Moreover, feeding cows according to individual needs improves production and nutrient utilization compared with feeding a single diet to all cows (Maltz et al., 2013). However, in order to accomplish this aim, individual data about milk yield and composition and body weight are needed. On-line milk composition analyser that measures milk fat, protein, and lactose in real time at each milking with concurrent measurements of BW are now available and might generate useful data to refine the feeding program of dairy cows (Maltz et al., 2009). A very recent study demonstrated that when both energy and protein efficiency were improved, land use, water use, and GHG emission reductions ranged from 23.4 to 35.5% (White, 2016).

Diet manipulation is proposed as one of the techniques for reducing ammonia from agriculture in the UNECE Guidance Documents (UNECE, 2014) but also in a recent review from FAO dealing with techniques for reducing non-CO₂ GHG emission from livestock facilities (Hristov et al. 2013)

Precision feeding, which combines genetics of the animal with feeding and grazing management, requires advanced technological facilities to precisely monitor the animal's needs and manage pastures and forage production appropriately, and can be rolled out in high-value farm systems that use highly technological farming systems. In pig production, the inclusion of dietary fibre reduces NH₃ emission by shifting the nitrogen from urine to faeces due to promotion of bacterial growth in the large intestine (Jarret et al., 2012). A significant reduction in NH₃ emission can also be achieved by lowering the dietary electrolyte balance or supplementing with acidifying salts such as benzoic acid or CaSO₄ (Philippe et al., 2011). For dairy cattle, decreasing N excretion can be achieved by an improvement of rumen protein metabolism leading to lower NH₃ and N₂O production. Two methods are recommended: firstly, decreasing crude protein content in the diet to 14% (from 17-18%) which is often practiced, so that less N is excreted whilst both maintaining milk production, and secondly, limiting the digestion of ingested food meal in the rumen. According to Lee et al. (2012), feeding cows to amino acid requirements may be a promising way decreasing N excretion. However, more efforts must be devoted to obtain optimal prediction of amino-acid requirements in lactating dairy cows. As a consequence, improved understanding of amino-acid utilization in dairy cattle needs more research effort (Arriola Apelo et al., 2014).

Feeding strategies aimed to reduce methane emission include: i) increasing dry matter intake; ii) adoption of grain processing techniques; iii) increasing concentrate/forage ratio; iv) increasing forage quality; v) lipid supplementation; vi) lowering rumen pH below 5.5.

The effects of these strategies have been recently reviewed by Knapp et al (2014). In terms of methane reduction, lowering rumen pH and increasing concentrate amount in the diet resulted in the highest reduction (up to 15-20%). However, these strategies may lead also to milk yield reduction, negative effects on animal welfare and adverse consequences on GHG emission at global level (increasing demand of grains for concentrates). The other strategies allow to obtain lower percentages of methane mitigation (from 2 to 6%), without worsening animal performances.

According to Pomar et al (2013) the essential elements for precision feeding in livestock production systems are

- the proper evaluation of the nutritional potential of feed ingredients,
- the precise determination of nutrient requirements,
- the formulation of balanced diets that limit the amount of excess nutrients, and
- the concomitant adjustment of the dietary supply and concentration of nutrients to match the evaluated requirements of each pig in the herd.

A recent report from Global Research Alliance on agricultural greenhouse gases (2013) indicates that precision feeding can have a direct effect on resource efficiency and emissions reduction only if it is supported well by training, education and knowledge transfer. The same report is also highlight the need for creating customised feeding programmes and feed supply chains, which main in turn rely on stable market conditions and commodity prices.

The Fig. 2 below summarises the main drivers for success, barriers for implementation and mitigation potential of precision feeding systems (Global Research Alliance, 2013; Pomar et al. 2013)




Fig. 2. Barriers for implementation and mitigation potential of precision feeding systems (Global Research Alliance, 2013; Pomar et al. 2013)

Farmers adoption

Although most farmers can see the benefits of using a more precise approach to manage their livestock production with additional information, the tools provided by precision farming and other information technologies have not yet moved into mainstream livestock management. The increased complexity of the systems inhibits easy adoption and makes calculations as to the financial benefits uncertain.

As indicated by many PLF experts, the main problem of commercialisation and fast implementation of PLF systems in livestock production chain is the lack of support mechanisms, knowledge transfer and a consistent service offering for farmers. Experts highlight the need for a service sector that will be able to (Banhazi et al. 2012).

- take care of technology components,
- interpret data captured by sensors,
- formulate and send simple, relevant advice to farmers on a regular basis,
- involve users in technology developments.

This service sector would need to use suitable business models that avoid high initial investment costs for farmers.


The lack of co-ordination between researchers, developers, market and farmers is the other important limiting factor of adoption rate of PLF technologies. Achieving better co-ordination between the developers and suppliers of PLF tools is very difficult, but would result in the development of better integrated systems. That in turn would result in greater commercialisation of PLF systems as integrated systems to serve the farmers better (Banhazi et al. 2012).

The decision to purchase and implement a precision dairy technology represents a significant investment for a producer, who often faces the challenge of choosing a technology that will serve their needs for several years. In a recent study among other aspects trying to identify the parameters currently measured on farms, find the considerations a farmer takes when selecting precision dairy farming technologies, and determine the parameters perceived by producers as most useful (Borchers, 2015). Producers were able to select multiple parameters because several technologies can monitor multiple parameters. Additionally, the potential exists for producers to have more than one technology. Producer responses indicated that the most commonly measured parameters by already adopted technologies were: daily milk yield (52.3%), cow activity (41.3%), not applicable (31.2%), and

mastitis (25.7%). The least used technologies were rumen pH (0.9%), respiration rate (1.8%), methane emissions (1.8%), body condition score (2.8%), and heart rate (3.7%) (Borchers, 2015).

In the same study (Borchers, 2015) an attempt was made to quantify the criteria that the farmers considering more important for acquiring a PLF technology. The farmers indicated benefit-to-cost ratio as most important, followed by total investment cost, simplicity and ease of use, proven performance through independent research, and availability of local support.

According to Banhazi et al (2012), in order to increase the adoption level of these new PLF technologies in livestock sector it is essential to develop a management system that ensures only the most essential procedures are carried out, they are all carried out correctly and consistently, and in a way that controls risk. The main principles of such system are (Banhazi et al. 2012):



Contribution of PLF to resource efficiency

A better use of resources contributes to lower the emission intensity. In this sense, Precision Livestock Farming has demonstrated to contribute to farm efficiency, particularly in dairy cattle. For beef cattle, however, PLF has experimented less development. The use of technology now focuses on the management of grazing. As reviewed by Ali et al. (2016), satellite remote sensing is a promising tool to properly manage grasslands in terms of animal production and ecosystem protection. For animal management, attention is now on telemetry technologies, particularly the use of GPS. This allows an automatic record of behaviour (particularly foraging), which combined with pasture characteristics may allow more accurate determination of animal intake and provide tools for decision making (Orr et al., 2012; Thompson et al., 2015). Finally, GPS location has been also used to estimate emissions from grazing animals by means of inverse modelling (McGinn et al., 2015).

Concerning emissions reduction precision feeding seems to have a direct impact on emission reduction. However, it is expected soon that emission reduction could be achieved also through smart ventilation systems and optimised barn climate.

Proposal for potential operational groups

- Precision Livestock Farming Technologies
- Integration of PLF with breeding and nutrition initiatives

- Demonstrating the viability of PLF in small and medium livestock farms
- Integrating PLF, the Internet of Things and Big Data

Proposals for (research) needs from practice

- A reliable and low cost system for measuring ventilation rate in livestock buildings
- Low cost sensors for detection of emission and concentration of ammonia and GHG
- Establish a common protocol for measuring ammonia and GHG in livestock operations
- Develop self-learning systems that optimise barn climate in naturally ventilated barns by e.g. altering barn openings => avoid high temperatures in the barn
- Hybrid ventilation system
- Targeted ventilation in animal occupied zones
- Sensors for in vivo detecting rumen environmental parameters (methane; H₂; CO₂; VFA)
- Sensors for monitoring animal behaviour at pasture (position, feeding activity, rumination)

☞ To be put in the [web form](#)

References

- Ali I., Cawkwell F., Dwyer E., Barrett B., Green S., 2016. Satellite remote sensing of grasslands: from observation to management—a review. *Journal of Plant Ecology*, doi:10.1093/jpe/rtw005
- Aerts J-M., Berckmans D., Decuyper E., Buyse J., Goedseels V. 1998. Modelling growth responses of broiler chickens to variations of the control input feed supply. *Proceedings of the 1998 ASAE Annual International Meeting, Orlando, USA, July 12–15, 15 pp. (Paper no. 98-4045).*
- Aerts J.-M., Lippens M., De Groote G., Buyse J., Decuyper E., Vranken E., Berckmans D. 2003. Recursive prediction of broiler growth response to feed intake by using a time-variant parameter estimation method. *Poult. Sci.* 82, 40–49.
- Arriola Apelo S., Knapp J., Hanigan M. 2014. Invited review: current representation and future trends of predicting amino acid utilization in the lactating dairy cow. *J. Dairy Sci.*
- Banhazi T.M., Rutley D., Parkin B., Lewis B. M. 2009. Field evaluation of a prototype sensor for measuring feed disappearance in livestock buildings. *Australian Journal of Multi-disciplinary Engineering*, 7(1): 27-38
- Banhazi T., Lehr M., Black H., Crabtree J.L., Schofield H., Tscharke P., Berckmans D. 2012. Precision Livestock Farming: an international review of scientific and commercial aspects. *International Journal of Agriculture and Biological Engineering*, 5 (3): 1-10
- Berckmans D. 2004. Automatic On-Line Monitoring of Animals by Precision Livestock, *International Society for Animal Hygiène- Saint-Malo*
- Berckmans D. 2007. Precision livestock farming, *Computers and Electronics in Agriculture*, 62 (1)
- Bikker J.P., van Laar H., Rump P., Doorenbos J., van Meurs K., Griffioen G.M., Dijkstra J. 2014. Technical note: Evaluation of an ear-attached movement sensor to record cow feeding behavior and activity. *Journal of Dairy Science*, 97: 2974-2979.
- Borchers M.R. 2015 "An evaluation of precision dairy farming technology adoption, perception, effectiveness, and use". *Theses and Dissertations--Animal and Food Sciences*. Paper 45.
- Cederberg C., Meyer D., Flysjö A., 2009. Life cycle inventory of greenhouse gas emissions and use of land and energy in Brazilian beef production. *SIK-Institutet för livsmedel och bioteknik*.
- Chizzotti M.L., Machado F.S., Valente E.E.L., Pereira L.G.R., Campos M.M., Tomich T.R., Coelho S.G., Ribas M.N. 2015. Technical note: Validation of a system for monitoring individual feeding behavior and individual feed intake in dairy cattle. *Journal of Dairy Science*, 98: 3438-3442.

EIP-AGRI Service Point 2015. Focus Group "Precision Farming", Final Report, 44 pp.

Eastwood C. R., Chapman, D. F., & Paine, M. S. 2012. Networks of practice for co-construction of agricultural decision support systems: Case studies of precision dairy farms in Australia. *Agricultural Systems*, 108, 10-18.

FAO 2013: Mitigation of greenhouse gas emissions in livestock production: A review of technical options for non-CO₂ emissions. FAO Animal Production and Health Paper No. 177. Gerber, P.J. et al. (eds), Food and Agriculture Organisation of the United Nations (FAO), Rome.

Frost A. R., Tillett R. D., Welch S. K. 2000. The development and evaluation of image analysis procedures for guiding a livestock monitoring sensor robot. *Computers and Electronics in Agriculture*, 28(3): 229-242

Gates R. S., Chao K., Sigrimis N. 2001. Identifying design parameters for fuzzy control of staged ventilation control systems. *Computers and Electronics in Agriculture*, 2001;31(1): 61-74.

Gerber P.J. et al. 2013: Tackling Climate Change Through Livestock: a global assessment of emissions and mitigation opportunities. Food and Agriculture Organisation of the United Nations (FAO), Rome

Gerber P.J., Hristov A.N., Henderson B., Makkar H., Oh J., Lee C., Meinen R., Montes F., Ott T., Firkins J. and Rotz A., 2013. Technical options for the mitigation of direct methane and nitrous oxide emissions from livestock: a review. *Animal*, 7: 220-234.

Global Research Alliance on agricultural greenhouse gases 2013. Reducing greenhouse gas emissions from livestock: Best practice and emerging options, p.46

Guerci M., Knudsen M.T., Bava L., Zucali M., Schönbach P., Kristensen T., 2013. Parameters affecting the environmental impact of a range of dairy farming systems in Denmark, Germany and Italy. *J. Clean. Prod.* 54, 133–141.

Hristov A.N., Oh J., Lee C., Meinen R., Montes F., Ott T., Firkins J., Rotz A., Dell C., Adesogan A., Yang W., Tricarico J., Kebreab E., Waghorn G., Dijkstra J. & Oosting S. 2013. Mitigation of greenhouse gas emissions in livestock production – A review of technical options for non-CO₂ emissions. Edited by Pierre J. Gerber, Benjamin Henderson and Harinder P.S. Makkar. FAO Animal Production and Health Paper No. 177. FAO, Rome, Italy.

Jarret G., Cerisuelo A., Peu P., Martinez J., Dourmad J.Y., 2012. Impact of pig diets with different fibre contents on the composition of excreta and their gaseous emissions and anaerobic digestion. *Agric. Ecosyst. Environ.* 160, 51-58.

Knapp J.R., Laur G.L., Vadas P.A., Weiss W.P., Tricarico J.M. 2014. Invited review: Enteric methane in dairy cattle production: quantifying the opportunities and impact of reducing emissions. *J. Dairy Sci.* 97:3231–3261.

Lee C., Hristov A., Heyler K., Cassidy T., Lapierre H., Varga G., Parys C. 2012. Effects of metabolizable protein supply and amino acid supplementation on nitrogen utilization, milk production, and ammonia emissions from manure in dairy cows. *J. Dairy Sci.* 95, 5253–5268.

Maltz E., Livshin N., Antler A., Edan Y., Matza S., Antman A. 2003. Variable milking frequency in large dairies: performance and economic analysis - models and experiments. in 1st European Precision Livestock Farming. Berlin, Germany: Wageningen Academic Publisher

Maertens W., Vangeyte J., Baert J., Jantuan A., Mertens K C., De Campeneere S., Pluk A., Opsomer O., Van Weyenberg S., Van Nuffel A., 2011. Development of a real time cow gait tracking and analysing tool to assess lameness using a pressure sensitive walkway: The GAITWISE system. *Biosystems Engineering*, 110 (1): 740-748

Maltz E., Antler A., Halachmi I., Schmilovitch Z. 2009. Precision concentrate rationing to the dairy cow using on-line daily milk composition sensor, milk yield and body weight. Proc. 4th ECPLF, Wageningen, the Netherlands. Wageningen Academic Publishers, Wageningen, the Netherlands.

Maltz E., Barbosa L.F., Bueno P., Scagion L., Kaniyamattam K., Greco L.F., De Vries A., Santos J.E.P. 2013. Effect of feeding according to energy balance on performance, nutrient excretion, and feeding behavior of early lactation dairy cows. *Journal of Dairy Science*, 96: 5249-5266.

- Mendes E.D.M., Carstens G.E., Tedeschi L.O., Pinchak W.E., Friend T.H. 2011. Validation of a system for monitoring feeding behavior in beef cattle. *Journal of Animal Science*, 89: 2904-2910.
- McGinn S.M., Flesch T.K., Coates T.W., Charmley E., Chen D., Bai M., Bishop-Hurley G. 2015. Evaluating dispersion modelling options to estimate methane emissions from grazing beef cattle. *Journal of Environmental Quality* 44, 97-102.
- Mertens K. 2009. An intelligent system for optimizing the production and quality of consumption eggs based on synergistic control. PhD thesis, Katholieke Universiteit Leuven, Belgium.
- Orr R.J., Tozer K.N., Griffith B.A., Champion R.A., Cook J.E., Rutter S.M. 2012. Foraging paths through vegetation patches for beef cattle in semi-natural pastures. *Applied Animal Behaviour Science* 141, 1-8
- Philippe F.X., Laitat M., Wavreille J., Bartiaux-Thill N., Nicks B., Cabaraux J.F. 2011. Ammonia and greenhouse gas emission from group-housed gestating sows depends on floor type, *Agriculture, Ecosystems & Environment*, 140 (3-4): 498-505
- Pomar C., Hauschild L., Zhang G H., Pomar J., Lovatto P A., 2001. Applying precision feeding techniques in growing-finishing pig operations. *Revista Brasileira de Zootecnia*, 38:226-237
- Shao B, Xin H. A real-time computer vision assessment and control of thermal comfort for group-housed pigs. *Computers and Electronics in Agriculture*, 2008; 62(1): 15-21
- Sloth K.H. Frederiksen D., Klimpel S., 2015. Estimated cow time-budgets using a Real Time Location System. *Precision Livestock Farming 2015 - Papers Presented at the 7th European Conference on Precision Livestock Farming, ECPLF 2015*
- Stewart M., Webster J. R., Verkerk G. A., Schaefer A. L., Colyn J. J. 2007. Non-invasive measurement of stress in dairy cows using infrared thermography. *Physiology & Behavior*, 92(3): 520-525.
- Thoma G. Popp J., Nutter D., Shonnard D., Ulrich R., Matlock M., Kim D.S., Neiderman Z., Kemper N., East C., Adom F. 2013. Greenhouse gas emissions from milk production and consumption in the United States: A cradle-to-grave life cycle assessment circa 2008. *Int. Dairy J.* 31(Suppl.1):S3-S14.
- Thompson D.J., Wheatley B.J., Church J.S., Newman R., Walker J. 2015. Comparing grazing and resting electivity of beef cattle for BC bunchgrass communities using GPS collars. *Canadian Journal of Animal Science* 95, 499-507.
- Van Hertem T., Bahr C., Tello A Schlageter., Viazzi S., Steensels M., Romanini C.E.B., Lokhorst C., Maltz E., Halachmi I., Berckmans D. 2016. Lameness detection in dairy cattle: single predictor v. multivariate analysis of image-based posture processing and behaviour and performance sensing, *Animal* 10 (9): 1525-1532
- UNECE 2014. Guidance Document on Preventing and Abating Ammonia Emissions from Agricultural. ECE/EB.AIR/120. 7 February 2014. UNECE, Geneva, p. 100.
- Wathes C.M., Kristensen H.H., Aerts J.M., Berckmans D. 2008. Is precision livestock farming an engineer's daydream or nightmare, an animal's friend or foe, and a farmer's panacea or pitfall? *Computers and Electronics in Agriculture* 64, 2-10.
- White R.R. 2016. Increasing energy and protein use efficiency improves opportunities to decrease land use, water use, and greenhouse gas emissions from dairy production. *Agricultural Systems*. 146: 20-29