EIP-AGRI Workshop 'Biosecurity at farm level: challenges for innovation' 22-23 January 2015 – Brussels, Belgium








Biosecurity in animal farming

Prof. Dr. Jeroen Dewulf

Jeroen.Dewulf@UGent.be

In collaboration with Maria Laanen, Merel Postma, Philippe Gelaude and Steven Sarrazin

What is biosecurity? All measures taken to prevent:



- 1. disease introduction
 - 6%
- 2. disease spread 3%
- 3. disease introduction and disease spread

91%



What is biosecurity



BIOSECURITY

The combination of all measures taken to reduce the risk of introduction and spread of diseases on herd, region, country,... level

'assessing risk and implementing measures to decrease that risk and to safeguard and improve health status on a farm'


What is biosecurity

EXTERNAL BIOSECURITY

- = Reduce introduction
 - endemic diseases
 - "exotic" diseases




= reduce spread





BIOSECURITY is (should be) the basis of any disease control program



- Better biosecurity ____ less disease
 - Better production results
 - reproduction
 - growth
 - feed conversion
 - uniformity
 - Less antimicrobial use
 - Higher prices when selling the animals



- Beter biosecurity less disease
 - Eradication programs ↑
 - Free / Obliged (Salmonella Action Plan)
 - Risk of exotic diseases
 - Public health, animal welfare, public opinion ('sustainable meat production')
 - Legislation


- Farmers motivators to take disease prevention measures (=558)
 - 1. To improve profit due to higher productivity
 - 2. To have farm stability
 - To improve quality and safety of produced products
 - 4. To improve animal welfare
 - 5. To fulfill the duty of keeping my animals healthy

Separation of infected and diseased animals


→ avoid both direct and indirect contact! (all-in/all-out, working lines, hospital pen, ...)



Direct disease contact:



Indirect disease contact:




Reduction of the general infection pressure

→ breaking the infection cyclus, reducing the burden on the immune system ↓ (cleaning, desinfection and empty period, vaccination, ...)











- Dependent upon herd situation (status, type,...)
- Perform well and consequent

- Based on knowledge of
 - disease transmission
 - survival of sources of infection
- General and species specific
- General and disease specific
 - Pulmonary diseases versus enteric diseases
 - Multifactorial diseases (PRRS & influenza, PMWS, PIA,...)

Amass S. (2005)	Pers.	Sperm	Faeces	And. / birdsl	Rodents	Insect.	<u>Aerosol</u>	Feed	<u>Water</u>	Fomites
M. hyopneumoniae							*		*	*
P. multocida	*		*				*		*	
Parvo		*	*							*
PRRSV	*	*	*	*		*	*		*	*
Aujeszky		*	*	*	*	*	*			*
Salmonella spp.	*		*	*	*	*	*	*	*	*
Strept. suis	*		*	*		*	*		*	*
Influenza	*		*	*			*			
Blaasjesziekte	*	*					*			*

Amass S. (2005)	Pers.	Sperm	Faeces	And. / vogel	Rodents	Insect.	<u>Aerosol</u>	<u>Feed</u>	Water	Fomites
APP							*			
B. bronchiseptica				*	*		*		*	
B. hyodysenteriae			*	*	*				*	
Brucella. suis	*	*		*						
Swine Fever	*	*	*	*		*	*			*
Clostr. perfringens							*		*	
E. coli	*		*		*	*	*		*	*
Food and mouth disease	*	*	*				*			*
Leptospirose	*	(*)		*	*				*	





Assume: risk of disease introduction on your herd through feed delivery is 1 out of 1000 and the feed delivery truck comes weekly, what is the annual risk?

28%


55%


17%



- 'Thousand times a small chance becomes a large chance'
 - Risk transmission route (p)
 - Frequency transmission route (n)
- $P = 1 (1-p)^n$
 - p= 0.1% (1 out of 1000)
 - n= 52 (e.g. weekly)
 - **5,06%**= $1 (1-0.001)^{52}$

Purchase of animals and sperm





Purchase of animals and sperm



Purchase of animals and sperm



pasture contact between animals



Disposal of animals and material





Disposal of animals and material







management and storage of carcasses



Supply of feed, water and goods
































Vermin and bird control








Vermin and bird control





External Biosecurity

location and surrounding









Disease management







Suckling pen

Suckling pen

1 week of age

Piglets after weaning


fatteners

All-in/ All-out



Separate age groups






Compartments and working lines








Pets





Materials







Needles and medicines





Cleaning and disinfection

1. Dry



3. Wet







Cleaning and disinfection

4. Dry



5. Disinfect




6. Empty



7. Check



Principles of biosecurity What is more important?



1. External biosecurity

24%

2. Internal biosecurity


7%

3. Equally important

69%




Positive correlation between scores for external and internal biosecurity



Biosecurity = complex

- No protocol suitable for every herd
- Balance biosecurity management
- Tool?

→ Scoring System





Scoring system and website Pigs and Poultry

Biocheck, prevention is better than cure!



www.biocheck.ugent.be

Biosecurity Scoring System

- Quantification of biosecurity status
 - Comparing of scores between different herds
 - Comparing of scores in time
 - Taking different risks into account



Scoring System

- External and internal biosecurity
- 6 subcategories per part
- 2-13 questions per subcategory
- Questionnaire in Dutch, English, French, German, Swedish,....





Scoring System

Weighted scores

Based on scientific research



- Risk for transmission: direct vs. indirect contact
- Weight factor for each subcategory and each question



Free online application: www.biocheck.ugent.be






Biocheck, prevention is better than cure!

Login Register

You are here: Home





Welcome

On this website, you can evaluate the quality of the biosecurity on your pig farm through a scientific based questionnaire and accompanying scoring system and this is completely **free** and **independent**.

Before you can fill in the survey, you have to log in first.

Furthermore, this website offers an extensive review of information concerning biosecurity and biosecurity measures on pig farms. You can find this information in the 'About Biocheck' section in the menu on the left.

Free literature on biosecurity is also available on this site. For an overview, click here!

Four times a year a "biocheck" newsletter is made containing each time an hot topic of biosecurity in pig production, explained in a very practical manner. If you want to receive this newsletter subscribe here.

This website and the accompanying scoring system have been developed by Prof. Jeroen Dewulf, Dr. Stefaan Ribbens and Drs. Maria Laanen with technical support of Steven Bruneel.

For remarks, questions, suggestions about this website or the scoring system, please mail to Steven.Bruneel@UGent.be.

Newsletter

- >> Subscribe
- >> Unsubscribe

Info

This questionnaire and scoring system is designed by the


Faculty of Veterinary Medicine



THE FACULTY HAS BEEN

in collaboration with Flemish Animal Health Organization









Biocheck report

Vakgroep Voortplanting, Verloskunde en Bedrijfsdiergeneeskunde

Your subtotal for external biosecurity is $\bf 59\%$ (avg. 65%) Your subtotal for internal biosecurity is $\bf 56\%$ (avg. 50%) The total score is $\bf 58\%$ (avg. 58%)

Nr	Description	Score	Avg
dernal bio	security		
Α	Purchase of animals and semen	62 %	87 %
В	Transport of animals, removal of manure and dead animals	61 %	66 %
C	Feed, water and equipment supply	53 %	39 %
D	Personnel and visitors	59 %	63 %
E	Vermin and bird control	30 %	60 %
F	Environment and region	80 %	54 %
S1888	77: X-17: X-		
0.000	Subtotal External biosecurity:	59%	65 %
ternal bios A	Subtotal External biosecurity:	59 % 40 %	65 % 56 %
ternal bios	Subtotal External biosecurity:		W-1000-01000
ternal bios A B	Subtotal External biosecurity: security Disease management	40 %	56 %
ternal bios A	Subtotal External biosecurity: security Disease management Farrowing and suckling period	4 <mark>0 %</mark> 64 %	56 % 62 %
ernal bios A B C	Subtotal External biosecurity: cecurity Disease management Farrowing and suckling period Nursery unit	40 % 64 % 43 %	56 % 62 % 57 %
ernal bios A B C D	Subtotal External biosecurity: Disease management Farrowing and suckling period Nursery unit Fattening unit	40 % 64 % 43 % 93 %	56 % 62 % 57 % 57 %
ernal bios A B C D E	Subtotal External biosecurity: Disease management Farrowing and suckling period Nursery unit Fattening unit Measures between compartments and the use of equipment	40 % 64 % 43 % 93 % 50 %	56 % 62 % 57 % 57 % 40 %







Biocheck report

Vakgroep Voortplanting, Verloskunde en Bedrijfsdiergeneeskunde

Bellow, a **figure** showing your results (red area) graphically compared to the average scores (green area), is given. The bigger the red area, the better the result. The names of the axes correspond to the numbering in report above.





Naming of the axes is linked to the numbering on the first page

For **more explanation** about biosecurity on pig herds and how to improve this, you can always look at the extensive information available on the **biocheck website** in the sections "about biocheck" and "download and links". You can also contact your **veterinarian** for advice on biosecurity measures on your farm.

www.biocheck.ugent.be




Biocheck, prevention is better than cure!



www.biocheck.ugent.be

Who should apply biosecurity Where is biosecurity most important?



1. large herds

21.1%

2. small herds

1.4%

3. independent of herd size

77.5%




Principles of biosecurity

• It's a numbers game







Biosecurity and herd and farmer characteristics




Laanen et al., 2013

Biosecurity and herd and farmer characteristics





Who is responsible for biosecurity Biosecurity is primarily a responsibility of...




1. farmer

92.5%

- 2. veterinarian
 - 3.0%
- 3. herd advisors
 - 1.5%
- 4. government
 - 3.0%



Biosecurity is a joint venture...



Biosecurity is a joint venture... ...often threatened by lack of time



