



Health effects attributable to exposure to ground-level ozone – 2000



Health-relevant ozone exposure expressed as SOMO35 (ppb.days), for the emissions of the year 2000. Calculation results for the meteorological conditions of 1997.

Source: IIASA


Health effects attributable to exposure to ground-level ozone – 2020



Health-relevant ozone exposure expressed as SOMO35 (ppb.days), for the emissions of the current legislation case of the “Climate policy” scenario in 2020. Calculation results for the meteorological conditions of 1997.


Source: IIASA

Health effects attributable to exposure to ground-level ozone – Scenario A



Health-relevant ozone exposure expressed as SOMO35 (ppb.days), for the emissions of the D23 (A) scenario in 2020. Calculation results for the meteorological conditions of 1997.
Source: IIASA


Health effects attributable to exposure to ground-level ozone – Scenario B



Health-relevant ozone exposure expressed as SOMO35 (ppb.days), for the emissions of the D23 (Scenario B) scenario in 2020. Calculation results for the meteorological conditions of 1997.


Source: IIASA

Health effects attributable to exposure to ground-level ozone – Scenario C



Health-relevant ozone exposure expressed as SOMO35 (ppb.days), for the emissions of the D23 (C) scenario in 2020. Calculation results for the meteorological conditions of 1997.
Source: IIASA

Health effects attributable to exposure to ground-level ozone – Thematic Strategy 2020



Health-relevant ozone exposure expressed as SOMO35 (ppb.days), for the emissions of the Thematic Strategy on Air Pollution in 2020. Calculation results for the meteorological conditions of 1997.

Source: IIASA