



EUROPEAN COMMISSION
HEALTH & CONSUMER PROTECTION DIRECTORATE-GENERAL

Directorate C - Scientific Opinions
C2 - Management of scientific committees II; scientific co-operation and networks

Scientific Committee on Food

SCF/CS/ADD/EDUL/215 Final
24 March 2003

Opinion of the Scientific Committee on Food

on

Erythritol

(opinion expressed on 5 March 2003)

Opinion on Erythritol

Terms of reference

To evaluate the safety of erythritol to be used as a food additive and to confirm its non-laxative effect as claimed by the petitioners. Additionally, to confirm the energy value of erythritol (0 to 0.2 kcal/g) as claimed by the petitioners.

Background

Erythritol is a four-carbon sugar alcohol (polyol) that has sweetness approximately 60-80% that of sucrose. It occurs naturally in minor amounts in some fruits (watermelon, pear and grape (Shindou et al., 1989). It also occurs in mushrooms, fermented foods (wine, sake, beer, soy sauce) (Shindou et al., 1988; Mitsubishi Chemical Corporation 1991) and cheese (Shindou and Sihizuka, 1996).

Erythritol has been used as a food ingredient in Japan since 1990 (Cerestar Holding B.V., Mitsubishi Chemical Corporation and Nikken Chemicals Co., Ltd, 1999). It has been approved in USA since 2001 (U.S. FDA, 2001).

Erythritol was evaluated by the Joint FAO/WHO Expert Committee on Food Additives (JECFA) on its fifty-third meeting and assigned an ADI "not specified" (JECFA, 1999a).

Technical data

The systematic name for erythritol is 1,2,3,4-Butanetetrol. It has the following chemical identity:

EINECS-number: 205-737-3

C.A.S.-number: 149-32-6

Chemical formula. $C_4H_{10}O_4$

Structural formula:

$$\begin{array}{c} \text{CH}_2\text{OH} \\ | \\ \text{H} - \text{C} - \text{OH} \\ | \\ \text{H} - \text{C} - \text{OH} \\ | \\ \text{CH}_2\text{OH} \end{array}$$

Formula weight: 122.12

Description: White, odourless, non-hygroscopic, heat-stable crystals. It has a sweetness approximately 60-80% that of sucrose.

Solubility: Freely soluble in water, slightly soluble in ethanol, insoluble in diethyl ether.

Melting range: Between 119° and 123°

Erythritol which is intended for use as a low calorie sweetener is produced by two different manufacturing processes, the Misubishi/Nikken method (M/N-method) and the Cerestar method (C-method) from wheat or corn starch by enzymatic hydrolysis yielding glucose. This is fermented by yeast-like fungi such as *Trichosporonoides megachiliensis* (M/N-method) or *Moniliella pollinis* (C-method). The fermentation broth is heated to kill the production organism and dead cells are removed by filtering. Once erythritol is separated from the fermentation broth, it is purified by ion exchange resin, activated charcoal, ultrafiltration and crystallisation. The final crystalline product is more than 99% pure (Cerestar Holding B.V., Mitsubishi Chemical Corporation and Nikken Chemicals Co., Ltd, 1999).

The purity criteria proposed by the applicants are in good agreement with analytical results obtained from different batches representing both manufacturing processes. It is noted that the proposed level of lead is twice as much as in the JECFA purity criteria (JECFA, 1999b).

Predicted exposure

The mean intake per person of erythritol estimated by the petitioners from natural occurrence in foods like wine, cheese, fruits, chocolate and mushrooms varies between countries, being e.g. 24 mg/day in USA and 105 mg/day in Japan. In comparison, the figure of 47 mg/day is a Danish estimate (DVFA, 2001).

The main food applications and maximum use levels of erythritol proposed by the petitioners include:

Food category	Maximum level of erythritol in product % by weight
Table top substitutes	99.9
Low-calorie beverages including sport drinks	3.5
Sugar-free chewing gum	60
Reduced calorie and sugar-free chocolate	50
Candies, not sugar-free: soft (fudges)	40
hard	50
Fondants and creams	60
Lozenges	99
Bakery (pastry) products (cookies biscuits, cakes pastries with cream filling/topping), not sugar-free	7-60

The calculations of estimated mean daily intake of erythritol using Danish disappearance and consumption data (DVFA, 2001) and the proposed main food applications and maximum use levels of erythritol in those are 18 to 27 g/day depending on sex and age, being the highest for 7 to 10 year old boys. When expressed as relative intake estimates these figures are for women 0.4 g/kg bw assuming an average body weight of 64 kg, for men 0.4 g/kg bw assuming an average body weight of 80 kg, for girls 1.3 g/kg bw with a reference body weight of 27.3 kg, and for boys 1.5 g/kg bw with a reference body weight of 27.8 kg. These are considered as “worst case “ calculations of the average consumption for adults and children as

they are based on the assumption that the erythritol intake derives from products where erythritol has replaced sugar.

In comparison, the U.S. Food and Drug Administration's own calculations of estimated daily intake of erythritol under conditions of use proposed by Cerestar, are 13 g/day at the mean and 30 g/day at the 90th percentile (U.S. FDA, 2001).

The petitioners provided lower estimates of intake based on several assumptions regarding the percentage of sugar-free products to be consumed. The estimates were 9.6 g/day and 11.2 g/day at the 90th percentile for the total population and for teenagers respectively.

Microbiological evaluation

The application from the petitioners involves the use of two fermentation processes for the production of erythritol each using a different fungus. *Trichosporonoides megachiliensis* Inglis & Sigler 1992, which is used in one of the processes was described from isolates associated with alfalfa leafcutter bees (Inglis & Sigler 1992) and the strain used in the process was isolated from soil in a sugar cane plant and selected following UV and NTG (nitrosomethyl guanidine) treatment. *Moniliella pollinis* (Hennebert & Verachtert) de Hoog & Gueho 1984, which is used in the other process, is the type strain of the organism and was isolated from pollen (Dooms *et al.* 1971; de Hoog & Gueho 1984).

Both *M.pollinis* and *T.megachiliensis* are osmophilic fungi and together with related species can contaminate high sugar foods such as syrups, jams and honey. Some *Moniliella* species are found in low pH foods (pickles, sauces) and both *Moniliella* spp. and *Trichosporonoides* spp. have been isolated from foods high in fat (e.g. margarine, ghee) (Samson & van Reenen-Hoekstra 1988).

Both fungi are able to grow at 35-37°C. A search of the literature, major fungus culture collection and taxonomic databases failed to reveal any documented evidence of *M.pollinis* or *T.megachiliensis* being pathogenic to humans. Related species have been isolated from clinical material although they appear to be rare occurrences (Kockova-Kratochvilova *et al.* 1987; McKenzie *et al.* 1984).

There is no indication in the literature that *T.megachiliensis* or *M.pollinis* produce toxic metabolites or antimicrobial compounds. However, both fungi have been reported to produce pigmented metabolites in culture (Dooms *et al.*, 1971; Inglis & Sigler 1992) although the petitioners have provided information indicating that colour changes in the fermentation broth are minor and can be explained by variation in broth components and the effect of heat treatment.

The fermentation broth produced by both fungi has been tested for toxicogenic activity. The isolate of *T.megachiliensis* was tested by feeding the fermentation broth to rats in doses up to 2000 mg dry weight per kg body weight with no indications of toxicity (Kashima Laboratory Mitsubishi Chemical Safety Institute Ltd., 1995). A repeated-dose (4-week) oral toxicity study of *M.pollinis* fermentation broth in rats demonstrated that dietary levels up to 2.5% (the highest dose tested) corresponding to an overall intake of 2.1 g/kg bw/day did not induce any treatment related changes (Lina, 2002).

According to the petitioners, use of the erythritol product will not expose consumers to the producing organisms, which will be destroyed by heat treatment of the fermentation broth and removed by filtration and purification. The petitioners provided end product microbiological specifications for their products together with some information indicating consistency between batches. There should be no microorganisms of public health significance if good manufacturing/hygienic practice and Hazard Analysis Critical Control Point principles are applied throughout the manufacturing processes.

No information was found in the literature concerning microbiological contamination problems associated with erythritol and the water content of the erythritol product (<0.2%) is too low to support microbial growth.

Absorption, distribution, excretion and biotransformation

In animals and humans, depending on doses, 60 to more than 90 % of ingested erythritol is rapidly absorbed from the small intestine and excreted unchanged in the urine (Bornet et al. 1996a, b; Dean et al., 1996; Hiele et al., 1993; Ishikawa et al., 1996; Lina et al., 1996; Nakayama, 1990a, b; Noda, 1994; Noda and Oku, 1990, 1992; Noda et al., 1988; Noda et al., 1996; Oku and Noda 1992a, b; Tetzlof et al., 1996; Til et al., 1996; van Ommen et al., 1990, 1996).

Consumption of erythritol with foods appears to delay absorption (Bornet et al. 1996b). Absorbed erythritol is rapidly distributed throughout the body in animals and humans, with peak plasma, serum and/or blood concentrations generally occurring within 1 hour of ingestion (Bornet et al., 1996a, b; Ishikawa et al., 1996; Nakayama, 1990a, b, c, d; Noda et al., 1994, 1996). In humans, plasma erythritol levels have been reported to reach a maximum of about 3 to 25 mmol/l within 30 to 120 min following ingestion of 0.3 to 1 g erythritol/kg bw (Noda et al., 1994; Bornet et al., 1996a, b). Bile concentrations of erythritol have been reported to be proportional to plasma erythritol concentrations (Westendorf and Czok, 1983). Erythritol has been reported to transfer across the human placenta (Jansson et al., 1993) and to pass slowly from the plasma to cerebrospinal fluid and brain of sheep (Dziegielewska et al., 1979). No metabolite of erythritol has been observed in rats (Noda and Oku, 1992; Noda et al., 1996; van Ommen et al., 1996) or in humans (Hiele et al., 1993; Noda et al., 1994) indicating that erythritol is not metabolised to a significant extent in the body. Unabsorbed erythritol does however undergo microbial fermentation in the colon to volatile short-chain fatty acids (Noda and Oku, 1990, 1992) or is excreted in the faeces. In the rat at high doses, a higher proportion of the ingested dose undergoes fermentation as the dosage is increased (Noda and Oku, 1992; Oku and Noda 1990a) and as a result of pre-adaptation to erythritol in the diet (Oku and Noda 1990a). In dogs less than 2% of ingested erythritol is excreted unchanged via faeces (Dean et al., 1996; Noda et al., 1996). Similar results have been reported in rats (Lina et al., 1996; Noda and Oku, 1990; Noda et al., 1996; Oku and Noda, 1990a, b; Til et al., 1996; Van Ommen et al., 1996). The biliary excretion of erythritol has been estimated in dogs at 1% (Lewis et al., 1982).

Animal studies

Acute toxicity

When tested for acute toxicity, erythritol is essentially non-toxic e.g. after oral administration LD₅₀ in dogs is greater than 5g/kg bw (Ozeki et al., 1988), LD₅₀ in rats is 13.1 g/kg bw for

males and 13.5 g/kg bw for females (Yamamoto et al. 1987) or greater than 18 g/kg bw (Beck et al., 1938). In these studies, the effects recorded in animals that died were those commonly associated with the dosing of large volumes of hypertonic solutions.

Sub-acute and sub-chronic toxicity

Increased thirst, increase in caecal and/or kidney weights, and transient diarrhoea in the 10% groups and occasionally in 5% groups were the most apparent adverse effects reported (the effects known to occur in rodents fed polyols) in two 28-day studies in rats (Oku and Noda, 1990a; Til and Modderman, 1996) and in a 28-day study in the rat (Shibata et al., 1991), conducted to further evaluate the findings of increased blood urea nitrogen (BUN) observed in some sub-chronic studies (Kamata, 1990a, b; Yamamoto et al., 1989; see below). In another 28-day toxicity study in rats, specifically designed to assess the potential effect of erythritol on the renal function, and in which the highest dose of erythritol was 5% in the diet, the only relevant and statistically significantly altered clinical parameter was an increased water intake in the high-dose group of the sham treated and nephrectomised rats compared to the controls (Kanai et al., 1992). This effect was considered to be evidence of a diuretic effect of erythritol.

In sub-chronic rodent studies (Kamata, 1990a; Til et al., 1991, 1992, 1996; Yamamoto et al., 1989) the following effects were seen in some or all studies: soft stool and/or diarrhoea and reduced body weight in groups exposed to high oral doses of erythritol (20% in the diet or ≥ 4 g/kg bw by gavage), an increased feed intake, increased water intake, changes in some clinico-biochemical parameters (e.g. increased alkaline phosphatase (ALP), decreased γ -glutamyltransferase (GGT), increased BUN, decreased plasma protein and chlorine, decreased sodium, decreased calcium and increased potassium), increased urination/urine volume, changes in several urinary parameters (e.g. increased excretion of sodium, potassium, calcium, chlorine, protein, GGT and *N*-acetyl glucosaminidase (NAG)), changes in organ weights such as increased weights of caecum, kidneys and adrenal glands and decreased thymus weight. Furthermore, a slight dilatation of renal tubules was reported in the high-dose group (8 g/kg bw/day) in one study (Yamamoto et al. 1989). All these effects were considered of non-specific nature being a result of physiological responses to the diuretic and osmotic actions of high doses of erythritol.

Treatment of dogs with erythritol in high doses (5 g/kg bw: Yamaguchi, 1990 and Kamata, 1990b; Dean and Jackson, 1992; Dean et al., 1996: 10% in the diet equivalent to 3.8 g/kg bw) in sub-chronic studies was associated with increased water intake and increased urination/urine volume irrespective of duration of the study and the route of administration (Yamaguchi, 1990: 13 weeks by gavage; Kamata, 1990b: 6 months intravenously; Dean and Jackson, 1992; Dean et al., 1996: 1 year in the diet). Furthermore, some renal changes were recorded in the 13-week study (Yamaguchi, 1990): eosinophilic degeneration, slight dilatation and pycnosis of tubules in 2 of 5 males in the high dose group (5g/kg bw) for each finding and epithelial desquamation, hydropic degeneration, slight necrosis of tubules in one of 5 males in the high dose group. In the 6-month study (Kamata, 1990b), a dose related increase in BUN was recorded but it was considered of no toxicological significance as (i) a renal functional test (*PSP clearance test*) did not reveal any evidence of renal function impairment, (ii) there were no histopathological changes in the kidneys, and (iii) there was no concomitant increase in creatinine concentrations to indicate the presence of renal damage. The results of the 1-year study indicated that ingestion up to 10% erythritol in the diet (equivalent of 3.8

g/kg bw/day) did not cause any overt signs of toxicity (Dean and Jackson, 1992; Dean et al., 1996).

Chronic toxicity and carcinogenicity

One chronic (78-weeks) study in rats with dietary levels of 0, 1, 3, or 10% erythritol (equal to 0.46, 1.4 and 5 g/kg bw/day for males and 0, 0.54, 1.7 and 5.7 g/kg bw/day for females) (Til and van Nesserooij, 1994) and another 2-year chronic toxicity/carcinogenicity study in rats with dietary levels of erythritol of 0, 2, 5, or 10% (equal to 0, 0.9, 2.2, and 4.6 g/kg bw/day for males and 0, 1.0, 2.6, and 5.4 g/kg bw/day) (Lina et al., 1994; 1996) demonstrated that erythritol did not affect survival and had no carcinogenic effect.

The spectrum of effects recorded in the 78-week toxicity study (Til and van Nesserooij, 1994) included soft faeces in the high-dose group during the two first weeks only, slight decrease in mean body weights of both sexes in the 10% erythritol group (the difference with the controls being statistically significant in males on days: 14, 21, 28, 77, 168 and from day 308 onwards), increase in water intake with increasing dietary levels of erythritol in both sexes (the differences with the controls being generally statistically significant in the 10% group and in the 3% and 1% groups on several occasions during the study), increased plasma alkaline phosphatase activity (ALP) (the difference with the controls reaching a statistical significance in week 13 for females and in weeks 26 and 78 for males in the 10% group), higher urine volume in the 10% group (the difference with the controls being statistically significant only in weeks 12 and 25), increased urinary calcium excretion over 16-hour period in both sexes in the 10% group (the difference with the controls being statistically significant at nearly all stages throughout the study^{1 2}), statistically significantly increased absolute and relative weights of the full caecum in the 3% and 10% groups and of the empty caecum in the 10% group, statistically significantly increased relative kidney weights, and absolute and relative spleen weights, and absolute thyroid weights in females in the 3% group.

The effects observed in a 2-year rat study with erythritol (Lina *et al.* 1994, 1996) were: reduced body weights (the difference with the controls being statistically significant for males in the 2% group at termination, in the 5% group from week 8 to the termination and in the 10% group from week 3 to the termination, and in females in the 10% group during most weeks after week 10), increased feed intake in both sexes in the 10% group (the difference being statistically significant on several occasions), increased water intake (the difference being statistically significant on most time points for males in the 10% group and occasionally for males in the 5% and for females in the 10% groups), a few statistical differences at various sampling times in haematological parameters and of blood chemistry parameters, consistent but not statistically significantly increase in plasma ALP of both sexes in the 10% group, increased the 24-hr urine production (the difference being statistically significant in the 5% group in week 26 and in the 10% group in weeks 26, 42, 50 and 78), decreased urine osmolarity in the 10% group (the difference being statistically significant in weeks 26, 42 and 50), changes in urinary pH (statistically significant decrease in the 5% group in week 26 and in the 10% group in weeks 26, 42, 50, 78, and statistically significant increase in the 5% and 10% groups at termination), increased urinary excretion of protein (the difference being

¹ Statistically significantly increased urinary calcium excretion/16hrs in the high-dose group was recorded on day 87 for both sexes, on day 179 for males only, on day 543 for females only.

² Increased calcium excretion could be due to increased absorption from the gut, a phenomenon reported with certain other low molecular weight organic compounds, which are fermented in the colon (poorly absorbed and poorly metabolised carbohydrates).

statistically significant in the 10% group in weeks 42 and 78), of low molecular weight protein (LMP) (the difference being statistically significant in the 10% group in weeks 42, 50 and 78), of enzymes (the difference being statistically significant for GGT in the 2% group in week 50, in the 5% group in weeks 26 and 50 and in the 10% group in weeks 26, 42, 50 and 78, and for NAG in the 5% group in weeks 26 and in the 10% group in weeks 26, 42, 50 and 78), and of electrolytes (the difference being statistically significant in the 10% group for sodium in weeks 26, 50, 78, for potassium in week 78, for calcium on all occasions measured, for phosphate in weeks 26, 42, 50 and 78 and in the 5% group for phosphate in week 26, for citrate on all occasions except for week 26), increased absolute and relative weights of full and empty caecum of both sexes in the 10% group (the difference being statistically significant at all time points i.e. weeks 52, 78 and 102) and in the 5% group (in weeks 72 and 102), increased relative kidney weights in the 10% group (the difference being statistically significant for males in weeks 52 and 78 and for females at termination), increased relative liver weights of males in the 10% group (the difference being statistically significant in week 78 and 102), increased weights of adrenals in the 10% group (the difference being statistically significant for females in the absolute and relative weights at termination and for males in the relative weights in week 78), increased relative weight of spleen of males in the 10% group (the difference being statistically significant in week 78). Histopathological findings revealed statistically increased nephrocalcinosis in the 5 and 10% groups in week 78, and statistically significantly increased pelvic nephrocalcinosis (pelvic epithelial mineralisation) in females of the 2%, 5%, and 10% groups).

Genotoxicity

Mutagenic potential of erythritol was investigated in two Ames' tests (Blijleven, 1990; Kawamura et al., 1996) and in an *in vitro* chromosome aberration test in Chinese hamster fibroblasts (Kawamura et al., 1996; Nakatsuru et al., 1988).

In the first Ames' test (Blijleven, 1990) erythritol, both in the absence and in the presence of an exogenous source of metabolic activation (Aroclor 1254-induced rat liver S9) showed no evidence of mutagenic activity at concentrations of up to 30.0 mg/plate when tested in five histidine-requiring *Salmonella typhimurium* strains (TA1535, TA1537, TA1538, TA98, TA100). Also in the second Ames' test (Kawamura et al., 1996) erythritol, both in the absence and in the presence of an exogenous source of metabolic activation, showed no evidence of mutagenic activity following incubation with histidine-requiring *Salmonella typhimurium* strains (TA98, TA100, TA1537) or with *Escherichia coli* strain WP2 *uvrA* at doses up to 5000 µg/plate. In strain TA1535, in the presence of S9, many of the erythritol treated groups showed a greater number of revertant colonies than the controls. These increases, however, failed to show a dose-response relationship, and tended to show considerable variation in number. As a result, the increased revertant numbers observed in TA1535 were not considered evidence of mutagenic activity.

In an *in vitro* chromosome aberration assay in Chinese hamster fibroblast cells, erythritol at concentrations up to 10.0 mM, both in the absence and in the presence of an exogenous source of metabolic activation, failed to produce a significant increase in the incidence of abnormal cells, polyploid cells, total chromosomal aberrations, break or exchange types aberrations, thus demonstrated no genotoxic activity (Kawamura et al., 1996; Nakatsuru et al., 1998).

Reproductive and developmental toxicity

In a peroral exposure by gavage of mice (doses 0, 1, 2, 4, or 8g/kg bw/day) the effects recorded in maternal animals of both sexes were sporadic diarrhoea and increased water intake at 4 and 8 g/kg bw/day, and dilatation of renal tubules in 2 of 24 males at 8g/kg bw/day. The only statistically significant differences in reproductive parameters, compared to the controls, were a lower implantation rate in the 4 g/kg bw group, and decreased a male: female ratio in the 8 g/kg bw/day group but there was no clear dose-response pattern and both parameters were within the normal historical range (Tateishi, 1989).

In the second study (Tateishi et al., 1992) an intravenous administration of erythritol to mice in doses of 0, 1, 1.73, or 3 g/kg bw/day had no effect on conventional reproductive performance parameters or on foetal development. The effects recorded in maternal animals were limited to the high-dose group and included the death of two males and one female, statistically significantly increased water intake in the female mice before mating, and a dilatation of the renal tubules and a dilatation of the Bowman's capsule in one male and a dilatation of the renal tubules in two females from the high-dose group.

In a 2-generation study in rats (Smits-van Prooije et al., 1996a; Waalkens-Berendsten et al., 1996), groups of 24 animals of both sexes were fed diets containing 0, 2.5, 5, or 10% erythritol for approximately 10 weeks before mating and during the gestation for two consecutive generations (F₀ and F₁) with one litter per generation. Diarrhoea was recorded only in the high-dose group of F₀ and F₁ generations during the first few days of treatment. Body weight of F₀ generation in the high-dose group was below those of controls (the difference being statistically significant for males only). However, the feed intake of F₀ generation in the high-dose group (both sexes) was statistically significantly reduced during the first week only. Thereafter the feed intake of both sexes was statistically significantly higher than in controls with the exception for females in weeks 2 and 3 of gestation and lactation. F₁-males and females exhibited the reduced body weight, the difference from controls being statistically significant for males during week 0-8 and 17-18, and for females during weeks 0-4. However, the rates of body weight gain of males and females of this dose group were not different from those of the controls. Erythritol did not affect reproductive performance of the parental rats (F₀ and F₁). There were no effects on the development of offspring. Histological examination did not reveal any abnormalities.

Results of the three studies indicated that erythritol even at high doses had no adverse effects on fertility or on the developing foetus.

Erythritol was administered intravenously to pregnant mice (mated with untreated males) in doses 0, 1, 2, or 4 g/kg bw/day on days 6-15 of gestation (Ota et al., 1990). Administration of erythritol at 4 g/kg bw/day to dams from F₀-generation was associated with maternal hypoactivity and staggering gait, the death of two dams, periodically decreased feed intake and increased water intake, slightly higher (but not statistically significantly) incidence of fetuses with external abnormalities and a statistically significantly higher incidence of foetuses with skeletal abnormalities. Erythritol in doses up to 4 g/kg bw/day had no effect on body weights of pups (F₁) during lactation and after weaning, or on developmental parameters, behaviour or reproductive performance of the F₁ generation.

When erythritol was administered to pregnant rats from day 0 to day 21 at dietary levels of 0, 2.5, 5, or 10% the only effects recorded were a statistically significantly reduced maternal body weight and body weight gain in week 2 of gestation in the high-dose group, and a statistically significantly higher number of postimplantation losses and higher placental weight in the low-dose group compared to the controls (Smits-van Prooije, 1993; Smits-van Prooije et al., 1996b).

Mated female rabbits received 0, 1, 2.24, or 5 g/kg bw/day of erythritol intravenously on days 6 to 18 of gestation (Hashima Laboratory, 1989; Shimizu et al., 1996). Maternal effects such as polyuria, auricular oedema and lethargy were observed in the high-dose group. Furthermore, water intake was statistically significantly higher than that in the controls from day 7 to 13 of gestation in the low- and mid-dose groups and from day 7 to 17 of gestation in the high-dose group. No effects were observed in the reproductive performance of the dams or on foetal development at any treatment level.

Human studies

Potential influence of erythritol on carbohydrate metabolism

In 5 male healthy volunteers, a single oral dose of erythritol (0.3g/kg bw as 20% aqueous solution) had no significant effect on serum glucose or insulin concentrations. Furthermore, ingestion of erythritol was not associated with any changes in serum cholesterol, triglycerides, free fatty acids, sodium, potassium or chloride levels (Noda et al., 1994). Urinary volume, osmotic pressure, concentration of sodium, potassium and chloride were not significantly different when compared to those after ingestion of the same dose of glucose. Approximately 90% of the ingested dose was excreted in the urine within 48 hours. Furthermore, plasma glucose and insulin levels were not affected in 3 male and 3 female healthy volunteers after a single oral dose of erythritol of 1g/kg bw dissolved in 250 ml of water (Bornet et al., 1996a), or in 12 male and 12 female healthy volunteers after single administration of 0.4 or 0.8 g/kg bw erythritol in form of a midmorning snack (Bornet et al. 1996b).

In 5 noninsulin-dependent diabetic patients (sex not stated), the results of a single dose study (20 g/person in 100 ml aqueous solution) indicated no significant effects of erythritol on carbohydrate metabolism (Ishikawa et al., 1992, 1996). Furthermore, a two-week daily administration of erythritol (20 g/person/day in solution throughout the day with the usual diet) to 11 noninsulin-dependent outpatients (3 males and 8 females) had no effect on blood glucose control (Ishikawa et al. 1996; Miyashita et al. 1993). These limited studies indicate that erythritol does not adversely affect carbohydrate metabolism.

Gastrointestinal tolerance

In studies investigating the laxative effect of erythritol in healthy volunteers after single oral doses up to 78 g (Bornet et al., 1996a) in aqueous solution given on an empty stomach the diarrhoea was not observed at 30 g/person corresponding to 0.46 g/kg bw (Umeki, 1992) and 0.47 g/kg bw (Takahashi, 1992a) in 6/6 and 8/8 males, respectively and to 0.57 g/kg bw in 4/4 females (Takahashi, 1992a). The minimum dose of erythritol causing laxation in these studies ranged from 0.6 to 0.7 g/kg bw. A single administration of erythritol incorporated into a jelly ingested by 14 males and 24 females 2 to 3 hours after a meal did not cause laxation at doses below 0.70 g/kg bw (Oku and Okazaki, 1996). Female subjects seemed to show a greater tolerance than males to the laxative effect of erythritol. Other reported symptoms associated

with diarrhoea were abdominal pain, nausea, intestinal rumbling and/or increased intestinal movements/cramps/spasms, flatulence and thirst. In all studies, the gastrointestinal symptoms showed recovery within 24 hrs after dosing.

In the studies with single daily doses of erythritol mentioned above (Umeki, 1992; Takahashi, 1992 a; Oku and Okazaki, 1996) sucrose and sorbitol were used as reference compounds for gastrointestinal tolerance. Sucrose in tested doses of 0.92 (Umeki, 1992), 1.0 (Takahashi, 1992a) and 1.2 g/kg bw (Oku and Okazaki, 1996), depending on the study, had no laxative effect. Sorbitol in aqueous solution caused laxation in all 6 male subject in a dose of 0.15 g/kg (Umeki, 1992) and in 3 of 8 male subjects at 0.16 g/kg bw (Takahashi, 1992a) while 0.19 g/kg bw had no effect in all 4 females (Takahashi, 1992a). When incorporated into a jelly, sorbitol doses of less than 0.25 g/kg bw caused laxation in 43% (6/14) of the males but not in females (0/24) (Oku and Okazaki, 1996). These results indicate that the laxative effect of erythritol is weaker than that of sorbitol.

When the effect of a repeated dosing was investigated in 8 male and 2 female healthy volunteers, ingestion of erythritol in two daily doses of 20 g in aqueous solution 2-3 hours after breakfast or lunch during 5 consecutive days (corresponding to 40 g/day equivalent to 0.64 and 0.74 g/kg bw in men and women respectively) caused gastrointestinal pain and diarrhoea in one man but not in other subjects (Takahashi, 1992b). In contrast, no signs of gastrointestinal intolerance were recorded in 12 male healthy volunteers when erythritol was given during 7 consecutive days in total daily doses of 0.3 g/kg bw on day 1, of 0.6 g/kg bw on day 2 and of 1 g/kg bw thereafter divided in five portions ingested with food or beverages (Tetzloff et al., 1996). However, when 1 g/kg bw was administered as a single dose in 250 ml water after an overnight fast to 3 male and 3 female healthy volunteers diarrhoea was observed in 2 out of 6 healthy subjects (sex not stated), while the other subjects had abdominal spasms, discomfort and flatulence (Bornet et al., 1996a).

Potential allergenicity of erythritol

Erythritol is a simple sugar alcohol and not known to undergo covalent binding to proteins. It is not a reactive compound and it is not metabolised to reactive metabolites (Bornet et al., 1992; Hiele et al., 1993; Noda et al., 1994). Therefore, it is unlikely that erythritol should cause allergic reactions when consumed with foods. Furthermore, it occurs naturally in some foods, which are not known to be common allergens. Additionally, no reactions, which would indicate any allergic sensitivity in humans, were reported in the above mentioned human studies with erythritol.

When the allergic potential of erythritol was investigated in the heterologous passive cutaneous anaphylaxis reaction in rats, erythritol elicited no hypersensitivity reactions or increases in IgE antibody production (Kawauchi et al., 1989a). Also in guinea pigs several antigenicity tests with erythritol (active systemic anaphylaxis test, homologous passive cutaneous test, active cutaneous anaphylaxis - delayed type hypersensitivity test and passive haemagglutination test) were negative, indicating no immunoreactive or sensitising effect of the compound (Kawauchi et al., 1989b).

There are three well-described cases of severe allergic or allergic-like adverse reactions after erythritol ingestion (Hino et al., 2000; Yunginger et al., 2001). All three patients had a history

of reactions after eating or drinking food with added erythritol: two women developed generalised urticaria and a man generalised urticaria or hypotension. Erythritol itself and not a contaminant seems to be responsible for the reaction; the pathophysiologic mechanisms (e.g. IgE or non IgE mediated) underlying these reactions remain obscure. According to the same authors (Yunginger et al., 2001) the estimated prevalence of adverse reactions to erythritol containing products is less than 1 per million people.

Energy value of erythritol

The caloric value of erythritol, as declared by the petitioners, is 0 - 0.2 kcal/g. The determination of the energy value was performed using a factorial method - a recognised approach for the determination of the energy values of fermentable non-digested carbohydrates (Livesey, 1992).

Some human studies (Noda et al., 1988, 1994; Ishikawa et al., 1992) indicated that 90% or more of erythritol was absorbed and not metabolised following doses of up to 20 g/day. Animal studies suggested a fractional absorption in the small intestine in the same range as described in humans (Nakayama, 1990a; Noda et al., 1996; Oku and Noda, 1990a). Metabolism of erythritol may occur to a very small extent, but no metabolite of this sugar alcohol has been observed in animals (Noda et al., 1996). The study of metabolism of erythritol in humans also suggested a very low rate of metabolism (if any) (Hiele et al., 1993). Erythritol that is not absorbed from the small intestine passes into the large intestine where it can be fermented by colonic microorganisms into short-chain fatty acids (SCFAs) (Noda and Oku, 1992). The conversion of erythritol to SCFAs and gases in the human colon is associated with a loss of about 25% of caloric value. Furthermore, the loss of caloric value of about 25% (from 14 to 30%) due to production of bacterial mass in colon should be added. Other minor losses such as heat production and faecal excretion are difficult to quantify. In all animal studies some faecal erythritol was noted. In studies with adapted rats where erythritol was quantified, faecal excretion of erythritol was usually between 1 and 5% of intake (Lina et al., 1994; Til et al., 1991). In vitro studies in humans (Barry et al., 1992; Hiele et al., 1993) suggested that erythritol is not fermented by human colonic micro-flora to the same extent as it is in rats, but no precise figure can be established.

On the basis of available information concerning the various components in the factorial equation for erythritol the caloric value for erythritol is confirmed to be less than 0.9 kJ/g; or less than 0.2 kcal/g, in humans given oral doses lower than 25g/day or 0.34 g/kg bw.

Comments

The level for lead proposed by the petitioners is twice as much as in the JECFA purity criteria (JECFA, 1999). It seems more appropriate to base the safety of erythritol as a food additive not on the lead level proposed by the petitioners, but on the JECFA criterion for lead (not more than 0.5 mg/kg) considering that erythritol may be used in relatively high quantities in certain foods based on the proposed application and anticipated maximum levels of erythritol in food products.

The effects of erythritol in animals have been investigated in a large number of studies (Dean and Jackson, 1992; Dean et al., 1996; Kamata, 1990a, b; Kanai et al., 1992; Lina et al., 1994,

1996; Oku and Noda, 1990a; Shibata et al., 1991; Smits-van Prooijje et al., 1996a, b; Smits-van Prooijje, 1993; Tateishi, 1989; Til and Modderman, 1996; Til and van Nesselroijl, 1994; Til et al., 1991, 1992, 1996; Waalkens-Berendsten et al., 1996; Yamaguchi, 1990; Yamamoto et al. 1989). They included transient occurrence of loose stool/diarrhoea, decreased body weight gain, increased water consumption, increased urine volume, increased urinary excretion of electrolytes (particularly sodium, potassium and calcium) and urinary enzymes (e.g. NAG, GGT), increased serum ALP and BUN, increased absolute or relative caecal weights, increased absolute or relative kidney weights and histopathological changes in kidneys such as a dilatation of renal tubules, calcium deposits in kidneys/pelvic nephrocalcinosis. These effects were recorded in rats and dogs when high doses of erythritol were used i.e. $\geq 5\%$ in the diet, ≥ 2.5 g/kg bw/day by gavage or ≥ 2.2 g/kg bw/day intravenously.

Other findings reported in sub-chronic and chronic studies mentioned above were minor isolated changes in haematological parameters or in blood chemistry, urinalysis and organ weights. These changes were either of small magnitude, or did not show a dose-response effect, or were not consistent across sex, time and study and therefore were considered to have no relationship to erythritol treatment.

The decreased body weight /body weight gain recorded in some of the sub-chronic and chronic rodent studies and in a two generation reproduction study is most likely attributable to the reduced caloric value of erythritol, as the feed intake was either comparable or slightly increased in the groups treated with high doses of erythritol when compared to the controls. In a short-term study, however, diarrhoea may also have played a role in decreasing the body weight.

The loose stools/diarrhoea in rats and effects on caecal weights in rats and mice treated orally with erythritol could be due to the loading of the large intestine of unabsorbed erythritol at high doses. The fact that the caecal enlargement was not observed in rats receiving erythritol intravenously supports this explanation.

The increased absolute and or /relative kidney weights reported in one of the short-term studies and in the sub-chronic and chronic rodent studies by the oral or intravenous routes of exposure and in one sub-chronic dietary dog study could be the result of the increased urine output recorded in these studies. A possible explanation might be that increased urine output increased the workload of the kidneys, increasing functional tissue activity and therefore increasing the kidney weight. This explanation seemed likely, as no increase in kidney weight was recorded in rats treated intravenously with erythritol for 6 months and allowed a 4-week recovery period, and based on the reported observation that diuresis resulting from a high-dose exposure of rodents to carbohydrates was accompanied by increased absolute and or relative kidney weights (Ogino et al., 1994). Furthermore, the slight dilatation of the renal tubules could be attributable to exposure to the substance with osmotic/diuretic activity.

Other effects such as increased total excretion of electrolytes and of urinary enzymes could also be attributable to the osmotic/diuretic activity of erythritol. In addition, these findings were not consistent from study to study, and were not correlated with any indication of renal disease.

The pelvic nephrocalcinosis recorded in the 2-year rat study was most likely associated with the recorded increase in calcium excretion. It has to be noted that pelvic nephrocalcinosis in that study was not limited to the 10% erythritol treatment but was also recorded for the 10% mannitol group. Furthermore, nephrocalcinosis has also been recorded in other rat studies with poorly absorbed / poorly metabolised carbohydrates (Bär, 1985) and it has been suggested to be associated with the increased calcium excretion as a result of increased calcium absorption due to the osmotic loading of the gastrointestinal tract (Bär, 1985).

The reported increases in serum ALP in rat studies could most likely be attributed to the osmotic activity of erythritol, which reached the large intestine after ingestion at high doses in a similar way as it was demonstrated with other osmotically active carbohydrates (Bär et al. 1995; Moser *et al.* 1980; Schaafsma and Visser 1980; Woutersen 1987).

The finding of an increased BUN recorded in some studies was further elucidated in a 28-day rat study and demonstrated to be associated with loss of electrolytes, particularly sodium, resulting from the diuretic action of erythritol.

Although no gene mutation test in mammalian cells was performed with erythritol the available negative results of Ames' tests and in *in vitro* chromosome aberration test in Chinese hamster fibroblasts were considered sufficient for evaluation of possible mutagenic activity of the compound considering that other polyols were not mutagenic.

The results of human studies on gastrointestinal tolerance indicated that the NOEL for gastrointestinal symptoms was between 0.5 to 1.0 g/kg bw. However, when erythritol was administered in water a dose of 0.5 g/kg bw was the lowest NOEL for diarrhoeic effect. The finding of a relatively higher incidence of gastrointestinal effects with acute bolus dosing in solutions and on an empty stomach may reflect greater passage of the osmotically active erythritol into the large intestine as a result of reduced absorption. The estimation of a daily intake of erythritol based on Danish disappearance and consumption data and under conditions of use proposed by petitioners indicates that the laxative threshold may be exceeded especially by young consumers and through ingestion of erythritol in beverages.

Conclusions

There are extensive animal studies and some human trials on erythritol. The Committee concluded that the effects seen in the animal studies were attributable to physiological and adaptive responses to the rapid absorption and excretion of erythritol and to the osmotic activity of unabsorbed erythritol and its fermentation products in the gut. The intestinal effects are common to all the polyols.

Erythritol does have a laxative effect, but at higher doses than other polyols. The NOEL for the laxative effect of erythritol in humans is around 0.5 g/kg bw for a single dose.

In accordance with the Committee's earlier opinion on other polyols it is considered inappropriate to establish a numerical ADI for erythritol.

The Committee considers that the use of erythritol as a food additive is acceptable. However, as with other polyols, this should not be interpreted as meaning the acceptance of unlimited

use in all foods at any technological level, because the laxative effect should be borne in mind.

The Committee confirms the caloric value of erythritol to be less than 0.9 kJ/g (or less than 0.2 kcal/g) for daily intakes not exceeding 25 g/day.

The Committee recommends that the limit for lead in the specifications should be not higher than 0.5 mg/kg.

References

- Bär A (1985). Safety assessment of polyol sweeteners. Some aspects of toxicity. Food Chem 16: 231-241.
- Barry JL, Hoebler C, Bonnet C, Rival M and David A (1992). In vitro fermentation of indigestible carbohydrates by human faecal flora. INRA, Station of Technology and Applied Nutrition, Nantes, France. Contract N° 34. 92.020.
- Beck F.F., Carr C.J. and Krantz J.C. (Jr.) (1938). Sugar alcohols: XVI – The fate of erythritol and erythritan in the animal body. Qrtrly J Pharmacy Pharmacol 11: 234-239.
- Bernier JJ and Pascal G (1990). Valeur energetique des polyols (sucres-alcools) <The energy value of polyols (sugar alcohols)>. Medecine et Nutrition 26: 221-238.
- Blijleven WGH. (1990). Examination of erythritol for mutagenic activity in the Ames test. Report no. V 90.306, (1990). TNO-CIVO Industries, Netherlands Organization for Applied Scientific Research, The Netherlands.
- Bornet F, Dauchy F, Chevalier A and Slagma G (1992). Étude du devenir metabolique, apres ingestion chez l'homme sain, d'un nuvelédulcorant de charge base caloré: l'erythritol. Gastroenterologie Clinic et Botanique 16: A 169.
- Bornet FRJ, Blayo A, Dauchy F and Slama G (1996a). Plasma and urine kinetics of erythritol after oral ingestion by healthy humans. Regul Toxicol Pharmacol 24: S280-S285.
- Bornet FRJ, Blayo A, Dauchy F, Slama G (1996b). Gastrointestinal response and plasma and urine determination in human subjects given erythritol. Regul Toxicol Pharmacol 24: S296-S302.
- Cerestar Holding BV, Mitsubishi Chemical Corporation and Nikken Chemicals Co., Ltd (1999). Application for assessment of erythritol prior to its authorization.
- Danish Veterinary and Food Administration (DVFA) (2001). Forsyningen af fødevarer 1955-1999 <Danish disappearance and consumption data 1955-1999> (in Danish). Fødevaredirektoratet (2001).
- Dean I and Jackson F (1992). Erythritol: one -year oral (dietary) toxicity study in dogs. Inveresk Research International, Tranent, Scotland, IRI Report no. 7900.
- Dean I, Jackson F and Greenough RJ (1996). Chronic (1-year) oral toxicity study of erythritol in dogs. Regul Toxicol Pharmacol 24: S254-S26.
- Dooms L, Hennebert GL and Verachttert H (1971). Polyol systhesis and taxonomic characters in the genus *Moniliella*. Antonie van Leeuwenhoek 37: 107-118.

Dziegielewska KM, Evans CAN, Malinowska DH, Mollgård K, Reynolds JM, Reynolds ML and Saunders NR (1979). Studies of the development of brain barrier systems to lipid insoluble molecules in foetal sheep. *J Physiol* 292: 207-231.

Hashima Laboratory (1989). Teratology study of NIK-242 in rabbits (intravenous dosing) Seg. II. Final Report. Hashima Laboratory, Japan.

Hiele M, Ghos Y, Rutgeerts P and Vantrappen G (1993). Metabolism of erythritol in humans: comparison with glucose and lactitol. *Br J Nutr* 69: 169-176.

Hino H, Kasai S, Hattori N and Kenjo K (2000). A case of allergic urticaria caused by erythritol. *J Dermatol* 27: 163-165.

De Hoog GS and Gueho E (1984). Deoxyribonucleic acid base composition and taxonomy of *Moniliella* and allied genera. *Antonie van Leeuwenhoek* 50: 135-141.

Inglis GD and Sigler L (1992). *Trichosporonoides megachiliensis*, a new hyphomycete associated with alfalfa leafcutter bees, with notes on *Trichosporonoides* and *Moniliella*. *Mycologia* 84: 555-570.

Ishikawa M, Hirose C, Tsujino D, Miyashita M, Kawashima Y and Nakamura T (1992). The effect of erythritol on glucose tolerance in diabetes patients. Yokohama-shi Seibu Hospital, St. Marianna University School of Medicine, Department of Metabolic Endocrinology and Department of Nutrition (Unpublished internal report).

Ishikawa M, Miyashita M, Kawashima Y, Nakamura T, Saitou N and Modderman J (1996). Effects of oral administration of erythritol on patients with diabetes. *Regul Toxicol Pharmacol* 24: S303-S308.

Jansson T, Powell TL and Illsley NP (1993). Non-electrolyte solute permeabilities of human placental microvillous and basal membranes. *J Physiol* 468: 261-274.

JECFA (1999a). Safety evaluation of certain food additives and contaminants. WHO Food Additive Series, No 44. pp. 15-70.

JECFA (1999b). Specifications for erythritol. FNP 52 Add 7. FAO, Rome.

Kamata S (1990a). A 6 month intravenous chronic toxicity study of NIK-242 in rats with 1 month recovery period: Final report. Study No. SR-8943. Safety Research Institute for Chemical Compounds Co., Ltd., (Internal report).

Kamata S (1990b). A 6 month intravenous chronic toxicity study of NIK-242 in Beagle dogs with 1 month recovery period. Final Report. Study No. SR-8872. Safety Research Institute for Chemical Compounds Co., Ltd., (Internal report).

Kanai M, Yamamoto H, Takahashi T, Onishi T and Shigeki Y (1992). A 4-week feeding toxicity study of erythritol in rats with reduced renal function. Division of Toxicology, Omiya Research Laboratory, Nikken Chemicals Co., Ltd., Japan (Internal report).

Kashima Laboratory Mitsubishi Chemical Safety Institute Ltd. (1995). A 28-day oral toxicity study of cell-free fermented broth in rats. Study No. 4L454, (Internal report).

Kawamura Y, Saito Y., Imamura M and Modderman JP (1996). Mutagenicity studies of erythritol in bacterial reversion assay systems and in Chinese hamster fibroblast cells. *Regul Toxicol Pharmacol* 24: S261-S263.

Kawauchi K, Kodama R, Noguchi K and Suetake K (1989a). Antigenicity study of NIK-242 in mice: heterologous passive cutaneous anaphylaxis (PCA) test with rats. Safety Laboratory, Panapharm Laboratories Co., Ltd., Japan.

Kawauchi K, Kodama R, Noguchi K and Suetake K. (1989b). Antigenicity study of NIK-242 in guinea pigs: active systemic anaphylaxis (ASA) test, homologous passive cutaneous test anaphylaxis (PCA) test, active cutaneous anaphylaxis – delayed type hypersensitivity (ACA-DTH) test and passive hemagglutination test. Safety Laboratory, Panapharm Laboratories Co., Ltd, Japan.

Kockova-Kratochilova M *et al.* (1987). *Moniliella suveolens* var. *Nigra*. *Mykosen* 30: 544-547.

Lewis MH, Baker AL, Dhorajiwala J and Moosa AR (1982). Secretin enhances [¹⁴C]erythritol clearance in unanesthetized dogs. *Digest Dis Sci* 27: 57-64.

Lina BAR (2002). Repeated-dose (28-day) oral toxicity study with erythritol fermentation broth from *Moniliella pollinis* in rats. TNO Nutrition and Food Research, Netherlands Organization for Applied Scientific Research, Zeist, Netherlands TNO Report no. V 4736.

Lina BAR, Bos-Kuijpers MHM and Til HP (1994). Chronic (2-year) oral toxicity and carcinogenicity study with erythritol in rats. TNO-CIVO Industries, Netherlands. Organization for Applied Scientific Research, Zeist, Netherlands TNO Report No. V 93.059.

Lina BAR, Bos-Kuipers MHM, Til HP and Bär A (1996). Chronic toxicity and carcinogenicity study of erythritol in rats. *Regul Toxicol Pharmacol* 24: S264-S279.

Livesey G (1992). The energy values of dietary fibre and sugar alcohols for man. *Nutr Res Rev* 5: 61-84.

MCKenzie RA, Connole MD, McGinnis MR and Lepelaar R (1984). Subcutaneous phaeohyphomycosis caused by *Moniliella suaveolens* in two cats. *Vet. Pathol.* 21: 582-586.

Mitsubishi Chemical Corporation (1991). Natural occurrence of erythritol in foods (Internal report).

Miyashita M, Kawashina Y and Nakamura T (1993). The effect of continuous administration of the sweetener erythritol on diabetes patients. Department of Nutrition

and Department of Metabolic Endocrinology, St Marianna University School of Medicine, (Unpublished report).

Moser RL, Peo ER, Jr, Crenshaw TD and Cunningham PJ (1980). Effect of dietary lactose on weight gain, food conversion, blood, bone and intestinal parameters in postweaning rats and swine. *J Anim Sci* 51: 89-99.

Nakatsuru S, Akiyama S and Saitou SA (1988). Screening for mutagenicity of NIK-242 on reverse mutation test in Bacteria and Chromosome Aberration Test in Cultured Cells. Division of Toxicology, Omiya Research Laboratory, Nikken Chemicals Co., Ltd., Japan, (Internal report).

Nakayama K (1990a). Pharmacokinetics after single oral administration of NIK-242 to dogs. Division of metabolism, Omiya Research Lab., Niken Chemicals Co., Ltd., Japan (Internal report).

Nakayama K (1990b). Blood cell incorporation, protein binding and excretions in urine, faeces, expired air and bile after single oral administration of C-NIK-242 in rats. Division of metabolism, Omiya Research Lab., Niken Chemicals Co., Ltd., Japan (Internal report).

Nakayama K (1990c). Radioactivity distribution after single oral administration of C-NIK-242 in rats. Division of metabolism, Omiya Research Lab., Nikken Chemicals Co., Ltd., Japan, (Internal report).

Nakayama K. (1990d). Blood radioactivity levels after single oral administration of C-NIK-242 in rats. Division of metabolism, Omiya Research Lab., Nikken Chemicals Co., Ltd., Japan (Internal report).

Noda K (1994). Analysis of metabolites in urine after single oral administration of NIK-242 to dogs. Division of metabolism, Omiya Research Lab., Niken Chemicals Co., Ltd., Japan (Internal report).

Noda K and Oku T (1990). The fate and availability of erythritol in rats. In: Hosoya N (Ed) International symposium on caloric evaluation of carbohydrates. Proceedings, Jan. 11-12, 1990, pp. 51-63. Research foundation for sugar metabolism, Tokyo

Noda K and Oku T (1992). Metabolism and disposition of erythritol after oral administration in rats. *J Nutr* 122: 1266-1272.

Noda K, Nakayama K and Modderman J (1996). Fate of erythritol after single oral administration to rats and dogs. *Regul Toxicol Pharmacol* 24: S206-S213.

Noda K, Nakayama K and Oku T (1994). Serum glucose and insulin levels and erythritol balance after oral administration of erythritol in healthy subjects. *Eur J Clin Nutr* 48: 286-292.

Noda K, Nakayama K, Inoue Y (1988). Study on excretion of erythritol after oral administration to human subjects. Division of metabolism, Omiya Research Lab. Nikken chemicals Co., Ltd., Japan, (Internal report).

Ogino Y, Okada S and Ota Z (1994). Effects of chronic, urea-induced osmotic diuresis on kidney weight and function in rats. *Diabetologia* 37: 225-231.

Oku T and Noda K (1990a). Influence of chronic ingestion of newly developed sweetener, erythritol on growth and gastrointestinal function of the rat. *Nutr Res* 10: 987-996.

Oku T and Noda K (1990b). Erythritol balance study and estimation of metabolizable energy of erythritol. In: Hosoya N (Ed) *International symposium on caloric evaluation of carbohydrates*. (Proceedings, Jan. 11-12, 1990) pp. 65-75. Research foundation for sugar metabolism, Tokyo.

Oku T and Okazaki M (1996). Laxative threshold of sugar alcohol erythritol in human subjects. *Nutr Res* 16: 577-589.

Ota T, Kato M and Nakagawa K (1990). Teratology study of NIK-242 in mice (intravenous dosing): Final Report. (1990) Hashima Research Laboratory, Nihon Bioresearch Center Inc., Division of Toxicology, Omiya Research Laboratory, Nikken Chemicals Co., Ltd., Japan (Internal report).

Ozeki M, Hirao A, Araki N, Yamaguchi T, Kimura H, Ito K, Shintani S, Morita K and Kitamura S (1988). Acute oral toxicity study of NIK-242 in dogs. Division of Toxicology, Omiya Research Laboratory, Nikken Chemicals Co., Ltd., Japan, (Internal report).

Samson RA & van Reenen-Hoekstra ES (1988). *Introduction to food-borne fungi*. Centraalbureau voor Schimmelcultures, Baarn, The Netherlands.

Schaafsma G and Visser R (1980). Nutritional interrelationship between calcium, phosphorus and lactose in rats. *J Nutr* 110:1101-1111.

Shibata M, Yamamoto S, Takahashi K, Kitamura S and Ichikawa N (1991). Study on increased BUN caused by repeated administration of erythritol in rats. Division of Toxicology, Omiya Research Laboratory, Nikken Chemicals Co., Ltd., Japan (Internal report).

Shimizu M, Katoh M, Imamura M and Modderman (1996). Teratology study of erythritol in rabbits. *Regul Toxicol Pharmacol* 24: S247-S253.

Shindou T, Sasaki Y, Miki H, Eguchi T, Hagiwara K and Ichikawa T (1989). Identification of erythritol by HPLC and GC_MS and quantitative measurement in pulps of various fruits. *J Agric Food Chem* 37: 1474-1476.

Shindou T, Sasaki Y, Miki H, Eguchi T, Hagiwara K and Ichikawa T (1988). Determination of erythritol in fermented foods by high performance liquid chromatography. *Sokuhin Eiseigaku Zasshi* 29 (6): 419-422.

Shindu T and Ishizuka H (1996). Quantitative determination of erythritol from Various Natural cheeses by HPLC. *Food Sci Technol* 2: 82-83.

Smits-van Prooije AE (1993). Oral embryotoxicity/teratogenicity study with erythritol in rats. Final Report. No. V 92.107. TNO Nutrition and Food Research, The Netherlands.

Smits-Van Prooije AE, Waalkens-Berendsen DH and Bär A (1996a). Dietary two-generation reproduction study with erythritol in rats. TNO report No. V92.552. TNO-CIVO Industries, Netherlands Organization for Applied Scientific Research, Zeist, The Netherlands.

Smits-van Prooije A.E., Waalkens-Berendsen DH and Bär A (1996b). Embryotoxicity and teratogenicity study with erythritol in rats. Regul Toxicol Pharmacol 24: S232-S236.

Takahashi C (1992a). Study on the maximum no-effect level of erythritol using transient diarrhoeal action as index. Abstract. Nikken Chemicals Co. Ltd., Japan (Internal report).

Takahashi C (1992b). Effect of continuous injection of erythritol with laxative action serving as index. Abstract. Nikken Chemicals Co.

Tateishi T (1989). Oral reproduction study of erythritol (NIK-242) with mice prior to and in the early stage of pregnancy. Final Report. Study No. 242040124. Omiya Research Laboratory, Nikken Chemicals Co., Ltd., Japan.

Tateishi T, Yamamoto S, Kagawa M, Mizutani M, Kosuga M, Takahashi K, Itoh K and Kasai Y (1992). Fertility study of NIK-242 in ICR strain mice (Intravenous dosing). Report. Toxicological Laboratory Omiya Research Laboratory, Nikken Chemicals Co., Ltd., Japan, (Internal Report).

Tetzloff W, Dauchy F, Medimagh S, Carr D and Bär A (1996). Tolerance to subchronic, high-dose ingestion of erythritol in human volunteers. Regul Toxicol Pharmacol 24: S286-S295.

Til HP, Falke HE, Kuper CF (1991). An explanatory subchronic feeding study with erythritol in adult male rats (final report). TNO report No. V90.003. TNO-CIVO Industries, Netherlands Organization for Applied Scientific Research, Zeist, Netherlands.

Til HP, Kuper CF and Bruyntjes JP (1992). Subchronic (13-week) feeding study with erythritol in mice (final report). TNO report No. V90.421. TNO-CIVO Industries, Netherlands Organization for Applied Scientific Research, Zeist, Netherlands.

Til HP and van Nesselrooij JHJ (1994). Chronic (78-week) oral toxicity study with erythritol in rats. Report No. V 93.367. TNO-CIVO Industries, Netherlands Organization for Applied Scientific Research, Zeist, Netherlands.

Til HP, Kuper CF, Falke HE and Bär A (1996) Subchronic oral toxicity studies with erythritol in mice and rats. Regul Toxicol Pharmacol 24: S221-S231.

Till HP and Modderman J (1996). Four-Week Oral Toxicity Study with Erythritol in Rats. Regul Toxicol Pharmacol 24: S214-S220.

U.S. Food and Drug Administration, Center for Food Safety and Applied Nutrition, Office of Food Additives Safety: Agency Response Letter: GRAS notice No. GRN 000076, September 2001 (<http://vm.cfsan.fda.gov/~rdb/opa-g76.html>) .

Umeki Y (1992). Study concerning transient diarrhoea induced by erythritol. (Unpublished report).

van Ommen B and de Bie ATHJ (1990). Disposition study with ¹⁴C-erythritol in rats. TNO Report No. V90.307. TNO-CIVO Industries, Netherlands organization for applied scientific research, Zeist, Netherlands.

van Ommen B, de Bie B and Bär A (1996). Disposition of ¹⁴C-erythritol in germfree and convention rats. Regul Toxicol Pharmacol 24: S198-S205.

Voedingsraad (1987). The energy value of sugar alcohols. Voedingsraad (Dutch Nutrition Council), Committee on polyalcohols, The Hague, Netherlands.

Waalkens-Berendsten DH, Smits-Van Prooije E, Wijnands MVM and Bär A (1996). Dietary two-generation reproduction study with erythritol in rats. Regul Toxicol Pharmacol 24: S237-S246.

Westendorf J and Czok G (1983). Die biliäre Ausscheidung choleretisch aktiver Zimtsäure-Derivate durch die Ratte. Zeitschrift für Ernährungswissenschaft 22: 255-270.

Woutersen RA (1987). Chronic toxicity and carcinogenicity of lactitol in rats comparison with lactose. In Low digestibility carbohydrates. (TNO-CIVO Workshop Proceedings, 27-28, 1986, Zeist, The Netherlands), Center for Agricultural publishing and Documentation, Pudoc, Wageningen, Netherlands, pp. 51-60.

Leegwater, VJ Feron and RJJ Hermus pp.51-60. Center for Agricultural Publishing and Documentation, Pudoc, Wageningen, The Netherlands.

Yamaguchi T (1990). A 13-week oral subacute toxicity study of NIK-242 in dogs with 4-week recovery period (PRL/35). Final report. Study No. 242020322. Division of Toxicology, Omiya Research Laboratory, Nikken Chemicals Co., Ltd., (Internal report).

Yamamoto H, Tateishi T, Sadamasu K, Kosuge M, Takahashi K, Nakano S and Kasai Y (1987). Acute intravenous, subcutaneous and oral toxicity study with NIK-242 in rats. Report. Division of Toxicology, Omiya Research Laboratory, Nikken Chemicals Co., Ltd., Japan, (Internal report).

Yamamoto H, Tateishi T, Touchi T, Kosuge M, Takahashi K, Takahashi T, Nakano S and Kasai Y (1989). 13-week oral subacute toxicity study of NIK-242 in rats with 4-week recovery period (PRL/34). Division of Toxicology, Omiya Research Laboratory, Nikken Chemicals Co., Ltd., Japan (Internal report).

Yunginger JW, Jones RT, Hirohito K, Katsuhiko S, Hefle SL and Taylor SL (2001). Allergic reactions after ingestion of erythritol-containing foods and beverages. *J Allergy Clin Immunol* 108: 650.

Appendix
List of short-term, sub-chronic, chronic, reproductive and developmental toxicity studies and no-effect and effect levels

Study	Reference	NOEL	LOEL
28-day toxicity study in rats	Oku and Noda, 1990a	Not established	5% in the diet
28-day toxicity study in rats	Til and Modderman, 1996	Not established	5% in the diet
Sub-chronic (13-week) feeding study with erythritol in mice	Til <i>et al.</i> , 1996	5% in the diet	10% in the diet
Sub-chronic (13-week) feeding study with erythritol in rats	Til <i>et al.</i> , 1996	5% in the diet	10% in the diet
A 13-week oral subacute toxicity study of NIK-242 with four week recovery period in rats. (gavage)	Yamamoto <i>et al.</i> , 1989	2 g/kg bw	4 g/kg bw
A 6 months intravenous chronic toxicity study of NIK-242 in rats with 1 month recovery period	Kamata, 1990a	1 g/kg bw	1.73 g/kg bw
A 13-week oral subacute toxicity study of NIK-242 in dogs with 4-week recovery period. (gavage)	Yamaguchi, 1990	1.25 g/kg bw	2.5 g/kg bw
A 6 months intravenous chronic toxicity study of NIK-242 in Beagle dogs with 1 month recovery period	Kamata, 1990b	Not established	1 g/kg bw
Chronic (1-year) oral toxicity study of erythritol in dogs	Dean <i>et al.</i> , 1996; Dean and Jackson, 1992	5% in the diet (corresponding to 1.7 g/kg bw)	10% in the diet (corresponding to 3.8 g/kg bw) (<i>the highest dose used</i>)
Chronic (78-week) oral toxicity study with erythritol in rats	Til and van Nesselrooij, 1994	3% in the diet (corresponding to 1.5 g/kg bw)	10% in the diet (corresponding to 5.4 g/kg bw) (<i>the highest dose used</i>)

Chronic (2-year) oral toxicity and carcinogenicity study with erythritol in rats	Lina <i>et al.</i> , 1996, Lina <i>et al.</i> , 1994	2% in the diet (corresponding to 1 g/kg bw)	5% in the diet (corresponding to 2.4 g/kg bw)
Oral reproduction study of erythritol (NIK-242) with mice prior to and in the early stage of pregnancy (gavage)	Tateishi, 1989	General toxicity: 2 g/kg bw Reproductive performance and foetal development: 8 g/kg bw	General toxicity: 4 g/kg bw Reproductive performance and foetal development: No LOEL as the highest dose was NOEL
Fertility study of NIK-242 in ICR strain mice (intravenous dosing)	Tateishi <i>et al.</i> , 1992	General toxicity: 1.73 g/kg bw Reproductive performance and foetal development: 3 g/kg bw	General toxicity: 3 g/kg bw Reproductive performance and foetal development: No LOEL as the highest dose was NOEL
Dietary two-generation reproduction study with erythritol in rats	Smits-van Prooijje <i>et al.</i> , 1996a, Waalkens-Berendsten <i>et al.</i> , 1996	10% in the diet (corresponding to 7.6 g/kg bw)	No LOEL as the highest dose was NOEL
Teratology study of NIK-242 in mice (intravenous dosing)	Ota <i>et al.</i> , 1990	Maternal and developmental toxicity: 2 g/kg bw	Maternal and developmental toxicity: 4 g/kg bw (the highest dose used)
Oral embryotoxicity/teratogenicity study with erythritol in rats	Smits-van Prooijje, 1993, Smits-van Prooijje <i>et al.</i> , 1996b	Maternal toxicity: 5% in the diet (corresponding to 3.4 g/kg bw) Developmental toxicity: 10% in the diet (corresponding to 6.7 g/kg bw)	Maternal toxicity: 10% in the diet (corresponding to 6.7 g/kg bw) Developmental toxicity: No LOEL as the highest dose was NOEL
Teratology study of erythritol in rabbits (intravenous dosing)	Shimizu <i>et al.</i> , 1996, Hashima Laboratory, 1989	Maternal toxicity: 2.24 g/kg bw Reproductive performance and foetal development: 5 g/kg bw	Maternal toxicity: 5 g/kg bw (the highest dose used) Reproductive performance and foetal development: No LOEL as the highest dose was NOEL