

**ANNEX I**

**SUMMARY OF PRODUCT CHARACTERISTICS**

## **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Flexicam 1.5 mg/ml oral suspension for dogs.

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each ml of Flexicam 1.5 mg/ml oral suspension contains:

### **Active substance:**

Meloxicam 1.5 mg  
(equivalent to 0.05 mg per drop)

### **Excipients:**

Sodium benzoate 1.5 mg  
(equivalent to 0.05 mg per drop)

For a full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

Oral suspension.

Flexicam is a pale green, uniform suspension containing 1.5 mg/ml meloxicam.

## **4. CLINICAL PARTICULARS**

### **4.1 Target species**

Dogs.

### **4.2 Indications for use, specifying the target species**

Alleviation of inflammation and pain in both acute and chronic musculo-skeletal disorders.

### **4.3 Contraindications**

Do not use in pregnant or lactating animals.

Do not use in animals suffering from gastrointestinal disorders such as irritation and haemorrhage, impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of individual hypersensitivity to the product.

Do not use in dogs less than 6 weeks of age.

### **4.4 Special warnings**

None.

### **4.5 Special precautions for use**

#### **Special precautions for use in animals**

If side effects occur, treatment should be discontinued and the advice of the veterinarian should be sought. Avoid use in any dehydrated, hypovolaemic or hypotensive animal, as there is a potential risk of increased renal toxicity.

## **Special precautions to be taken by the person administering the veterinary medicinal product to animals**

People with known hypersensitivity to NSAIDs should avoid contact with the veterinary medicinal product.

In case of accidental ingestion, seek medical advice immediately and show the package insert or the label to the physician.

### **4.6 Adverse reactions (frequency and seriousness)**

Typical adverse drug reactions of NSAIDs such as loss of appetite, vomiting, diarrhoea, faecal occult blood and apathy have occasionally been reported. These side effects occur generally within the first treatment week and are in most cases transient and disappear following termination of the treatment but in very rare cases may be serious or fatal.

### **4.7 Use during pregnancy, lactation or lay**

The safety of the veterinary medicinal product has not been established during pregnancy and lactation (see 4.3).

### **4.8 Interaction with other medicinal products and other forms of interaction**

Other NSAIDs, diuretics, anticoagulants, aminoglycoside antibiotics and substances with high protein binding may compete for binding and thus lead to toxic effects. Flexicam must not be administered in conjunction with other NSAIDs or glucocorticosteroids.

Pre-treatment with anti-inflammatory substances may result in additional or increased adverse effects and accordingly a treatment-free period with such drugs should be observed for at least 24 hours before commencement of treatment. The treatment-free period, however, should take into account the pharmacokinetic properties of the products used previously.

### **4.9 Amounts to be administered and administration route**

To be administered mixed with food.

Initial treatment is a single dose of 0.2 mg meloxicam/kg body weight on the first day. Treatment is to be continued once daily by oral administration (24-hour intervals) at a maintenance dose of 0.1 mg meloxicam/kg body weight.

Particular care should be taken with regard to the accuracy of dosing.

The suspension can be given using either the drop dispenser (for very small breeds) or the Flexicam measuring syringe provided in the package. The dispenser provides 0.05 mg meloxicam per drop (ie. a dose of 0.1 mg meloxicam/kg body weight corresponds to 2 drops/kg body weight). The syringe fits onto the bottle and has a kg-body weight scale which corresponds to the maintenance dose (ie. 0.1 mg meloxicam/kg body weight). Thus for the first day, twice the maintenance volume will be required.

Shake well before use.

A clinical response is normally seen within 3-4 days. Treatment should be discontinued after 10 days at the latest if no clinical improvement is apparent.

Avoid introduction of contamination during use.

### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

In the case of overdosage symptomatic treatment should be initiated.

#### **4.11 Withdrawal periods**

Not applicable.

### **5. PHARMACOLOGICAL PROPERTIES**

Pharmacotherapeutic group: Non-steroidal, anti-inflammatory drug (NSAID).  
ATCvet Code: QM01AC06.

#### **5.1 Pharmacodynamic properties**

Meloxicam is a non-steroidal anti-inflammatory drug (NSAID) of the oxicam class which acts by inhibition of prostaglandin synthesis, thereby exerting anti-inflammatory, analgesic, anti-exudative and antipyretic effects. It reduces leukocyte infiltration into the inflamed tissue. To a minor extent it also inhibits collagen-induced thrombocyte aggregation. *In vitro* and *in vivo* studies demonstrated that meloxicam inhibits cyclooxygenase-2 (COX-2) to a greater extent than cyclooxygenase-1 (COX-1).

#### **5.2 Pharmacokinetic particulars**

##### Absorption

Meloxicam is completely absorbed following oral administration and maximal plasma concentrations are obtained after approximately 7.5 hours. When the product is used according to the recommended dosage regime, steady state concentrations of meloxicam in plasma are reached on the second day of treatment.

##### Distribution

There is a linear relationship between the dose of meloxicam administered and plasma concentration observed in the therapeutic dose range. Approximately 97% of meloxicam is bound to plasma proteins.

The volume of distribution is 0.31/kg.

##### Metabolism

Meloxicam is predominantly found in plasma and is also a major biliary excretion product whereas urine contains only traces of the parent compound. Meloxicam is metabolised to an alcohol, an acid derivative and to several polar metabolites. All major metabolites have been shown to be pharmacologically inactive.

##### Elimination

Meloxicam is eliminated with a half-life of 24 hours. Approximately 75% of the administered dose of meloxicam is eliminated via faeces and the remainder via urine.

### **6. PHARMACEUTICAL PARTICULARS**

#### **6.1 List of excipients**

Dispersible cellulose  
Xanthan gum  
Sodium benzoate  
Glycerol  
Xylitol  
Sodium lauryl sulphate  
Citric acid monohydrate  
Sodium citrate

Honey flavour  
Simethicone emulsion  
Purified water

## **6.2 Incompatibilities**

Not applicable.

## **6.3 Shelf life**

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years.  
Shelf-life after first opening the immediate packaging: 9 months.

## **6.4 Special precautions for storage**

This veterinary medicinal product does not require any special storage conditions.

## **6.5 Nature and composition of immediate packaging**

Polyethylene bottle containing 10, 32 or 100 ml with a polyethylene dropper, a tamper proof child resistant closure and a polypropylene measuring syringe.

## **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Dechra Veterinary Products A/S  
Mekuvej 9  
7171 Uldum  
Denmark

## **8. MARKETING AUTHORISATION NUMBERS**

EU/2/06/058/001  
EU/2/06/058/002  
EU/2/06/058/003

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

10.04.2006

**10 DATE OF REVISION OF THE TEXT**

18.03.2010

Detailed information on this veterinary medicinal product is available on the website of the European Medicines Agency (European Medicines Agency) <http://www.ema.europa.eu/>.

**PROHIBITION OF SALE, SUPPLY AND/OR USE**

Not applicable.

## 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Flexicam 5 mg/ml solution for injection for dogs and cats.

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ml of Flexicam 5 mg/ml solution for injection contains:

### **Active substance:**

Meloxicam 5 mg

### **Excipients:**

Ethanol, anhydrous 150 mg

For a full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Solution for injection.

## 4. CLINICAL PARTICULARS

### 4.1 Target species

Dogs and cats.

### 4.2 Indications for use, specifying the target species

#### Dogs:

Alleviation of inflammation and pain in both acute and chronic musculo-skeletal disorders. Reduction of post-operative pain and inflammation following orthopaedic and soft tissue surgery.

#### Cats:

Reduction of post-operative pain after ovariohysterectomy and minor soft tissue surgery.

### 4.3 Contraindications

Do not use in pregnant or lactating animals.

Do not use in animals suffering from gastrointestinal disorders such as irritation and haemorrhage, impaired hepatic, cardiac or renal function and haemorrhagic disorders.

Do not use in case of hypersensitivity to the active substance or to any of the excipients.

Do not use in animals less than 6 weeks of age nor in cats of less than 2 kg.

Do not use an oral follow-up therapy using meloxicam or other NSAIDs in cats, as no safe dosage for repeated oral administration has been established.

### 4.4 Special warnings for each target species

For post-operative pain relief in cats, safety has only been documented after thiopental/halothane anaesthesia.

#### **4.5 Special precautions for use**

##### **Special precautions for use in animals**

If side effects occur, treatment should be discontinued.

Avoid use in any dehydrated, hypovolaemic or hypotensive animal, as there is a potential risk of increased renal toxicity.

##### **Special precautions to be taken by the person administering the veterinary medicinal product to animals**

Accidental self-injection may give rise to pain.

People with known hypersensitivity to meloxicam should avoid contact with the veterinary medicinal product.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

#### **4.6 Adverse reactions (frequency and seriousness)**

Typical adverse drug reactions of NSAIDs such as loss of appetite, vomiting, diarrhoea, faecal occult blood and apathy have occasionally been reported. In dogs, these side effects occur generally within the first treatment week and are in most cases transient and disappear following termination of the treatment but in very rare cases may be serious or fatal.

#### **4.7 Use during pregnancy, lactation or lay**

The safety of the veterinary medicinal product has not been established during pregnancy and lactation (see 4.3).

#### **4.8 Interaction with other medicinal products and other forms of interaction**

Other NSAIDs, diuretics, anticoagulants, aminoglycoside antibiotics and substances with high protein binding may compete for binding and thus lead to toxic effects. Flexicam must not be administered in conjunction with other NSAIDs or glucocorticosteroids. Concurrent administration of potential nephrotoxic drugs should be avoided. In animals at anaesthetic risk (e.g. aged animals) intravenous or subcutaneous fluid therapy during anaesthesia should be taken into consideration. When anaesthesia and NSAID are concomitantly administered, a risk for renal function cannot be excluded.

Pre-treatment with anti-inflammatory substances may result in additional or increased adverse effects and accordingly a treatment-free period with such drugs should be observed for at least 24 hours before commencement of treatment. The treatment-free period, however, should take into account the pharmacokinetic properties of the products used previously.

#### **4.9 Amounts to be administered and administration route**

##### Dogs:

Musculo-skeletal disorders:

Single subcutaneous injection at a dosage of 0.2 mg meloxicam/kg body weight (i.e. 0.4 ml/10 kg body weight).

Flexicam 1.5 mg/ml oral suspension may be used for continuation of treatment at a dosage of 0.1 mg meloxicam/kg body weight, 24 hours after administration of the injection.

Reduction of post-operative pain (over a period of 24 hours):

Single intravenous or subcutaneous injection at a dosage of 0.2 mg meloxicam/kg body weight (i.e. 0.4 ml/10 kg body weight) before surgery, for example at the time of induction of anaesthesia.



#### Cats:

Reduction of post-operative pain:

Single subcutaneous injection at a dosage of 0.3 mg meloxicam/kg body weight (i.e. 0.06 ml/kg body weight) before surgery, for example at the time of induction of anaesthesia.

Particular care should be taken with regard to the accuracy of dosing.

Avoid introduction of contamination during use.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

In the case of overdosage symptomatic treatment should be initiated.

#### **4.11 Withdrawal periods**

Not applicable.

### **5. PHARMACOLOGICAL PROPERTIES**

Pharmacotherapeutic group: Anti-inflammatory and antirheumatic products, non-steroids (oxicams).  
ATCvet code: QM01AC06.

#### **5.1 Pharmacodynamic properties**

Meloxicam is a non-steroidal anti-inflammatory drug (NSAID) of the oxicam class, which acts by inhibition of prostaglandin synthesis, thereby exerting anti-inflammatory, analgesic, anti-exudative and antipyretic effects. It reduces leukocyte infiltration into the inflamed tissue. To a minor extent it also inhibits collagen-induced thrombocyte aggregation. *In vitro* and *in vivo* studies demonstrated that meloxicam inhibits cyclooxygenase-2 (COX-2) to a greater extent than cyclooxygenase-1 (COX-1).

#### **5.2 Pharmacokinetic particulars**

##### Absorption

Following subcutaneous administration, meloxicam is completely bioavailable and maximal mean plasma concentrations of 0.73 µg/ml in dogs and 1.1 µg/ml in cats were reached approximately 2.5 hours and 1.5 hours post administration, respectively.

##### Distribution

There is a linear relationship between the dose administered and plasma concentration observed in the therapeutic dose range in dogs. More than 97% of meloxicam is bound to plasma proteins. The volume of distribution is 0.3 l/kg in dogs and 0.09 l/kg in cats.

##### Metabolism

In dogs, meloxicam is predominantly found in plasma and is also a major biliary excretion product whereas urine contains only traces of the parent compound. Meloxicam is metabolised to an alcohol, an acid derivative and to several polar metabolites. All major metabolites have been shown to be pharmacologically inactive.

##### Elimination

Meloxicam is eliminated with a half-life of 24 hours in dogs and 15 hours in cats. Approximately 75% of the administered dose is eliminated via faeces and the remainder via urine.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Ethanol anhydrous  
Poloxamer 188  
Glycofurol  
Meglumine  
Glycine  
Sodium Chloride  
Sodium Hydroxide  
Water for injection

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years.  
Shelf-life after first opening the immediate packaging: 28 days.

### **6.4. Special precautions for storage**

This veterinary medicinal product does not require any special storage conditions.

### **6.5 Nature and composition of immediate packaging**

Colourless type I glass injection vial of 10 ml, closed with a grey EPDM rubber stopper and sealed with a flip-off aluminium violet seal.

### **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Dechra Veterinary Products A/S  
Mekuvej 9  
7171 Uldum  
Denmark

## **8. MARKETING AUTHORISATION NUMBER**

EU/2/06/058/004

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

09.12.2008

**10 DATE OF REVISION OF THE TEXT**

18.03.2010

Detailed information on this veterinary medicinal product is available on the website of the European Medicines Agency (European Medicines Agency) <http://www.ema.europa.eu/>.

**PROHIBITION OF SALE, SUPPLY AND/OR USE**

Not applicable.

## **ANNEX II**

- A. MANUFACTURING AUTHORISATION HOLDERS RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS OF THE MARKETING AUTHORISATION REGARDING SUPPLY OR USE**
- C. CONDITIONS OR RESTRICTIONS OF THE MARKETING AUTHORISATION WITH REGARD TO SAFE AND EFFECTIVE USE**
- D. STATEMENT OF THE MRLs**

**A. MANUFACTURING AUTHORISATION HOLDERS RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturers responsible for batch release

Flexicam 1.5 mg/ml oral suspension for dogs:

Fisher Clinical Services UK Ltd.  
Langhurstwood Road  
Horsham  
West Sussex  
RH12 4QD  
United Kingdom

Dechra Veterinary Products A/S  
Mekuvej 9  
7171 Uldum  
Denmark

Flexicam 5 mg/ml solution for injection for dogs and cats:

Accord Healthcare Limited  
Sage House 1<sup>st</sup> Floor, 319 Pinner Road  
North Harrow, Middlesex HA1 4HF  
United Kingdom

**B. CONDITIONS OR RESTRICTIONS OF THE MARKETING AUTHORISATION REGARDING SUPPLY OR USE**

To be supplied only on veterinary prescription.

**C. CONDITIONS OR RESTRICTIONS OF THE MARKETING AUTHORISATION WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE PRODUCT**

Not applicable.

**D. STATEMENT OF THE MRLs**

Not applicable.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**10 ml bottle**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Flexicam 1.5 mg/ml oral suspension for dogs

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Meloxicam 1.5 mg/ml

**3. PHARMACEUTICAL FORM**

Oral suspension

**4. PACKAGE SIZE**

10 ml

**5. TARGET SPECIES**

Dogs

**6. INDICATION(S)**

**7. METHOD AND ROUTE OF ADMINISTRATION**

Shake well before use.  
To be administered mixed with food.  
Avoid introduction of contamination during use.

**8. WITHDRAWAL PERIOD**

Not applicable.

**9. SPECIAL WARNINGS, IF NECESSARY**

Do not use in pregnant or lactating animals.  
Read the package leaflet before use.



**10. EXPIRY DATE**

EXP: { month/year }

Shelf-life of opened bottle: 9 months.

**11. SPECIAL STORAGE CONDITIONS**

Keep out of reach and sight of children.

Shelf-life of opened bottle: 9 months.

Do not use after the expiry date stated on the carton and the bottle.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Dispose of in accordance with local requirements.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable**

For animal treatment only - veterinary medicinal product subject to prescription.

**14. THE WORDS “KEEP OUT OF THE REACH AND SIGHT OF CHILDREN”**

Keep out of the reach and sight of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Dechra Veterinary Products A/S  
Mekuvej 9  
7171 Uldum  
Denmark

**16. MARKETING AUTHORISATION NUMBER**

EU/2/06/058/001

**17. MANUFACTURER’S BATCH NUMBER**

Lot: { number }

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

32 ml bottle

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Flexicam 1.5 mg/ml oral suspension for dogs

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Meloxicam 1.5 mg/ml

**3. PHARMACEUTICAL FORM**

Oral suspension

**4. PACKAGE SIZE**

32 ml

**5. TARGET SPECIES**

Dogs

**6. INDICATION(S)**

**7. METHOD AND ROUTE OF ADMINISTRATION**

Shake well before use.  
To be administered mixed with food.  
Avoid introduction of contamination during use.

**8. WITHDRAWAL PERIOD**

Not applicable.

**9. SPECIAL WARNINGS, IF NECESSARY**

Do not use in pregnant or lactating animals.  
Read the package leaflet before use.

**10. EXPIRY DATE**

EXP: { month/year }

Shelf-life of opened bottle: 9 months.

**11. SPECIAL STORAGE CONDITIONS**

Keep out of reach and sight of children.

Shelf-life of opened bottle: 9 months.

Do not use after the expiry date stated on the carton and the bottle.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Dispose of in accordance with local requirements.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable**

For animal treatment only – veterinary medicinal product subject to prescription.

**14. THE WORDS “KEEP OUT OF THE REACH AND SIGHT OF CHILDREN”**

Keep out of the reach and sight of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Dechra Veterinary Products A/S  
Mekuvej 9  
7171 Uldum  
Denmark

**16. MARKETING AUTHORISATION NUMBER**

EU/2/06/058/002

**17. MANUFACTURER’S BATCH NUMBER**

Lot: { number }

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**100 ml bottle**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Flexicam 1.5 mg/ml oral suspension for dogs

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Meloxicam 1.5 mg/ml

**3. PHARMACEUTICAL FORM**

Oral suspension

**4. PACKAGE SIZE**

100 ml

**5. TARGET SPECIES**

Dogs

**6. INDICATION(S)**

**7. METHOD AND ROUTE OF ADMINISTRATION**

Shake well before use.  
To be administered mixed with food.  
Avoid introduction of contamination during use.

**8. WITHDRAWAL PERIOD**

Not applicable.

**9. SPECIAL WARNINGS, IF NECESSARY**

Do not use in pregnant or lactating animals.  
Read the package leaflet before use.

**10. EXPIRY DATE**

EXP: {month/year}

Shelf-life of opened bottle: 9 months.

**11. SPECIAL STORAGE CONDITIONS**

Keep out of the reach and sight of children.

Shelf-life of opened bottle: 9 months.

Do not use after the expiry date stated on the carton and the bottle.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Dispose of in accordance with local requirements.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable**

For animal treatment only – veterinary medicinal product subject to prescription.

**14. THE WORDS “KEEP OUT OF THE REACH AND SIGHT OF CHILDREN”**

Keep out of the reach and sight of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Dechra Veterinary Products A/S

Mekuvej 9

7171 Uldum

Denmark

**16. MARKETING AUTHORISATION NUMBER**

EU/2/06/058/003

**17. MANUFACTURER’S BATCH NUMBER**

Lot: {number}

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**10 ml glass vial**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Flexicam 5 mg/ml solution for injection for dogs and cats  
Meloxicam

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Meloxicam 5 mg/ml  
Ethanol 150 mg/ml

**3. PHARMACEUTICAL FORM**

Solution for injection

**4. PACKAGE SIZE**

10 ml

**5. TARGET SPECIES**

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

**8. WITHDRAWAL PERIOD**

**9. SPECIAL WARNING(S), IF NECESSARY**

Do not use in pregnant or lactating animals.

**10. EXPIRY DATE**

EXP: {month/year}

Shelf-life after first opening the container: 28 days.

Once broached, use by:

**11. SPECIAL STORAGE CONDITIONS**

This veterinary medicinal product does not require any special storage conditions.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Dispose of in accordance with local requirements.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable**

For animal treatment only - to be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE REACH AND SIGHT OF CHILDREN”**

Keep out of the reach and sight of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Dechra Veterinary Products A/S

Mekuvej 9

7171 Uldum

Denmark

**16. MARKETING AUTHORISATION NUMBER**

EU/2/06/058/004

**17. MANUFACTURER'S BATCH NUMBER**

Lot: {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**10 ml bottle**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Flexicam 1.5 mg/ml oral suspension for dogs

**2. QUANTITY OF THE ACTIVE SUBSTANCE**

Meloxicam 1.5 mg/ml

**3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

10 ml

**4. ROUTE(S) OF ADMINISTRATION**

**5. WITHDRAWAL PERIOD**

Not applicable.

**6. BATCH NUMBER**

Lot: {number}

**7. EXPIRY DATE**

EXP: {month/year}

Shelf-life of opened bottle: 9 months.

**8. THE WORDS "FOR ANIMAL TREATMENT ONLY"**

For animal treatment only.



**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

32 ml bottle

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Flexicam 1.5 mg/ml oral suspension for dogs

**2. QUANTITY OF THE ACTIVE SUBSTANCE**

Meloxicam 1.5 mg/ml

**3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

32 ml

**4. ROUTE(S) OF ADMINISTRATION**

**5. WITHDRAWAL PERIOD**

Not applicable.

**6. BATCH NUMBER**

Lot: {number}

**7. EXPIRY DATE**

EXP: {month/year}

Shelf-life of opened bottle: 9 months.

**8. THE WORDS "FOR ANIMAL TREATMENT ONLY"**

For animal treatment only.

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**100 ml bottle**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Flexicam 1.5 mg/ml oral suspension for dogs

**2. QUANTITY OF THE ACTIVE SUBSTANCE**

Meloxicam 1.5 mg/ml

**3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

100 ml

**4. ROUTES OF ADMINISTRATION**

Shake well before use.  
To be administered mixed with food.  
Avoid introduction of contamination during use.

**5. WITHDRAWAL PERIOD**

Not applicable.

**6. BATCH NUMBER**

Lot: {number}

**7. EXPIRY DATE**

EXP: {month/year}  
Shelf-life of opened bottle: 9 months.

**8. THE WORDS "FOR ANIMAL TREATMENT ONLY"**

For animal treatment only.

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**10 ml glass vial**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Flexicam 5 mg/ml solution for injection for dogs and cats  
Meloxicam

**2. QUANTITY OF THE ACTIVE SUBSTANCE(S)**

**3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

10 ml

**4. ROUTES OF ADMINISTRATION**

Dogs: i.v. or s.c.  
Cats: s.c.

**5. WITHDRAWAL PERIOD**

**6. BATCH NUMBER**

Lot: {number}

**7. EXPIRY DATE**

EXP: {month/year}

**8. THE WORDS "FOR ANIMAL TREATMENT ONLY"**

For animal treatment only.

**B. PACKAGE LEAFLET**

**PACKAGE LEAFLET**  
**Flexicam 1.5 mg/ml oral suspension for dogs**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

**Marketing authorisation holder:**

Dechra Veterinary Products A/S  
Mekuvej 9  
7171 Uldum  
Denmark

**Manufacturer for the batch release:**

Fisher Clinical Services UK Ltd.  
Langhurstwood Road  
Horsham  
West Sussex  
RH12 4QD  
United Kingdom

Dechra Veterinary Products A/S  
Mekuvej 9  
7171 Uldum  
Denmark

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Flexicam 1.5 mg/ml oral suspension for dogs.

**3. STATEMENT OF THE ACTIVE SUBSTANCE AND OTHER INGREDIENTS**

Meloxicam 1.5 mg/ml (equivalent to 0.05 mg per drop).

**4. INDICATIONS**

Alleviation of inflammation and pain in both acute and chronic musculo-skeletal disorders.

**5. CONTRAINDICATIONS**

Do not use in pregnant or lactating animals.

Do not use in animals suffering from gastrointestinal disorders such as irritation and haemorrhage, impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of individual hypersensitivity to the product.

Do not use in dogs less than 6 weeks of age.

**6. ADVERSE REACTIONS**

Typical adverse drug reactions of NSAIDs such as loss of appetite, vomiting, diarrhoea, faecal occult blood and apathy have occasionally been reported. These side effects occur generally within the first treatment week and are in most cases transient and disappear following termination of the treatment but in very rare cases may be serious or fatal.

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

## 7. TARGET SPECIES

Dogs.

## 8. DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION

Initial treatment is a single dose of 0.2 mg meloxicam/kg body weight on the first day. Treatment is to be continued once daily by oral administration (at 24- hour intervals) at a maintenance dose of 0.1 mg meloxicam/kg body weight.


Shake well before use. To be administered mixed with food.

The suspension can be given using either the drop dispenser (for very small breeds) or the Flexicam measuring syringe provided in the package (see below). The dispenser provides 0.05mg meloxicam per drop (i.e. a dose of 0.1 mg meloxicam/kg body weight corresponds to 2 drops/kg body weight). The syringe fits onto the bottle and has a kg- body weight scale which corresponds to the maintenance dose (i.e. 0.1 mg meloxicam/kg body weight). Thus for the first day, twice the maintenance volume will be required.

### Dosing procedure using the measuring syringe:



Shake bottle well. Push down and unscrew bottle top. Attach the dosing syringe to the bottle by gently pushing the end onto the top of the bottle.



Turn the bottle/syringe upside down. Pull the plunger out until the black line on the plunger corresponds to your dog's bodyweight in kilograms.



Turn the bottle right way up and with a twisting movement separate the dosing syringe from the bottle.



By pushing the plunger in, empty the contents of the syringe onto the food.

A clinical response is normally seen within 3-4 days. Treatment should be discontinued after 10 days at the latest if no clinical improvement is apparent.  
Avoid introduction of contamination during use.

## **9. ADVICE ON CORRECT ADMINISTRATION**

Particular care should be taken with regard to the accuracy of dosing. Please carefully follow the instructions of the veterinarian.

## **10. WITHDRAWAL PERIOD**

Not applicable.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the reach and sight of children.  
Do not use after the expiry date stated on the carton and the bottle after EXP.  
Shelf-life after first opening the container: 9 months.

## **12. SPECIAL WARNINGS**

If side effects occur, treatment should be discontinued and the advice of a veterinarian should be sought. Avoid use in any dehydrated, hypovolaemic or hypotensive animal, as there is a potential risk of increased renal toxicity.

Other NSAIDS, diuretics, anticoagulants, aminoglycoside antibiotics and substances with high protein binding may compete for binding and thus lead to toxic effects. Flexicam must not be administered in conjunction with other NSAIDS or glucocorticosteroids.

Pre-treatment with anti-inflammatory substances may result in additional or increased adverse effects and accordingly a treatment-free period with such drugs should be observed for at least 24 hours before commencement of treatment. The treatment-free period, however, should take into account the pharmacokinetic properties of the products used previously.

In the case of overdosage symptomatic treatment should be initiated.

People with known hypersensitivity to NSAIDS should avoid contact with the veterinary medicinal product.

In case of accidental ingestion, seek medical advice immediately and show this package insert or the label to the physician.

## **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste material derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

18.03.2010

Detailed information on this veterinary medicinal product is available on the website of the European Medicines Agency (European Medicines Agency) <http://www.ema.europa.eu/>.

**15. OTHER INFORMATION**

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.



**PACKAGE LEAFLET**  
**Flexicam 5 mg/ml solution for injection for dogs and cats**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

**Marketing authorisation holder:**

Dechra Veterinary Products A/S  
Mekuvej 9  
7171 Uldum  
Denmark

**Manufacturer for the batch release:**

Accord Healthcare Limited  
Sage House 1<sup>st</sup> Floor, 319 Pinner Road  
North Harrow, Middlesex HA1 4HF  
United Kingdom

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Flexicam 5 mg/ml solution for injection for dogs and cats.  
Meloxicam.

**3. STATEMENT OF THE ACTIVE SUBSTANCE AND OTHER INGREDIENTS**

1 ml of solution for injection contains 5 mg of meloxicam.  
Other substances: Ethanol anhydrous 150 mg/ml.

**4. INDICATIONS**

Dogs:

Alleviation of inflammation and pain in both acute and chronic musculo-skeletal disorders. Reduction of post-operative pain and inflammation following orthopaedic and soft tissue surgery.

Cats:

Reduction of post-operative pain after ovariohysterectomy and minor soft tissue surgery.

**5. CONTRAINDICATIONS**

Do not use in pregnant or lactating animals.

Do not use in animals suffering from gastrointestinal disorders such as irritation and haemorrhage, impaired hepatic, cardiac or renal function and haemorrhagic disorders.

Do not use in case of hypersensitivity to the active substance or to any of the excipients.

Do not use in animals less than 6 weeks of age nor in cats of less than 2 kg.

Do not use an oral follow-up therapy using meloxicam or other NSAIDs in cats, as no safe dosage for repeated oral administration has been established.

## **6. ADVERSE REACTIONS**

Typical adverse drug reactions of NSAIDs such as loss of appetite, vomiting, diarrhoea, faecal occult blood and apathy have occasionally been reported. In dogs, these side effects occur generally within the first treatment week and are in most cases transient and disappear following termination of the treatment but in very rare cases may be serious or fatal.

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

Dogs and cats.

## **8. DOSAGE FOR EACH SPECIES, ROUTES AND METHOD OF ADMINISTRATION**

Dogs: single administration of 0.2 mg meloxicam/kg body weight (i.e. 0.4 ml/10 kg).

Cats: single administration of 0.3 mg meloxicam/kg body weight (i.e. 0.06 ml/kg).

### Dogs:

Musculo-skeletal disorders: single subcutaneous injection.

Flexicam 1.5 mg/ml oral suspension may be used for continuation of treatment at a dosage of 0.1 mg meloxicam/kg body weight, 24 hours after administration of the injection.

Reduction of post-operative pain (over a period of 24 hours): single intravenous or subcutaneous injection before surgery, for example at the time of induction of anaesthesia.

### Cats:

Reduction of post-operative pain after ovariohysterectomy and minor soft tissue surgery: single subcutaneous injection before surgery, for example at the time of induction of anaesthesia.

Avoid introduction of contamination during use.

## **9. ADVICE ON CORRECT ADMINISTRATION**

Particular care should be taken with regard to the accuracy of dosing.

## **10. WITHDRAWAL PERIOD**

Not applicable.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the reach and sight of children.

This veterinary medicinal product does not require any special storage conditions.

Do not use after the expiry date stated on the carton and the vial after EXP.

Shelf-life after first opening the container: 28 days.

## **12. SPECIAL WARNINGS**

If side effects occur, treatment should be discontinued. Avoid use in any dehydrated, hypovolaemic or hypotensive animal, as there is a potential risk of increased renal toxicity.

Other NSAIDs, diuretics, anticoagulants, aminoglycoside antibiotics and substances with high protein binding may compete for binding and thus lead to toxic effects. Flexicam must not be administered in conjunction with other NSAIDs or glucocorticosteroids. Concurrent administration of potential nephrotoxic drugs should be avoided. In animals at anaesthetic risk (e.g. aged animals) intravenous or subcutaneous fluid therapy during anaesthesia should be taken into consideration. When anaesthesia and NSAID are concomitantly administered, a risk for renal function cannot be excluded.

Pre-treatment with anti-inflammatory substances may result in additional or increased adverse effects and accordingly a treatment-free period with such drugs should be observed for at least 24 hours before commencement of treatment. The treatment-free period, however, should take into account the pharmacokinetic properties of the products used previously.

In the case of overdosage symptomatic treatment should be initiated.

For post-operative pain relief in cats, safety has only been documented after thiopental/halothane anaesthesia.

The safety of the veterinary medicinal product has not been established during pregnancy and lactation.

### **User warnings**

Accidental self-injection may give rise to pain. People with known hypersensitivity to meloxicam should avoid contact with the veterinary medicinal product.

In case of accidental self-injection, seek medical advice immediately and show this package insert or the label to the physician.

## **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste material derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

## **14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

18.03.2010

Detailed information on this product is available on the website of the European Medicines Agency (European Medicines Agency) <http://www.ema.europa.eu/>.

## **15. OTHER INFORMATION**

### Pack sizes

Single 10 ml injection vial.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.