

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Zoonotic Influenza Vaccine Seqirus suspension for injection in pre-filled syringe.
Zoonotic influenza vaccine (H5N1) (surface antigen, inactivated, adjuvanted).

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Influenza virus surface antigens (haemagglutinin and neuraminidase)* of strain:

A/turkey/Turkey/1/2005 (H5N1)-like strain (NIBRG-23) (clade 2.2.1) 7.5 micrograms**
per 0.5 ml dose

* propagated in fertilised hens' eggs from healthy chicken flocks

** expressed in micrograms haemagglutinin (HA).

Adjuvant MF59C.1 containing:

squalene	9.75 milligrams per 0.5 ml
polysorbate 80	1.175 milligrams per 0.5 ml
sorbitan trioleate	1.175 milligrams per 0.5 ml
sodium citrate	0.66 milligrams per 0.5 ml
citric acid	0.04 milligrams per 0.5 ml

Excipient with known effect:

The vaccine contains 1.899 milligrams of sodium and 0.081 milligrams of potassium per 0.5 ml dose.

Zoonotic Influenza Vaccine Seqirus may contain trace residues of egg and chicken proteins, ovalbumin, kanamycin, neomycin sulphate, formaldehyde, hydrocortisone and cetyltrimethylammonium bromide which are used during the manufacturing process (see section 4.3).

For the full list of excipients see section 6.1.

3. PHARMACEUTICAL FORM

Suspension for injection (injection).

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Active immunisation against H5N1 subtype of Influenza A virus.

This indication is based on immunogenicity data from healthy subjects from the age of 18 years onwards following administration of two doses of the vaccine containing H5N1 subtype strain (see sections 4.4 and 5.1).

Zoonotic Influenza Vaccine Seqirus should be used in accordance with official recommendations.

4.2 Posology and method of administration

Posology

Adults and elderly (18 years of age and above)

One dose of 0.5 ml.

A second dose of 0.5 ml should be given after an interval of at least 3 weeks.

Zoonotic Influenza Vaccine Seqirus has been evaluated in healthy adults (18-60 years of age) and healthy elderly (over 60 years of age) following a 1, 22 day primary vaccination schedule, and booster vaccination (see sections 4.8 and 5.1).

There is limited experience in elderly over 70 years of age (see section 5.1).

In the event of an officially declared influenza pandemic due to A/H5N1 virus, individuals previously vaccinated with one or two doses of Zoonotic Influenza Vaccine Seqirus that contained HA antigen derived from a different clade of the same influenza subtype as the declared influenza pandemic strain may receive a single dose of Zoonotic Influenza Vaccine Seqirus instead of two doses that are required in previously unvaccinated individuals (see section 5.1).

Paediatric population

The safety and efficacy of Zoonotic Influenza Vaccine Seqirus in subjects under 18 years of age have not yet been established.

Currently available data in subjects aged 6 months to 18 years of age are described in section 5.1 but no recommendation on a posology can be made.

No data are available in children aged less than 6 months.

Method of administration

For intramuscular injection. The preferred injection site is the deltoid muscle.

The vaccine should under no circumstances be administered intravascularly or intradermally.

4.3 Contraindications

History of an anaphylactic (i.e. life-threatening) reaction to any of the constituents or trace residues (egg and chicken proteins, ovalbumin, kanamycin and neomycin sulphate, formaldehyde, hydrocortisone and cetyltrimethylammonium bromide) of this vaccine.

However, in a pandemic situation caused by the strain included in this vaccine, it may be appropriate to give this vaccine to individuals with a history of anaphylaxis as defined above, provided that facilities for resuscitation are immediately available in case of need.

4.4 Special warnings and precautions for use

Traceability

In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded.

General recommendations

Hypersensitivity and anaphylaxis

Caution is needed when administering this vaccine to individuals with a known hypersensitivity to the active substance, to any of the excipients listed in section 6.1 and to residues (eggs and chicken proteins, ovalbumin, kanamycin and neomycin sulphate, formaldehyde, hydrocortisone and cetyltrimethylammonium bromide).

As with all injectable vaccines, close observation for 15 min is recommended and appropriate medical treatment should always be readily available in case of a rare anaphylactic event following the administration of the vaccine.

Concurrent illness

Vaccination should be postponed in individuals with acute febrile illness or infection.

Thrombocytopenia and coagulation disorders

There are no data with Zoonotic Influenza Vaccine Seqirus using the subcutaneous route of administration. Therefore, healthcare providers need to assess the benefits and potential risks of administering the vaccine to individuals with thrombocytopenia or any bleeding disorder that would contraindicate intramuscular injection unless the potential benefit outweighs the risk of bleedings.

Limitations of vaccine effectiveness

There is no immune correlate of protection established for influenza A caused by H5N1 strain. Based on humoral immune responses after two doses of Zoonotic Influenza Vaccine Seqirus, a protective immune response may not be elicited in all vaccinees. In addition, antibody responses in patients with endogenous or iatrogenic immunosuppression may be insufficient to provide protection (see section 5.1).

No safety, immunogenicity or efficacy data to support interchangeability of Zoonotic Influenza Vaccine Seqirus with other H5N1 monovalent vaccines.

Anxiety-related reactions

Syncope (fainting) can occur following, or even before, any vaccination as a psychogenic response to the needle injection. This can be accompanied by several neurological signs such as transient visual disturbance, paraesthesia and tonic-clonic limb movements during recovery. It is important that procedures are in place to avoid injury from faints.

Cross-reactivity immunity

Some degree of cross-reactive immunity has been observed against H5N1 viruses of clades different to that of Zoonotic Influenza Vaccine Seqirus. However, the degree of protection that may be elicited to H5N1 strains of other clades is unknown (see section 5.1).

Excipients with known effect

Sodium

This vaccine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

Potassium

This vaccine contains potassium, less than 1 mmol (39 mg) per dose, that is to say essentially 'potassium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

Zoonotic Influenza Vaccine Seqirus may be co-administered with non-adjuvanted seasonal influenza vaccines, and immunisation should be carried out on separate limbs.

There are no data on co-administration of Zoonotic Influenza Vaccine Seqirus with other vaccines. If co-administration with another vaccine is considered, immunisation should be carried out on separate limbs. It should be noted that the adverse reactions may be intensified.

The immunological response may be diminished if the patient is undergoing immunosuppressant treatment.

4.6 Fertility, pregnancy and lactation

Pregnancy

Limited data were obtained from women who became pregnant during the course of clinical trials with Zoonotic Influenza Vaccine Seqirus or similar pandemic H1N1 vaccine adjuvanted with MF59C.1.

However, it is estimated that during the 2009 H1N1 pandemic more than 90 000 women were vaccinated during pregnancy with an H1N1 pandemic vaccine similar to Zoonotic Influenza Vaccine Seqirus which contains the same amount of adjuvant MF59C.1 as Zoonotic Influenza Vaccine Seqirus.

Post-marketing spontaneously reported adverse events and an interventional study do not suggest direct or indirect harmful effects of H1N1 vaccine exposure on pregnancy.

In addition, two large observational studies designed to assess the safety of H1N1 vaccine exposure in pregnancy showed no increase in the rates of gestational diabetes, preeclampsia, abortions, stillbirth, low birth weight, prematurity, neonatal deaths, and congenital malformations among almost 10 000 vaccinated pregnant women and their offspring compared with unvaccinated controls.

Since Zoonotic Influenza Vaccine Seqirus is expected not to be used in an emergency situation, its administration during pregnancy might be deferred as a precautionary approach.

Healthcare providers need to assess the benefit and potential risks of administering the vaccine to pregnant women taking into consideration official recommendations.

Breast-feeding

There are no data regarding the use of Zoonotic Influenza Vaccine Seqirus during breast-feeding. The potential benefits to the mother and risks to the infant should be considered before administering Zoonotic Influenza Vaccine Seqirus during breast-feeding.

Fertility

There are no data concerning human fertility. A study in rabbits did not indicate reproductive or developmental toxicity of Zoonotic Influenza Vaccine Seqirus (see section 5.3).

4.7 Effects on ability to drive and use machines

Some of the undesirable effects mentioned under section 4.8 may affect the ability to drive or operate machinery.

4.8 Undesirable effects

Summary of the safety profile

The incidence of adverse reactions has been evaluated in seven clinical trials in healthy subjects involving over 4300 adults and elderly receiving Zoonotic Influenza Vaccine (H5N1) (at least 7.5 µg HA, adjuvanted). There were 3872 subjects 18-60 years of age, 365 subjects 61-70 years of age, and 89 subjects greater than 70 years of age. The safety profile across clinical studies using Zoonotic Influenza Vaccine (H5N1) containing either the A/turkey/Turkey/1/2005 or the A/Vietnam/1194/2004 strain is comparable.

Consistent with the data observed by trial for solicited reactions, there was a general trend towards decreased reports of local reactions after the second vaccination compared with the first injection. Irrespective of antigen dose, almost all systemic reactions were reported on the day of vaccination (day 1) or during the 3 days immediately following.

Data on safety of a booster dose of Zoonotic Influenza Vaccine (H5N1) are limited to three trials (V87P1, V87P2 and V87P1E1) that included 116 adults (18-60 years) and 56 elderly subjects (≥ 61 years). No increase in reactions was reported when a booster dose was administered 6 months-18 months later, after the initial dosing series. A slight increase in reactions in adults was reported when a booster dose was administered 18 months after the initial dosing series. In the elderly, the reported reactions increased with the third booster dose only when compared with the second dose.

Tabulated list of adverse reactions

The adverse reaction rates reported after any vaccination doses (i.e. 1st, 2nd or booster) were similar and are listed according to the following MedDRA frequency convention and system organ class: Very common ($\geq 1/10$); Common ($\geq 1/100$ to $< 1/10$); Uncommon ($\geq 1/1\ 000$ to $< 1/100$); Rare ($\geq 1/10\ 000$ to $< 1/1\ 000$); Very rare ($< 1/10\ 000$).

Table 1. Adverse drug reactions

MedDRA System Organ class	Very common ($\geq 1/10$)		Common ($\geq 1/100$ to $< 1/10$)	Rare ($\geq 1/10\ 000$ to $< 1/1\ 000$)
Nervous system disorders	Headache			
Gastrointestinal disorders			Nausea	
Skin and subcutaneous tissue disorders			Sweating	
Musculoskeletal and connective tissue disorders	Myalgia		Arthralgia	
General disorders and administration site conditions	Injection site swelling, Injection site pain, Injection site induration, Injection site redness, Fatigue, Malaise, Chills		Injection site ecchymosis, Fever	Anaphylaxis

The majority of these side effects usually disappear within 1-2 days without treatment.

Clinical trials in special populations

Adverse reactions in special populations have been evaluated in two clinical trials, V87_25 and V87_26, involving adult (18-60 years) and elderly (≥ 61 years) subjects who were either healthy or with underlying medical conditions or immunosuppressive conditions.

Table 2. Adverse reactions in special populations

	Study V87_25	Study V87_26
--	--------------	--------------

	Medical conditions		Healthy		Immunocompromised		Healthy	
	Adults (20-60 years)*	Elderly (61-84 years)*	Adults (19-60 years)*	Elderly (61-79 years)*	Adults (20-60 years)*	Elderly (61-84 years)*	Adults (18-59 years)*	Elderly (61-91 years)*
No. of subjects	N=145	N=149	N=59	N=58	N=147	N=148	N=58	N=62

*actual age range of population enrolled

Across studies V87_25 and V87_26, the safety of Zoonotic Influenza Vaccine (H5N1) in healthy adult and elderly subjects was consistent with existing safety data from previous clinical trials. However, in immunocompromised subjects 18 through 60 years of age, slightly higher rates of nausea (13.0%) were reported. In addition, higher rates of arthralgia (up to 23.3%) were reported in both adult and elderly subjects, who were immunocompromised or with underlying medical conditions.

The following solicited adverse reactions were additionally collected in these two studies and reported with the following frequencies across subjects who received Zoonotic Influenza Vaccine (H5N1) irrespective of age or health status: diarrhoea (up to 11.9%), loss of appetite (up to 10.9%) and vomiting (up to 1.7%). In both studies, subjects with underlying medical and immunosuppressive conditions reported higher frequencies of diarrhoea, loss of appetite and vomiting compared to healthy subjects (irrespective of age).

Post-marketing surveillance

No post-marketing surveillance data are available for Zoonotic Influenza Vaccine Seqirus. The following adverse events were reported from post-marketing surveillance with an H1N1 pandemic vaccine similar to Zoonotic Influenza Vaccine Seqirus which contains the same amount of adjuvant MF59C.1 as Zoonotic Influenza Vaccine Seqirus, approved for use in children 6 months of age and above, adults and the elderly:

Blood and lymphatic system disorders

Lymphadenopathy.

Immune system disorders

Allergic reactions, anaphylaxis including dyspnoea, bronchospasm, laryngeal oedema, in rare cases leading to shock.

Nervous system disorders

Headache, dizziness, somnolence, syncope. Neurological disorders, such as neuralgia, paraesthesia, convulsions and neuritis.

Cardiac disorders

Palpitation, tachycardia.

Respiratory disorders

Cough.

Gastrointestinal disorders

Gastrointestinal disorders such as nausea, vomiting, abdominal pain and diarrhoea.

Skin and subcutaneous tissue disorders

Generalised skin reactions including pruritus, urticaria or non-specific rash, angioedema.

Musculoskeletal, connective tissue and bone disorders

Muscular weakness, pain in extremities.

General disorders and administration site conditions

Asthenia.

The following additional adverse events were reported from post-marketing surveillance with seasonal non-adjuvanted trivalent vaccines in all age groups and a seasonal trivalent MF59-adjuvanted subunit influenza vaccine approved for use in elderly individuals 65 years of age and older:

Blood and lymphatic system disorders

Thrombocytopenia (in some cases reversible platelet counts less than 5 000 mm³).

Nervous system disorders

Neurological disorders, such as encephalomyelitis, and Guillain Barré syndrome.

Vascular disorders

Vasculitis which may be associated with transient renal involvement.

Skin and subcutaneous tissue disorders

Erythema multiforme.

General disorders and administration site conditions

Extensive swelling of injected limb lasting more than one week, injection-site cellulitis-like reaction (some cases of swelling, pain, and redness extending more than 10 cm and lasting more than 1 week).

Paediatric population

The incidence of Zoonotic Influenza Vaccine (H5N1) (A/Vietnam/1194/2004) adverse reactions has been evaluated in one clinical trial (V87P6) in children (6 months to 17 years old). Regardless of age, reactogenicity was higher after the first dose than after the second vaccination. Reactogenicity after the third dose, administered 12 months following the first dose, was higher than after both first and second doses. The percentages of subjects reporting local reactions were higher in the older age groups, mainly due to the higher reports for pain. In toddlers, erythema and tenderness were the most commonly reported solicited local reactions; irritability and unusual crying were the most commonly reported solicited systemic reactions. In children and adolescents, pain was the most frequently

reported solicited local reaction, and fatigue and headache were the most commonly reported solicited systemic reactions. Across all ages, low percentages of subjects reported fever.

Table 3. Incidence of adverse reactions in paediatric study V87P6

	Injection 1	Injection 2	Injection 3
	Zoonotic Influenza Vaccine (H5N1)		
Toddlers (6-<36 months)	N=145	N=138	N=124
Any	76%	68%	80%
Local	47%	46%	60%
Systemic	59%	51%	54%
Fever $\geq 38^{\circ}\text{C}$ ($\geq 40^{\circ}\text{C}$)	0%	0%	0%
Any Other Adverse Event	54%	49%	35%
Children (3-<9 years)	N=96	N=93	N=85
Any	72%	68%	79%
Local	66%	58%	74%
Systemic	32%	33%	45%
Fever $\geq 38^{\circ}\text{C}$ ($\geq 40^{\circ}\text{C}$)	4%	2%	6%
Any Other Adverse Event	36%	31%	19%
Adolescents (9-<18 years)	N=93	N=91	N=83
Any	91%	82%	89%
Local	81%	70%	81%
Systemic	69%	52%	69%
Fever $\geq 38^{\circ}\text{C}$ ($\geq 40^{\circ}\text{C}$)	0%	1%	2%
Any Other Adverse Event	30%	27%	22%

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

4.9 Overdose

No case of overdose has been reported.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Influenza vaccine ATC Code J07BB02.

Clinical efficacy and safety

Clinical trials with Zoonotic Influenza Vaccine (H5N1) have been conducted with either A/Vietnam/1194/2004 (H5N1) (clade 1) or A/turkey/Turkey/1/2005 (H5NI) vaccine strain (clade 2.2.1).

Immune response to Zoonotic Influenza Vaccine (H5N1) A/Vietnam/1194/2004 (H5N1) and A/turkey/Turkey/1/2005 (H5N1)

Adults (18-60 years)

A phase II clinical trial (V87P1) was conducted with Zoonotic Influenza Vaccine (H5N1) (A/Vietnam/1194/2004) in 312 healthy adults. Two doses of Zoonotic Influenza Vaccine (H5N1) were administered three weeks apart to 156 healthy adults. Immunogenicity was assessed in 149 subjects. In phase III clinical trial (V87P13) 2693 adult subjects were enrolled and 2566 received two doses of Zoonotic Influenza Vaccine (H5N1) (A/Vietnam/1194/2004) administered three weeks apart. Immunogenicity was assessed in a subset (N=197) of subjects. In a third clinical trial (V87P11) 194 adult subjects were enrolled and received two doses of Zoonotic Influenza Vaccine (H5N1) (A/turkey/Turkey/1/2005) administered three weeks apart. Immunogenicity was assessed in 182 subjects.

The seroprotection rate*, seroconversion rate** and the seroconversion factor*** for anti-HA antibody to H5N1 A/Vietnam/1194/2004 and to H5N1 A/turkey/Turkey/1/2005 in the adults measured by SRH assay is reported below:

Table 4. Seroprotection rate, seroconversion rate and seroconversion factor for anti HA antibody to H5N1 A/Vietnam/1194/2004 and to H5N1 A/turkey/Turkey/1/2005

Anti-HA antibody (SRH)	Study V87P1 A/Vietnam/1194/2004 21 days after 2 nd dose N=149	Study V87P13 A/Vietnam/1194/2004 21 days after 2 nd dose N=197	Study V87P11 A/turkey/Turkey/1/2005 21 days after 2 nd dose N=182
Seroprotection rate (95%CI)*	85% (79-91)	91% (87-95)	91% (85-94)
Seroconversion rate (95%CI)**	85% (78-90)	78% (72-84)	85% (79-90)
Seroconversion factor (95%CI)***	7.74 (6.6-9.07)	4.03 (3.54-4.59)	6 (5.2-6.93)

Anti-HA antibody (SRH)	Study V87P13 A/Vietnam/1194/2004 21 days after 2 nd dose N=69	Study V87P13 A/Vietnam/1194/2004 21 days after 2 nd dose N=128	-
Baseline Serostatus	< 4 mm ²	≥ 4 mm ²	-
Seroprotection rate (95%CI)*	87% (77-94)	94% (88-97)	-
Seroconversion rate (95%CI)**	87% (77-94)	73% (65-81)	-
Seroconversion factor (95%CI)***	8.87 (7.09-11)	2.71 (2.38-3.08)	-

* Seroprotection: SRH area ≥ 25 mm²

** Seroconversion was defined as an SRH area ≥25 mm² for subjects who were seronegative at baseline (Day 1 SRH area ≤4 mm²) or a significant (at least 50%) increase in SRH area for subjects who were seropositive at baseline (Day 1 SRH area >4 mm²)

*** geometric mean ratios (GMRs) of SRH

MicroNeutralisation (MN) results against homologous A/Vietnam/1194/2004) indicate a seroprotection and seroconversion rate ranging from 67% (60-74) to 85% (78-90) and 65% (58-72) to 83% (77-89), respectively. Immune response to vaccination assessed by MN assay is in line with results obtained with SRH.

In Study V87P11 MN results against homologous A/turkey/Turkey/1/2005 indicate a seroprotection and seroconversion rate of 85% (79-90) and 93% (89-96), respectively. Immune response to vaccination assessed by MN assay is in line with results obtained with SRH.

Persistence of antibodies after primary vaccination in this population was assessed by hemagglutination inhibition (HI), SRH, and MN assays. Compared to the antibody levels obtained at day 43 after completion of primary vaccination schedules, antibody levels at day 202 were reduced by 1/5 to 1/2 from their prior levels.

Elderly (≥ 61 years)

Table 5. The seroprotection rate*, seroconversion rate and the seroconversion factor*** for anti-HA antibody to H5N1 (A/Vietnam/1194/2004 and to A/ turkey/Turkey/1/2005) in subjects aged 61 years and older^a measured by SRH assay assessed in three clinical studies:**

Anti-HA antibody (SRH)	Study V87P1 A/Vietnam/1194/2004 21 days after 2 nd dose N=84 ^a	Study V87P13 A/Vietnam/1194/2004 21 days after 2 nd dose N=210 ^b	Study V87P11 A/turkey/Turkey/1/2005 21 days after 2 nd dose N=132 ^c
Seroprotection rate (95%CI)*	80% (70-88)	82% (76-87)	82% (74-88)
Seroconversion rate (95%CI)**	70% (59-80)	63% (56-69)	70% (61-77)
Seroconversion factor (95%CI)***	4.96 (3.87-6.37)	2.9 (2.53-3.31)	3.97 (3.36-4.69)

Anti-HA antibody (SRH)	Study V87P13 A/Vietnam/1194/2004 21 days after 2 nd dose N=66	Study V87P13 A/Vietnam/1194/2004 21 days after 2 nd dose N=143
Baseline Serostatus	< 4 mm ²	≥ 4 mm ²
Seroprotection rate (95%CI)*	82% (70-90)	82% (75-88)
Seroconversion rate (95%CI)**	82% (70-90)	54% (45-62)
Seroconversion factor (95%CI)***	8.58 (6.57-11)	1.91 (1.72-2.12)

^a Limited number of subjects were above 70 years of age; N=123

^b Ages 62-88 years; ^c Ages 61-68 years; ^d Ages 61-89 years

* Seroprotection: SRH area ≥ 25 mm²

** Seroconversion was defined as an SRH area ≥ 25 mm² for subjects who were seronegative at baseline (Day 1 SRH area ≤ 4 mm²) or a significant (at least 50%) increase in SRH area for subjects who were seropositive at baseline (Day 1 SRH area > 4 mm²)

*** GMRs of SRH

MN results against homologous A/Vietnam/1194/2004 indicate a seroprotection and seroconversion rate ranging from 57% (50-64) to 79% (68-87) and 55% (48-62) to 58% (47-69), respectively. MN results, similar to SRH results, demonstrated strong immune response after completion of priming vaccination series in a population of elderly subjects.

In Study V87P11, MN results against homologous A/turkey/Turkey/1/2005 indicate a seroprotection and seroconversion rate of 68% (59-75) and 81% (74-87), respectively. Immune response to vaccination assessed by MN assay is similar to SRH results.

Based on data obtained from trials V87P1, V87P11 and V87_13, persistence of antibodies after primary vaccination in elderly subjects as assessed by HI, SRH, and MN tests reduced from 1/2 to 1/5th of their post-vaccination level at day 202 as compared to day 43 after completion of primary schedules. Up to 50% (N=33) of the elderly subjects aged 62 to 88 years immunised with Zoonotic Influenza Vaccine (H5N1) in trial V87P1 were seroprotected at six months.

A third (booster) dose of Zoonotic Influenza Vaccine (H5N1) was administered 6 months onwards after the primary vaccination. Results are shown by SRH.

Table 6. The seroprotection rate*, seroconversion rate and the seroconversion factor*** for anti-HA antibody to H5N1 A/Vietnam/1194/2004 measured by SRH assays**

	Study V87P1 Adults booster after 2 nd dose	Study V87P2 Adults booster after 2 nd dose	Study V87P1 Elderly booster after 2 nd dose
SRH	N=71	N=13	N=38
Seroprotection rate (95%CI)*	89% (79-95)	85% (55-98)	84% (69-94)
Seroconversion rate (95%CI)**	83% (72-91)	69% (39-91)	63% (46-78)
Seroconversion factor (95%CI)***	5.96 (4.72-7.53)	2.49 (1.56-3.98)	5.15 (3.46-7.66)

* Seroprotection: SRH area ≥ 25 mm²

** Seroconversion was defined as an SRH area ≥ 25 mm² for subjects who were seronegative at baseline (Day 1 SRH area ≤ 4 mm²) or a significant (at least 50%) increase in SRH area for subjects who were seropositive at baseline (Day 1 SRH area >4 mm²)

*** GMRs of SRH

Cross reactivity data in adults

Cross-reactive immune response elicited by A/Vietnam/1194/2004 against A/turkey/Turkey/1/2005 and A/Indonesia/5/2005

Some heterologous immune response against A/turkey/Turkey/1/2005 (NIBRG23; clade 2.2.1) and A/Indonesia/5/2005 (clade 2.1) was detectable both after the second and third vaccinations, indicating cross-reactivity of the clade 1 vaccine against clade 2 strains.

Table 7. Seroprotection rate*, seroconversion rate and the seroconversion factor*** for anti-HA antibodies to H5N1 A/turkey/Turkey/1/2005 after the 2nd dose in adults 18-60 years of age, measured by SRH and HI assays:**

	Anti-HA antibody	Study V87P12 21 days after 2 nd dose N=60	Study V87P3 21 days after 2 nd dose N=30	Study V87P13 21 days after 2 nd dose N=197
SRH	Seroprotection rate (95%CI)*	65% (52-77)	90% (73-98)	59% (52-66)
	Seroconversion rate (95%CI)**	65% (52-77)	86% (68-96)	49% (42-56)
	Seroconversion factor(95%CI)***	4.51 (3.63-5.61)	7.67 (6.09-9.67)	2.37 (2.1-2.67)
		N=60	N=30	N=197
HI	Seroprotection rate (95%CI) ^o	28% (17-41)	24% (10-44)	23% (18-30)
	Seroconversion rate (95%CI) ^o	28% (17-41)	21% (8-40)	19% (14-25)
	Seroconversion factor (95%CI) ^{oo}	2.3 (1.67-3.16)	1.98 (1.22-3.21)	1.92 (1.64-2.25)

* Seroprotection: SRH area ≥ 25 mm²

** Seroconversion was defined as an SRH area ≥ 25 mm² for subjects who were seronegative at baseline (Day 1 SRH area ≤ 4 mm²) or a significant (at least 50%) increase in SRH area for subjects who were seropositive at baseline (Day 1 SRH area >4 mm²)

*** GMRs of SRH

^o measured by HI assay ≥ 40

°° GMRs of HI

MN results for the three clinical studies in the Table above revealed a seroprotection rate and seroconversion rate against A/turkey/Turkey/2005 ranging from 10% (2-27) to 39% (32-46) and 10% (2-27) to 36% (29-43) respectively. MN results yielded a GMR against A/turkey/Turkey/2005 ranging from 1.59 to 2.95.

Cross-reactive immune response elicited by A/turkey/Turkey/1/2005 against A/Indonesia/5/2005 and A/Vietnam/1194/2004

Heterologous immune response against A/Indonesia/5/2005 (clade 2.1) was detectable in study V87P11 after the second vaccination, indicating cross-reactivity of the clade 2.2.1 vaccine against clade 2.1 strains.

Table 8. Seroprotection rate*, seroconversion rate and the seroconversion factor*** for anti-HA antibodies to H5N1 A/ Indonesia/5/2005 and A/Vietnam/1194/2004 after the 2nd dose in adults (18-60 years) and elderly (≥61years), measured by SRH and HI assays**

Anti-HA antibody		V87P11 Adults (18-60 years) N=182		V87P11 Elderly (≥61-89 years) ^a N=132	
		A/Indonesia/ 5/2005	A/Vietnam/ 1194/2004	A/Indonesia/ 5/2005	A/Vietnam/ 1194/2004
SRH	Seroprotection rate (95%CI)*	83 (77-88)	62 (54-69)	61 52-69	45 (37-54)
	Seroconversion rate (95%CI)*	79 (72-85)	60 (53-68)	64 (56-73)	44 (35-53)
	Seroconversion factor (95%CI)**	6.24 (5.44-7.16)	4.45 (3.85-5.14)	3.87 (3.31-4.53)	3.03 (2.56-3.58)
		N=194		N=148	
HI	Seroprotection rate (95%CI) [°]	50 (43-57)	47 (40-55)	34 (26-42)	39 (31-48)
	Seroconversion rate (95%CI) [°]	49 (42-56)	44 (37-51)	32 (25-41)	34 (26-42)
	Seroconversion factor (95%CI) °°	4.71 (3.74-5.93)	4.25 (3.36-5.37)	2.69 (2.18-3.32)	2.8 (2.2-3.55)

^a Actual age range of population enrolled

* Seroprotection: SRH area ≥25 mm²

** Seroconversion was defined as an SRH area ≥25 mm² for subjects who were seronegative at baseline (Day 1 SRH area ≤4 mm²) or a significant (at least 50%) increase in SRH area for subjects who were seropositive at baseline (Day 1 SRH area >4 mm²)

*** GMRs of SRH

[°] measured by HI assay ≥ 40

°° GMRs of HI

MN results for A/Indonesia/5/2005 revealed a seroprotection rate of 38% (31-45) in adults (18-60 years) and 14% (8-20) in elderly (≥61 years); a seroconversion rate of 58% (50-65) in adults and 30% (23-38) in elderly and finally a GMR of 4.67 (3.95-5.56) in adults and 2.19 (1.86-2.58) in elderly.

MN results for A/Vietnam/1194/2004 revealed a seroprotection rate of 10% (6-16) in adults (18-60 years) and 6% (3-11) in elderly (≥61 years); a seroconversion rate of 19% (13-25) in adults and 7% (4-13) in elderly and finally a GMR of 1.86 (1.63-2.12) in adults and 1.33 (1.17-1.51) in elderly.

Long term booster immune memory

A single vaccination with Zoonotic Influenza Vaccine (H5N1) (A/Vietnam/1194/2004) induced high and rapid serological response in subjects primed 6-8 years previously with two doses of a different surrogate H5vaccine, having same formulation as Zoonotic Influenza Vaccine (H5N1) but using the strain H5N3.

In a phase I clinical trial (V87P3) adult subjects aged 18-65 years primed 6-8 years previously with 2 doses of MF59-adjuvanted H5N3 vaccine/A/Duck/Singapore/97, were administered 2 booster doses of Zoonotic Influenza Vaccine (H5N1) (A/Vietnam/1194/2004). SRH results after the first dose, that mimic pre-pandemic priming plus single heterologous booster dose, revealed seroprotection and seroconversion rates of 100% (74- 100) and an 18-fold increase in SRH area (GMR).

Alternative vaccination schedules

In a clinical trial evaluating 4 different vaccination schedules in 240 subjects 18 to 60 years of age, where the second dose occurred either 1, 2, 3 or 6 weeks after the first Zoonotic Influenza Vaccine (H5N1) (A/Vietnam/1194/2004) dose, all vaccine schedule groups after 3 weeks from the 2nd vaccination achieved high levels of antibodies as evaluated with SRH. SRH seroprotection rates ranged from 86% to 98%, seroconversion rates from 64% to 90%, and GMR ranged from 2.92 to 4.57. The magnitude of immune response was lower in the group who received the 2nd dose 1 week later and higher in the groups with longer interval schedules.

Subjects with underlying medical or immunosuppressive conditions

Immunogenicity of Zoonotic Influenza Vaccine (H5N1) (A/turkey/Turkey/1/2005) in adults (18 to 60 years) and elderly (≥ 61 years) subjects with underlying medical conditions (Study V87_25) or immunosuppressive conditions (mainly HIV-infected subjects) (Study V87_26) in comparison to healthy adults (18-60 years) and elderly (≥ 61 years), was evaluated in two randomised, phase III controlled clinical trials (with a seasonal trivalent inactivated MF59-adjuvanted subunit influenza vaccine approved for use in elderly subjects 65 years of age and older as a comparator). In trial V87_25 and V87_26, 96 and 67 subjects, respectively, were over the age of 70 years. In both trials, immunogenicity of Zoonotic Influenza Vaccine (H5N1) was shown by HI, SRH and MN assays following both the first and second dose.

Table 9. Geometric mean area*, seroprotection rate*, seroconversion rate* and the seroconversion factor for anti-HA antibody to H5N1 A/turkey/Turkey/1/2005 measured by SRH assays 21 days after the 2nd dose**

Study V87_25				
	Adults (20- 60 years) ^a	Adults (19- 60 years) ^a	Elderly (61-84 years) ^a	Elderly (61-79 years) ^a
Anti-HA antibody (SRH)	Medical Conditions N=140	Healthy N=57	Medical Conditions N=143	Healthy N=57
Geometric Mean Area (95%CI)*	31.07 (27.43-35.19)	58.02 (48.74-69.06)	29.34 (26.07-33.01)	27.78 (22.57-34.18)
Seroprotection rate (95%CI)*	65.00 (56.5-72.9)	89.47 (78.5-96)	58.74 (50.2-66.9)	57.89 (44.1-70.9)
Seroconversion rate (95%CI)*	72.86 (64.7-80)	98.25 (90.6-99.96)	64.34 (55.9-72.2)	66.67 (52.9-78.6)
Seroconversion factor (95%CI)**	3.33 (2.94-3.77)	6.58 (5.53-7.83)	2.37 (2.10-2.66)	2.96 (2.41-3.64)
Study V87_26				
	Adults (20- 60 years) ^a	Adults (18-59 years) ^a	Elderly (61-84 years) ^a	Elderly (61-91 years) ^a

Anti-HA antibody (SRH)	Immuno-compromised N=143	Healthy N=57	Immuno-compromised N=139	Healthy N=62
Geometric Mean Area (95%CI)*	26.50 (22.49-31.22)	48.58 (40.01-58.99)	26.85 (23.01-31.33)	23.91 (18.89-30.26)
Seroprotection rate (95%CI)*	60.84 (52.3-68.9)	87.72 (76.3-94.9)	58.99 (50.3-67.3)	53.23 (40.1-66)
Seroconversion rate (95%CI)*	61.54 (53-69.5)	89.47 (78.5-96)	64.75 (56.2-72.7)	56.45 (43.3-69)
Seroconversion factor (95%CI)**	3.16 (2.69-3.73)	7.10 (5.85-8.62)	3.15 (2.70-3.68)	2.83 (2.24-3.58)

^a actual age range of population enrolled

* measured by SRH assay seroprotection: SRH area ≥ 25 mm², seroconversion: SRH area ≥ 25 mm² for subjects with a baseline SRH area ≤ 4 mm² or a minimum 50% increase in SRH area for subjects with >4 mm².

** geometric mean ratios of SRH

HI results for the two clinical studies revealed lower values than those reported in previous studies. Seroconversion rates against homologous A/turkey/Turkey/1/2005 ranged from 37.50% to 43.10% in healthy adults, and from 19.18% to 26.47% in adults with immunosuppressive or underlying medical conditions, respectively; seroconversion rates ranged from 21.43% to 30.65% in healthy elderly subjects, and from 24.49% to 27.86% in elderly subjects with immunosuppressive or underlying medical conditions. Similar trends were observed for seroprotection rates in both studies.

MN results against homologous A/turkey/Turkey/1/2005 indicate a seroconversion rate of 66.67% in healthy adults, and ranging from 33.57% to 54.14% in adults with immunosuppressive or underlying medical conditions, respectively; seroconversion rates ranged from 24.39% to 29.03% in healthy elderly subjects, and from 31.65% to 39.42% in elderly subjects with immunosuppressive or underlying medical conditions. Similar trends were observed for seroprotection rates in both studies.

In both studies V87_25 and V87_26, the lower levels of antibodies (as measured by HI, SRH and MN assays) and reduced seroprotection rates in adults and elderly (≥ 61 years old) subjects with underlying medical or immunosuppressive conditions, suggest that Zoonotic Influenza Vaccine (H5N1) may not elicit the same level of protection against A/H5N1 strain as compared to healthy adults (see section 4.4). These studies provided limited immunogenicity data in subjects with some underlying medical (in particular, renal impairment and peripheral cardiovascular disease) and immunosuppressive conditions (in particular, transplant recipients and patients under cancer treatment). In these trials, lower levels of antibodies and reduced seroprotection rates against homologous H5N1 A/turkey/Turkey/1/2005 were also measured in healthy elderly subjects, as compared to healthy adults, though previous studies showed induction of sufficiently immunogenic responses against H5N1 strains (see above for information on elderly).

Available data in paediatric population

A clinical trial (V87P6) was conducted with Zoonotic Influenza Vaccine (H5N1) (A/Vietnam/1194/2004) in 471 children from 6 months to 17 years of age. Two doses of Zoonotic Influenza Vaccine (H5N1) were administered three weeks apart and a third dose 12 months following the first dose. After 3 weeks from the 2nd vaccination (day 43) all age groups (i.e. 6-35 months, 3-8 years and 9-17 years) achieved high levels of antibodies to

(A/Vietnam/1194/2004) as evaluated with SRH and HI assays as presented in table below. In this trial no vaccine related SAEs were observed.

Table 10. Seroprotection rate*, seroconversion rate and the seroconversion factor*** for anti-HA antibodies to H5N1 A/Vietnam/1194/2004 measured by HI and SRH assays 21 days after the 2nd dose in paediatric subjects 6 months to 18 years of age**

		Toddlers (6-<36 months)	Children (3-<9 years)	Adolescents (9-<18 years)
		N=134	N=91	N=89
HI	% SP (95% CI) Day 43	97% (92-99)	97% (91-99)	89% (80-94)
	GMR Day 43 to Day 1	129 (109-151)	117 (97-142)	67 (51-88)
	% SC (95% CI) Day 43	97% (92-99)	97% (91-99)	89% (80-94)
SRH		N=133	N=91	N=90
	% SP (95% CI) Day 43	100% (97-100)	100% (96-100)	100% (96-100)
	GMR (95% CI) Day 43 to Day 1	16 (14-18)	15 (13-17)	14 (12-16)
	% SC (95% CI) Day 43	98% (95-100)	100% (96-100)	99% (94-100)

SP= seroprotection

SC= seroconversion

MN results against a A/Vietnam/1194/2004 indicate a seroprotection rate of 99% (95%CI: 94-100), a seroconversion rate ranging from 97% (95%CI: 91-99) to 99% (95%CI: 96-100) and a GMR ranging from 29 (95%CI: 25-35) to 50 (95%CI: 44-58).

The European Medicines Agency has deferred the obligation to submit the results of studies with Zoonotic Influenza Vaccine (H5N1) in one or more subsets of the paediatric population in active immunisation against H5N1 subtype of Influenza A virus. See section 4.2 for information on paediatric use.

Information from non-clinical studies

Efficacy against challenge with virus homologous and heterologous to vaccine strains was evaluated in the ferret model. Zoonotic Influenza Vaccine (H5N1) (A/Vietnam/1194/2004) and Zoonotic Influenza Vaccine (H5N1) (A/turkey/Turkey/1/2005) were tested. Animals received one or two doses of vaccine containing 3.75 or 7.5 micrograms of antigen, followed by challenge with a lethal dose of A/Vietnam/1203/04 virus.

All animals receiving 2 doses of Zoonotic Influenza Vaccine (H5N1) were protected, and 94% of animals receiving a single dose of Zoonotic Influenza Vaccine (H5N1) were protected. 87% of animals challenged with virus heterologous to the vaccine strain after 2 doses of vaccine were protected, and a single dose of heterologous vaccine protected 56% of the animals.

In a similar study, intranasal challenge was delayed until approximately 4 months after the second dose of vaccine was administered. In this study 100% of animals were protected against homologous challenge, and 81% of animals were protected against heterologous challenge. Vaccination protected animals from lethal challenge even when HI antibody titres were low or undetectable.

Efficacy against challenge with the heterologous virus A/Indonesia/5/2005 was also tested. Ferrets received one or two doses of vaccine (A/Vietnam/1194/2004). Two doses of vaccine protected 92% of animals, and a single dose of vaccine protected 50% of animals against challenge with the A/Indonesia/5/2005 virus. Lung damage was reduced in vaccinated groups. Viral shedding and viral titres in lungs were also reduced, suggesting that vaccination may reduce the risk of viral transmission.

5.2 Pharmacokinetic properties

Not applicable.

5.3 Preclinical safety data

Non-clinical data obtained with Zoonotic Influenza Vaccine (H5N1) and with seasonal influenza vaccine containing MF59C.1 adjuvant reveal no special hazard for humans based on conventional studies of repeated dose toxicity, local tolerance, female fertility, and reproductive and developmental toxicity (through the end of the lactation period).

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sodium chloride
Potassium chloride (E508)
Potassium dihydrogen phosphate (E340)
Disodium phosphate dihydrate (E339)
Magnesium chloride hexahydrate (E511)
Calcium chloride dihydrate (E509)

Water for injections.

For the adjuvant, see section 2

6.2 Incompatibilities

In the absence of compatibility trials, this medicinal product must not be mixed with other medicinal products.

6.3 Shelf life

2 years.

6.4 Special precautions for storage

Store in a refrigerator (2°C - 8°C).
Do not freeze.
Store in the original package in order to protect from light.

6.5 Nature and contents of container

0.5 ml in pre-filled syringe (type I glass) with plunger-stopper (bromo-butyl rubber).

Packs of 1 or 10 pre-filled syringes.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

Gently shake before use.

After shaking, the normal appearance of Zoonotic Influenza Vaccine Seqirus is a milky-white suspension.

Visually inspect the suspension prior to administration. In case of any particles and/or abnormal appearance, the vaccine should be discarded.

Any unused vaccine and waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Seqirus S.r.l.
Via del Pozzo 3/A, S. Martino
53035 Monteriggioni (SI)
Italy.

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/23/1761/001-002

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation:

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu/>.

ANNEX II

- A. MANUFACTURERS OF THE BIOLOGICAL ACTIVE SUBSTANCE AND MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

A. MANUFACTURERS OF THE BIOLOGICAL ACTIVE SUBSTANCE AND MANUFACTURER RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturers of the biological active substance

Seqirus Vaccines Ltd
Gaskill Road, Speke, Liverpool
L24 9GR
UK

Name and address of the manufacturer responsible for batch release

Seqirus Netherlands B.V.
Paasheuvelweg 28
1105BJ Amsterdam
Netherlands

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to medical prescription.

- **Official batch release**

In accordance with Article 114 of Directive 2001/83/EC, the official batch release will be taken by a state laboratory or a laboratory designated for that purpose.

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

- **Periodic Safety Update Reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

PSUR submission when Zoonotic Influenza Vaccine Seqirus is used during an influenza pandemic:

During a pandemic situation, the annual frequency of PSUR submission may not be adequate for the safety monitoring of a pandemic vaccine for which high levels of exposure are expected within a short period of time. Such situation requires rapid notification of safety information that may have the greatest implications for benefit-risk balance in a pandemic. Prompt analysis of cumulative safety information, in light of extent of exposure, will be crucial for regulatory decisions and protection of the population to be vaccinated.

In consequence, as soon as the pandemic is declared and the zoonotic vaccine is used, the Marketing Authorisation Holder (MAH) shall submit more frequent simplified PSURs with a periodicity defined in the Risk Management Plan (RMP).

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

- **Risk Management Plan (RMP)**

The Marketing Authorization Holder shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

If the dates for submission of a PSUR and the update of a RMP coincide, they can be submitted at the same time.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARDBOARD BOX

1. NAME OF THE MEDICINAL PRODUCT

Zoonotic Influenza Vaccine Seqirus Suspension for injection in pre-filled syringe.
Zoonotic influenza vaccine (H5N1) (surface antigen, inactivated, adjuvanted)

2. STATEMENT OF ACTIVE SUBSTANCE

One dose of 0.5 ml contains: Influenza virus surface antigens (haemagglutinin and neuraminidase), propagated in fertilised hens' eggs from healthy chicken flocks, of strain:

A/turkey/Turkey/1/2005 (H5N1)-like strain (NIBRG-23) (clade 2.2.1) 7.5 micrograms haemagglutinin

Adjuvant: MF59C.1 oil in water emulsion containing squalene, as the oil phase, stabilised with polysorbate 80, sorbitan trioleate, sodium citrate and citric acid.

3. LIST OF EXCIPIENTS

Sodium chloride
Potassium chloride (E508)
Potassium dihydrogen phosphate (E340)
Disodium phosphate dihydrate (E339)
Magnesium chloride hexahydrate (E511)
Calcium chloride dihydrate (E 509)
Water for injections.

4. PHARMACEUTICAL FORM AND CONTENTS

Suspension for injection.

1 pre-filled syringe (0.5 ml)
10 pre-filled syringes (0.5 ml)

5. METHOD AND ROUTE OF ADMINISTRATION

To be administered intramuscularly into the deltoid muscle.

Read the package leaflet before use.

Gently shake before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP.:

9. SPECIAL STORAGE CONDITIONS

Store in a refrigerator.

Do not freeze.

Keep the pre-filled syringe in the outer carton in order to protect from light

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

Dispose of in accordance with local requirements.

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Seqirus S.r.l.

Via del Pozzo 3/A, S. Martino

53035 Monteriggioni (SI)

Italy.

12. MARKETING AUTHORISATION NUMBER

EU/1/23/1761/001 1 prefilled syringe

EU/1/23/1761/002 10 prefilled syringes

13. BATCH NUMBER

Lot:

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA
--

PC:

SN:

NN:

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

LABEL FOR SYRINGE

1. NAME OF THE MEDICINAL PRODUCT AND ROUTE OF ADMINISTRATION

Zoonotic Influenza Vaccine Seqirus injection
Zoonotic influenza vaccine H5N1
IM

2. METHOD OF ADMINISTRATION

Intramuscular use

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT

0.5 ml

6. OTHER

B. PACKAGE LEAFLET

Package leaflet: Information for the user

Zoonotic Influenza Vaccine Seqirus suspension for injection in pre-filled syringe Zoonotic influenza vaccine (H5N1) (surface antigen, inactivated, adjuvanted)

Read all of this leaflet carefully before you receive this vaccine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or nurse.
- If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Zoonotic Influenza Vaccine Seqirus is and what it is used for
2. What you need to know before you receive Zoonotic Influenza Vaccine Seqirus
3. How Zoonotic Influenza Vaccine Seqirus is given
4. Possible side effects
5. How to store Zoonotic Influenza Vaccine Seqirus
6. Contents of the pack and other information

1. What Zoonotic Influenza Vaccine Seqirus is and what it is used for

Zoonotic Influenza Vaccine Seqirus is a vaccine for use in adults from 18 onwards, intended to be given in the context of outbreaks of zoonotic influenza viruses (coming from birds) with pandemic potential to prevent flu caused by viruses similar to the vaccine strain reported in section 6.

Zoonotic influenza viruses occasionally infect humans, and can cause disease ranging from mild upper respiratory infection (fever and cough) to rapid progression to severe pneumonia, acute respiratory distress syndrome, shock and even death. Human infections are primarily caused by contact with infected animals, but do not spread easily between people.

Zoonotic Influenza Vaccine Seqirus is intended also to be given when there is anticipation of a possible pandemic due to the same or a similar strain.

When a person is given the vaccine, the immune system (the body's natural defence system) will produce its own protection (antibodies) against the disease. None of the ingredients in the vaccine can cause flu.

As with all vaccines, Zoonotic Influenza Vaccine Seqirus may not fully protect all persons who are vaccinated.

2. What you need to know before you are given Zoonotic Influenza Vaccine Seqirus

You should not receive Zoonotic Influenza Vaccine Seqirus:

- if you have previously had a sudden life-threatening allergic reaction to any ingredient of Zoonotic Influenza Vaccine Seqirus (listed in section 6) or to any of the substances that may be present in trace amounts as follows: egg and chicken protein, ovalbumin, formaldehyde, kanamycin and neomycin sulphate (antibiotics), hydrocortisone or cetyltrimethylammonium bromide (CTAB). Signs of an allergic reaction may include itchy skin rash, shortness of breath and swelling of the face or tongue. However, in a pandemic situation, it may be appropriate for you to be vaccinated with Zoonotic Influenza Vaccine Seqirus provided that appropriate medical treatment is immediately available in case of an allergic reaction.

Warnings and precautions

Talk to your doctor or nurse before having this vaccine

- if you have had any allergic reaction to any ingredient contained in the vaccine, to egg and chicken protein, ovalbumin, formaldehyde, kanamycin and neomycin sulphate (antibiotics), hydrocortisone or cetyltrimethylammonium bromide (CTAB) (see section 6. Further information).
- if you have a severe infection with fever (over 38°C). If this applies to you then your vaccination will usually be postponed until you are feeling better. A minor infection such as a cold should not be a problem, but your doctor or nurse should advise whether you could still be vaccinated with Zoonotic Influenza Vaccine Seqirus.
- if you are having a blood test to look for evidence of infection with certain viruses. In the first few weeks after vaccination with Zoonotic Influenza Vaccine Seqirus the results of these tests may not be correct. Tell the doctor requesting these tests that you have recently been given Zoonotic Influenza Vaccine Seqirus.
- In the presence of immune deficiencies Zoonotic Influenza Vaccine Seqirus may be administered but a protective immune response may not be elicited.

Please inform your doctor or nurse if you have a bleeding problem or bruise easily.

Fainting can occur following, or even before, any needle injection. Therefore tell the doctor or nurse if you fainted with a previous injection.

Zoonotic Influenza Vaccine Seqirus may not fully protect everyone who is vaccinated, especially elderly individuals and those with weakened immune systems, such as HIV patients, or those with underlying long term medical problems, such as diabetes, lung disease or heart problems. Tell your doctor if you have a weak immune system or an underlying long term medical problem.

In any of these cases, TELL YOUR DOCTOR OR NURSE, as vaccination may not be recommended, or may need to be delayed.

Other medicines and Zoonotic Influenza Vaccine Seqirus

Tell your doctor or nurse if you are taking or have recently taken or might take any other medicines, including medicines obtained without a prescription or have recently received any other vaccine.

Zoonotic Influenza Vaccine Seqirus can be given at the same time as non-adjuvanted seasonal influenza vaccines. There is no information on administration of Zoonotic Influenza Vaccine Seqirus with non-influenza vaccines. If administration of Zoonotic Influenza Vaccine Seqirus with other vaccines can not be avoided, the vaccines should be injected into separate limbs. In such cases, you should be aware that the side effects may be more intense.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or nurse for advice before receiving this vaccine. Your doctor needs to assess the benefits and potential risks of giving you the vaccine.

Driving and using machines

Some effects mentioned under section 4. "Possible side effects" may affect the ability to drive or use machines.

Zoonotic Influenza Vaccine Seqirus contains sodium and potassium.

This vaccine contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially 'sodium free'.

This vaccine contains potassium, less than 1 mmol (39 mg) per dose, i.e. essentially ‘potassium free’.

3. How Zoonotic Influenza Vaccine Seqirus given

Your doctor or nurse will administer the vaccine in accordance with official recommendations. The vaccine will be injected into the muscles of the upper arm (deltoid muscle). The vaccine should never be given into a vein.

Adults from 18 onwards:

One dose of 0.5 ml will be given. A second dose of 0.5 ml should be given after an interval of at least 3 weeks.

There is limited experience in elderly over 70 years of age.

Use in children

Children from 6 months to 17 years of age

There is limited experience in children between 6 months and 17 years of age. Vaccination is currently not recommended in this age group.

Children aged less than 6 months of age

Vaccination is currently not recommended in this age group.

If you have any further questions on the use of this medicinal product, ask your doctor or nurse.

4. Possible side effects

Like all medicines, Zoonotic Influenza Vaccine Seqirus can cause side effects, although not everybody gets them.

Allergic reactions may occur following vaccination, in rare cases leading to shock. Doctors are aware of this possibility and have emergency treatment available for use in such cases.

The side effects listed below have occurred with a vaccine similar to Zoonotic Influenza Vaccine Seqirus in clinical studies in adults, including the elderly:

Very common (affects more than 1 user in 10):

- Pain at the site of injection
- Hardening of the skin at the injection site
- Injection site redness
- Injection site swelling
- Aching muscles
- Headache
- Fatigue
- Generally feeling unwell
- Shivering

Common (affects 1 to 10 users in 100):

- Bruising of the skin at the injection site
- Aching joints
- Fever and nausea
- Sweating

Rare (affects 1 to 10 users in 10 000):

- Anaphylaxis (severe allergic reactions)

These side effects usually disappear within 1-2 days without treatment. If they persist, CONSULT YOUR DOCTOR.

Undesirable effects in patients with underlying long term medical problems such as diabetes, lung disease or heart problems and weakened immune systems (immunocompromised) such as HIV patients

Nausea, aching joints, diarrhoea and loss of appetite were reported very commonly in this population. In addition, vomiting was commonly reported.

Side effects from clinical study in children (6 months to 17 years of age)

General side effects reported very commonly in the 6 months to 35 months of age group were injection site redness, muscle ache, irritability and unusual crying. Very commonly reported reactions in the 36 months to 17 years of age group were pain, headache and fatigue.

Other side effects observed after routine use:

The side effects listed below have occurred in the days or weeks after vaccination with another H1N1 vaccine similar to Zoonotic Influenza Vaccine Seqirus. These side effects may occur with Zoonotic Influenza Vaccine Seqirus.

- Generalised skin reactions including
 - Itching
 - Urticaria (hives)
 - Rash or swelling of the skin and mucous membranes
 - Angioedema (abnormal swelling of the skin, usually around the eyes, lips, tongue, hands or feet, due to an allergic reaction)
- Disorders of the gut such as
 - Nausea
 - Vomiting
 - Abdominal pain
 - Diarrhoea
- Headache, dizziness, drowsiness, fainting
- Neurological disorders such as
 - Severe stabbing or throbbing pain along one or more nerves
 - Tingling
 - Fits
 - Neuritis (inflammation of nerves)
- Swollen lymph nodes, palpitations (irregular or forceful heart beat), tachycardia (faster than normal heart beat), weakness, pain in the extremities, cough and asthenia (unusual weakness).
- Allergic reactions possibly with shortness of breath, wheezing, swelling of the throat, or leading to a dangerous decrease of blood pressure, which, if untreated, may lead to shock. Doctors are aware of this possibility and have emergency treatment available for use in such cases.

Data in children and adolescents suggest a slight decrease in side effects after the second dose of the vaccine, with no increase in rates of fever.

In addition, the side effects listed below have occurred in the days or weeks after vaccination with vaccines given routinely every year to prevent seasonal flu. These side effects may occur with Zoonotic Influenza Vaccine Seqirus.

- Low blood platelet count which can result in bleeding or bruising
- Vasculitis (inflammation of the blood vessels which can cause skin rashes, joint pain and kidney problems)
- Erythema multiforme (type of allergic skin reaction that occurs in response to medications, infections, or illness)
- Neurological disorders such as encephalomyelitis (inflammation of the central nervous system), and a type of paralysis known as Guillain-Barré Syndrome
- Swelling, pain and redness at the injection site extending to more than 10 cm and lasting more than one week (Injection site cellulitis-like reaction)
- Extensive swelling of injected limb lasting more than one week

Reporting of side effects

If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Zoonotic Influenza Vaccine Seqirus

Keep this vaccine out of the sight and reach of children.

Do not use Zoonotic Influenza Vaccine Seqirus after the expiry date which is stated on the carton and the label. The expiry date refers to the last day of that month.

Store in a refrigerator (2°C - 8°C).

Do not freeze.

Store in the original package in order to protect from light.

Do not throw any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Zoonotic Influenza Vaccine Seqirus contains

- Active Substance:
Influenza virus surface antigens (haemagglutinin and neuraminidase)* of strain:

A/turkey/Turkey/1/2005 (H5N1)-like strain (NIBRG-23) (clade 2.2.1) 7.5 micrograms**
per 0.5 ml dose

* propagated in fertilised hens' eggs from healthy chicken flocks
** expressed in microgram haemagglutinin.
- Adjuvant MF59C.1:
The vaccine contains per 0.5 ml 9.75 mg squalene, 1.175 mg polysorbate 80, 1.175 mg sorbitan trioleate, 0.66 mg sodium citrate and 0.04 mg citric acid.
- Other ingredients:

The other ingredients are: sodium chloride, potassium chloride, potassium dihydrogen phosphate, disodium phosphate dihydrate, magnesium chloride hexahydrate, calcium chloride dihydrate and water for injections.

What Zoonotic Influenza Vaccine Seqirus looks like and contents of the pack

Zoonotic Influenza Vaccine Seqirus is a suspension for injection in a pre-filled syringe. The suspension is a milky-white liquid. It is provided in a ready-to-use pre-filled syringe, containing a single dose of 0.5 ml for injection.

Packs of 1 or 10 pre filled syringes.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

Seqirus S.r.l.
Via del Pozzo 3/A, S. Martino
53035 Monteriggioni (SI)
Italy.

Manufacturer

Seqirus Netherlands B.V.
Paasheuvelweg 28
1105BJ Amsterdam
Netherlands

This leaflet was last revised in

Detailed information on this medicine is available on the European Medicines Agency web site:
<http://www.ema.europa.eu>.