



Scientific Committee on Health and Environmental Risks

SCHER

Opinion on  
risk assessment on indoor air quality



Approved by the SCHER at the 17<sup>th</sup> plenary of 29 May 2007 after public consultation

### About the Scientific Committees

Three independent non-food Scientific Committees provide the Commission with the scientific advice it needs when preparing policy and proposals relating to consumer safety, public health and the environment. The Committees also draw the Commission's attention to the new or emerging problems which may pose an actual or potential threat.

They are: the Scientific Committee on Consumer Products (SCCP), the Scientific Committee on Health and Environmental Risks (SCHER) and the Scientific Committee on Emerging and Newly-Identified Health Risks (SCENIHR) and are made up of external experts.

In addition, the Commission relies upon the work of the European Food Safety Authority (EFSA), the European Medicines Evaluation Agency (EMA), the European Centre for Disease prevention and Control (ECDC) and the European Chemicals Agency (ECHA).

### SCHER

Questions relating to examinations of the toxicity and ecotoxicity of chemicals, biochemicals and biological compound whose use may have harmful consequences for human health and the environment.

In particular, the Committee addresses questions related to new and existing chemicals, the restriction and marketing of dangerous substances, biocides, waste, environmental contaminants, plastic and other materials used for water pipe work (e.g. new organics substances), drinking water, indoor and ambient air quality. It addresses questions relating to human exposure to mixtures of chemicals, sensitisation and identification of endocrine disrupters.

### Scientific Committee members

Herman Autrup, Peter Calow, Wolfgang Dekant, Helmut Greim, Wojciech Hanke, Colin Janssen, Bo Jansson, Hannu Komulainen, Ole Ladefoged, Jan Linders, Inge Mangelsdorf, Marco Nuti, Jerzy Sokal, Anne Steenhout, Jose Tarazona, Emanuela Testai, Marco Vighi, Matti Viluksela

### Contact:

European Commission  
Health & Consumer Protection DG  
Directorate C: Public Health and Risk Assessment  
Unit C7 - Risk Assessment  
Office: B232 B-1040 Brussels  
[Sanco-Sc8-Secretariat@ec.europa.eu](mailto:Sanco-Sc8-Secretariat@ec.europa.eu)

© European Commission 2007

The opinions of the Scientific Committees present the views of the independent scientists who are members of the committees. They do not necessarily reflect the views of the European Commission. The opinions are published by the European Commission in their original language only.

[http://ec.europa.eu/health/ph\\_risk/risk\\_en.htm](http://ec.europa.eu/health/ph_risk/risk_en.htm)

## **ACKNOWLEDGMENTS**

Members of the working group are acknowledged for their valuable contribution to this opinion:

Prof. Wojciech Hanke, Nofer Institute of Occupational Medicine, Poland

Prof. Bo Jansson, Stockholm University, Sweden

Dr. Hannu Komulainen, National Public Health Institute, Finland

Prof. Ole Ladefoged, Institute of Food Safety and Nutrition, Denmark

Dr. Inge Mangelsdorf, Fraunhofer Institute for Toxicology and Experimental Medicine, Germany

Prof. Anne Steenhout, Université Libre de Bruxelles, Belgium

Dr. E. Testai, Istituto Superiore di Sanità, Italy

### External experts

Prof. Gunnar Damgård Nielsen, National Research Centre for Working Environment, Denmark

Prof. Philip Harteman, Faculté de Médecine, Université de Nancy, France

Dr. Aino Nevalainen, National Public Health Institute, Finland

Keywords: SCHER, scientific opinion, indoor air, air pollution, household chemicals, building, dampness

Opinion to be cited as:

SCHER Opinion on risk assessment on indoor air quality, 29 May 2007

## TABLE OF CONTENTS

ACKNOWLEDGMENTS .....	3
1. Background .....	5
2. Terms of reference .....	6
3. Opinion .....	6
3.1. Question 1.....	7
3.1.1. Risk Assessment for indoor air .....	7
3.1.1.1. Hazard identification.....	7
3.1.1.2. Dose-response assessment .....	9
3.1.1.3. Exposure assessment .....	9
3.1.1.4. Risk characterisation .....	11
3.1.2. Vulnerable groups .....	11
3.1.3. Mixture/combined effects .....	12
3.2. Question 2.....	14
3.2.1. Adequacy of current information .....	14
3.2.2. Data requirements and gaps of knowledge.....	15
3.2.3. Existing measurement standards.....	16
3.3. Question 4.....	16
3.3.1. Concerns in relation to specific chemical compounds .....	17
3.3.2. Concerns in relation to household-chemicals and other products (e.g. decorating materials, cleaners, furnishing, etc.).....	19
3.3.3. Concerns in relation to building dampness/moisture and microbial growth .....	20
4. Conclusions .....	21
5. General conclusions and recommendations.....	22
6. References .....	23
7. List of abbreviations.....	33

## 1. BACKGROUND

Air quality is one of the major environmental health concerns for Europe. The main goal of the Community policy on air pollution is to achieve levels of air quality that do not result in unacceptable risks to human health.

Until now, much progress has been made in Europe in tackling outdoor air pollutants with Community legislation on emission sources available since 1970 and on air quality standards since 1980. This legislation has been continuously updated. The most recent EU quality standards are defined within the Air quality framework directive from 1996<sup>1</sup> and subsequent directives<sup>2</sup>. The air quality directives require Member States to set up and maintain a system for assessing outdoor air quality and to draw up action plans to reach the objectives of the EC directives. Limit values have been adopted for outdoor air concentration of the most common pollutants such as sulphur dioxide, lead and particulate matter<sup>3</sup>. In 2001, the Commission also launched an air quality programme Clean Air for Europe (CAFÉ)<sup>4</sup> to prepare a long term strategy on air pollution, which was adopted by the Commission on 21 September 2005.

In 2002, four priority areas to be tackled with urgent actions were identified in the EU's Sixth Community Environment Action Programme<sup>5</sup>. One of the priority areas is the Environment and health and quality of life. In 2003, the European Commission adopted a new Strategy on Environment and Health<sup>6</sup> with the overall aim to reduce diseases caused by environmental factors in Europe. Among others, air quality is one of the main identified problems of environmental pollution related to health problems such as respiratory diseases, asthma and allergies.

This strategy was followed by the EU Action Plan on Environmental and Health<sup>7</sup>. In this plan, the "action 12" intends to develop work on improving indoor air quality.

These initiatives recognize the importance of a complete integrated strategy on air pollution which considers not only the outdoor air but also includes possible solutions concerning indoor air pollutants. As reported in the communication on a European Environment and Health Strategy, a broad stakeholder involvement is also foreseen which includes a consultative group on "Environment and Health". The Commission will use this group to consult on analyses of environment and health data, and risk management measures. The analyses in question will rely in part on the work of the SCHER to identify missing links in existing monitoring systems.

Indoor exposures to air pollutants may occur in both private and public indoor environments such as homes, offices, schools, and transport systems.

---

<sup>1</sup> OJ L296/55 21 November 1996

<sup>2</sup> OJ L163/41, 29 June 1999, OJ L313/12, 13 December 2000, OJ L67/14 9 March 2002, L23/3 26 January 2005

<sup>3</sup> OJ L163/41 29 June 1999

<sup>4</sup> COM (2001) 245 final

<sup>5</sup> OJ L 242 10/9/2002

<sup>6</sup> COM (2003) 338 final

<sup>7</sup> COM (2004) 416 final

Most indoor air pollutants consist of chemicals released, for example, the use of cleaning products, air fresheners, pesticides and emissions from furniture and construction materials, heating and cooking. In addition, outdoor sources may contribute to indoor air pollution. Aspects such as thermal insulation and ventilation rates may also play a role. Microbiological contaminants which may induce allergies and asthma also require consideration as indoor air pollutants. Examples of potential serious effects include respiratory disorders, including asthma and cancer.

## **2. TERMS OF REFERENCE**

To provide a basis for assessment of risks to human health from indoor air quality, and a sound scientific basis for the development and implementation of policies, the SCHER is asked:

- 1.** To identify a Risk Assessment Strategy to support policy on indoor air quality. The Committee is particularly asked to take into account potentially vulnerable groups of population such as children, pregnant women and elderly. The committee is also asked to consider the practicality of a risk assessment which takes into account on combined exposure and cumulative effects of specific air pollutants.
- 2.** To identify the adequacy of current information and data requirements for filling-in gaps on aspects such as exposure/effect and dose/response relationships, existing measurement standards and gaps in knowledge which will help to guide further research and monitoring efforts.
- 3.** To consider risks associated with the use of air fresheners:

SCHER has given a separate opinion on this point (SCHER 2006).

- 4.** To identify potential areas of concern in relation to:
  - specific chemical compounds taking into account the recent outcome of the INDEX report prepared by DG JRC
  - household – chemicals and other products (e.g. decorating materials, cleaners, furnishings, etc.)
  - building dampness/moisture and microbial growth (moulds, bacteria)

## **3. OPINION**

Indoor air constitutes a complex case for risk assessment and management due to a wide variety of pollutants, exposure levels, different possible health outcomes, differences in sensitivity of the population, cultural habits, way of living, building stock and climate across Europe.

Global trends such as climate change and rise of energy costs may have important effects on indoor air quality through for example, extreme weather conditions and need to additional thermal insulation and decreased ventilation. These may lead to too high or too low indoor temperatures or to dampness problems.

Possible health risks of indoor air pollution and aspects of risk assessment have been addressed recently in European and international working groups and projects (WHO 1997, ECA 2000, INDEX 2005, California EPA 2005).

The opinion is aimed to cover indoor environments where the general public may be exposed to pollutants, such as private homes and public buildings e.g., schools, day care centres, offices and places of leisure. Transport vehicles (vehicle compartments) are specific indoor environments. The same risk assessment principles can be applied for them but the means of risk management may differ. Industrial exposures, including professional cleaning, in indoor environments are excluded because they do not represent the exposure of the general public; and specific exposure limits for contaminants are established. In addition, the opinion does not cover active smoking but environmental tobacco smoke (ETS) is included.

The SCHER emphasises that the opinion covers only risk assessment principles which are recommended to be applied in indoor environments and does not cover risk management, the practical means to decrease/prevent adverse health effects.

The opinion draft was on public comments through the internet and was finalized after consideration of the comments received.

### **3.1. Question 1**

**To provide a basis for assessment of risks to human health from indoor air quality, and a sound scientific basis for the development and implementation of policies, the SCHER is asked to identify a Risk Assessment Strategy to support policy on indoor air quality. The Committee is particularly asked to take into account potentially vulnerable groups of population such as children, pregnant women and elderly. The committee is also asked to consider the practicality of a risk assessment which takes into account on combined exposure and cumulative effects of specific air pollutants.**

#### **Answer to Question 1**

##### **3.1.1. Risk Assessment for indoor air**

The SCHER recommends to use the basic paradigm for toxicological risk assessment, (Commission Directive 93/67/EEC; Council Regulation (EEC) 793/93; TGD, 2003) in support of the indoor air quality policy. It takes into account all necessary elements in health risk assessment: the hazards, their dose-response, exposure and results in science-based risk characterisation (Commission Regulation (EC) No. 1488/94; Technical Guidance Document on Risk Assessment). Where possible, the margin of safety/exposure should be defined or the exposure should be compared with relevant health based guideline values for risk characterisation. In the case this is not possible a scientific based hazard characterisation should be attempted.

Where diseases, risk of diseases, symptoms or complaints are obviously related to unacceptable indoor conditions (e.g. serious over crowdedness, extreme heat, lack of ventilation, excessive dirtiness, excessive microbial growth and dampness) and combustion products, no further detailed risk assessment may be needed and risk management may directly be performed.

##### **3.1.1.1. Hazard identification**

A number of factors in the indoor environment can affect well-being and health. The main factors related to indoor air are

- Chemicals for intended use or unintentional emissions from different sources
- Radon
- Particles
- Microbes

- Pets and pests
- Humidity
- Ventilation
- Temperature

### *Chemicals*

Data requirement for hazard identification of chemicals is described in the TGD. Information on the acute and sub-acute health effects can be obtained from toxicological data-bases and sources. However, information on possible health effects after long term exposures (carcinogenic, mutagenic, reproductive effects) is not available for many of the chemicals reported to be present in indoor air. The relevance of such data gaps needs to be discussed on case-by-case.

Sensory irritation of the eyes and upper airways is a specific effect often related to exposure to indoor air pollutants (Alarie 1973; Alarie et al. 1998, Nielsen 1991; Doty et al. 2004) and needs particular attention. Sensory irritation may be induced by specific chemicals, but also by factors such as too low relative humidity of air.

Malodours are generally undesirable in indoor environments (Wolkoff et al. 2006a). Odours *per se* do not cause toxicological effects but may increase the reporting of symptoms (for example, headache, nausea, eye and throat irritation) (Wolkoff et al. 2006a). Other symptoms of odour exposures are hyperventilation or conditioned responses (Shusterman 2002).

Odours and sensory irritation and conditioned responses due to these challenges are common causes of complaints regarding poor indoor air quality.

### *Radon*

Radon in indoor air has been associated with lung cancer (WHO 1998, Darby et al., 2005). Radon gas diffuses through soil into residences in areas where bedrock contains in excess uranium. Radon evaporates also from household water into air upon warming. High radon concentrations have been measured indoors locally in several countries.

### *Particles*

Particles (PM<sub>10</sub>, PM<sub>2.5</sub>, fine particles, ultrafine particles) in ambient air have been associated with adverse health effects, including respiratory and cardiovascular effects (WHO 2003, WHO 2005b). Particles from outdoor air may contribute to particle load in indoor air, but there are also indoor sources such as combustion (Lam et al., 2006), cooking, and particles may be formed by reactions between ozone and some VOCs (Wainman et al., 2000, Sarwar et al., 2004, Afsari et al., 2005). Man-made nanoparticles are increasingly used in consumer products but their impact as indoor air pollutants is not yet known and finally assessed.

### *Microbes*

Microbial agents may play a role in the development of asthma and allergic airway diseases. Many fungi contain allergens, and sensitization is possible by indoor exposure to fungi due to dampness and mould growth (see answer to Question 4c).

Additionally, virus infections may be transmitted by indoor air. In the first years of life, virus infections are common causes of wheezing. Some viruses are associated with an increase in asthma and allergy incidence (Schaub et al. 2006).

### *Pets and pests*

Pests, house dust mites and cockroaches are important sources of indoor allergens (aeroallergens) (D'Amato et al. 1998; Platts-Mills et al. 2000; Nielsen et al. 2002). Exposure may be different due to cultural habits, and cockroach and mouse allergens may be more important in inner cities (Eggleston 2001, Phipatanakul et al. 2000a; 2000b). Exposures to aeroallergens may cause sensitization with production of IgE antibodies. The IgE antibodies promote development of the allergic airway diseases, rhinitis and asthma (Beasley et al. 2000). The scientific literature on contact with pets and their effects on development of asthma is contradicting (for review e.g. Chan-Yeung and Becker 2006). However, allergic individuals should avoid exposures to allergens that elicit the allergic reactions.

### *Humidity*

Humidity in indoor air has an optimal window. Too low humidity results in increased reporting of skin symptoms (dryness, rash), eye irritation and nasal dryness (Reinikainen and Jaakkola, 2003, Wolkoff et al., 2006b). In excessive humidity, water condenses onto (cold) surfaces, causing water damage and mould problems. High humidity also favours the growth of dust mites.

### *Ventilation*

Ventilation is one of the most important contributing factors to indoor air quality. Low ventilation rates, as well as the carbon dioxide (CO<sub>2</sub>) concentrations in indoor air, have been associated with health and perceived air quality outcomes (Seppänen et al., 1999). Ventilation rate has been associated also with work performance in office work and academic performance of school children (Seppänen et al., 2005, Schaughnessy et al., 2006, Wargocki et al. 2007). Thermal insulation contributes to building tightness which usually increases the need of controlled ventilation.

### *Temperature*

Proper temperature is the basic indoor air requirement. Extreme indoor temperatures are a serious health hazard (Healy 2003, Kosatsky 2005) and too high or low temperatures perceived unpleasant. Too high temperature e.g. exacerbates the effects of insufficient humidity (Reinikainen and Jaakkola, 2001).

### **3.1.1.2. Dose-response assessment**

Information about exposure-response relationship is crucial for risk assessment. Regarding local effects such as irritation, air concentrations of pollutants are directly relevant for effects assessment. For evaluation of possible systemic effects, air concentrations of pollutants may be transformed to internal doses using established values for breathing volumes and alveolar retention. When data from occupational exposures are used, they must be adjusted for differences in exposure duration (general population 20-21 hours indoors vs. a work shift of 8 hours, and a longer exposure period of the life span) and the limitations of the occupational observations need to be considered for the exposed population; occupational exposures, in general, do not include infants, children and the elderly. Both acute and chronic effects should be taken into account in the dose-response assessment (INDEX 2005).

Biomarkers, when available, may be used to establish the dose-response relationships.

### **3.1.1.3. Exposure assessment**

More than 900 different organic compounds have been detected in indoor air (SCALE, 2004a) in addition to fine and ultrafine particles and biological material (microbes, allergens). Concentrations of pollutants in indoor air are determined by a number of factors

including type and emission rates from sources, ventilation rate, adsorption/absorption of compounds on/in materials (sinks and possible secondary sources).

Differences in cultural habits of people throughout Europe may result in large qualitative and quantitative differences in indoor air quality. The results of the EXPOLIS-INDEX study concerning exposure to European indoor environments in Athens, Basel, Grenoble, Milan, Helsinki, Oxford and Prague (e.g. VOCs and PM<sub>2.5</sub>) showed larger within city variation in exposure than average between city variation (EXPOLIS-INDEX, 2004). The health consequence of these variations is insufficiently understood.

Emissions of chemicals may occur from building materials (e.g. Bornehag et al., 2005a), cooking activities (e.g. Afshari et al., 2005), cleaning activities (Nazaroff and Weschler 2004, Singer et al., 2006), heating and combustion of biomass fuels in general (e.g. the opinion on air fresheners, SCHER 2006) but the exposure patterns are not sufficiently known.

Models have been developed to predict the emissions and distribution patterns of pollutants. Such models are essential for the development indoor exposure and risk assessment (ECA 2006, Kephelopoulos et al., 2007). However, no comprehensive general model has yet emerged and been validated.

Exposures to indoor air pollutants may occur directly by inhalation, or indirectly by ingestion of e.g., dust, while volatile compounds such as formaldehyde and benzene are mainly present in the gas phase. Less volatile substances are also to some extent bound to particles and dust, and exposure via those media may contribute to the total exposure. Many semivolatiles such as phthalates, flame retardants, PAHs, chlorophenols, pesticides, organotins and metals may adsorb to house dust (Butte and Heinzow, 2002). Particles may abrade from materials containing the chemicals of interest, e.g. PVC particles containing phthalates. House dust forms a long-term sink, may be regularly resuspended and represent a secondary source for pollutants. The particle size of house dust is, however, typically large and only a fraction of the dust is respirable (Butte and Heinzow, 2002, Maertens et al., 2004). Ingestion is likely the main exposure route for house dust (Butte and Heinzow, 2002) in small children due to licking and biting on articles and "hand to mouth" pattern. Small children spend a considerable part of the time on the floor and may thus be more exposed to resuspended dust than adults. The particle/dust pathway has been shown to be an important exposure route e.g. for polybrominated diphenyl ethers (PBDEs). Exposure to dust may account for about 14% of the external exposure of those substances for adults, but over 90% for children in the age of 1 to 4 years (Shoeib et al, 2004, Stapleton et al, 2005; Wilford et al, 2005). However, there is insufficient information in general to what extent the exposure to dust actually leads to uptake of pollutants in children.

In the absence of exactly measured data, the exposure should be estimated using validated models and information on local conditions. Modelling of chemical substances has the longest tradition (WHO 2005a). At present, there are no reliable models for exposure to indoor air microbial exposure.

SCHER states that it is important to evaluate effects of inhalation exposure but also underlines the importance to assess the contribution of indoor air exposures to total exposure to a chemical by all routes (ingestion, inhalation and dermal) for assessment of systemic effects to obtain an overall risk assessment and to estimate the contribution of indoor air pollution. The frequent focus on VOCs and other selected compounds in measurement campaigns may neglect a possible impact of compounds with high toxicity present in low concentrations or of compounds which are difficult to detect by the procedures applied. Moreover, the SCHER recommends to use realistic exposure scenarios and to avoid conclusions on exposures based on the content of potentially toxic chemicals in

materials such as wallpaper or furnishings. Biomonitoring will make a very valuable contribution to the exposure assessment.

#### **3.1.1.4. Risk characterisation**

For some pollutants occurring in indoor air, international (WHO 2000) and national (EPA, OEHHA, ATSDR, UBA, Health Canada, cited e.g. in the INDEX-report, INDEX 2005) health-based exposure limit/guidelines exist. They may not be aimed specifically for indoor air but may be used considering the specific exposure situation.

In risk characterisation, the whole exposure range should be included, not only average or median exposures. The variations in exposures may expand over several orders of magnitude. Analysis of EXPOLIS study data on VOCs has shown that the group at the upper 95% end of the distribution may get exposed significantly more than the median group (Edwards et al., 2005). On the other hand, the most sensitive subgroups may react at notably lower exposures. Accordingly, the range of plausible risk estimates (not only the central estimate) is useful. The precision and the uncertainty related to risk estimates should also be given in assessments.

Indication of the association between exposure and health effects may be difficult when the primary causing factor/agent is not known though the association is evident. This is typical for indoor air problems caused by microbes. For those cases, exclusion of other contributing factors may strengthen the association.

#### **3.1.2. Vulnerable groups**

For the opinion, vulnerable groups are represented by children, pregnant women, elderly persons over 65 years of age, and persons suffering from asthma or other respiratory diseases, and cardiovascular diseases. For some pollutants (e.g. microbes) other health compromises (immunodeficiency) may render people more vulnerable. Genetic traits, nutritional status and life-style factors may also contribute (TNO and RIVM, 2006).

The assumption of different susceptibility of vulnerable groups (children, pregnant, elderly) to pollutants is based on age-dependent differences in physiology and toxicokinetics and varying responses due to existing diseases and genetic factors (e.g. IPCS 1993, Tamburini et al., 2002, Pediatrics 2004).

Vulnerability to chemical toxicity after birth may be highest during the first 6 months (Scheuplein et al, 2002, Ginsberg et al., 2004) and continue for years before maturation. However, children may also be less sensitive and tolerate higher doses of chemical substances than adults, depending on the age and the compound (Schneider et al., 2002, Dourson et al., 2002). Higher exposure due to specific exposure patterns (e.g. hand-to-mouth activity in children) may increase the risk for children. Air pollutants may affect adversely foetal and infant lung development, cause post-neonatal infant mortality due to respiratory diseases, cause cough and bronchitis and aggravate asthma (WHO 2005b). The effect on lung function during development has been observed below the NOEL of effects of single air pollutants in adults, suggesting a higher susceptibility of children. The causative pollutants have not been identified but the association to adverse effects has been detected most consistently with outdoor particulate matter (PM), nitrogen dioxide and ozone (WHO 2005b). Studies addressing specifically the indoor environment, where the concentrations are different, are so far limited. Children's higher susceptibility is known for lead and environmental tobacco smoke (Tamburini et al., 2002, DiFranza et al., 2004); some concern has been expressed also for organophosphate pesticides (Grandjean and Landrigan, 2006, see also bullet 4a. in the opinion).

Altered physiology and toxicokinetics (e.g. reduced renal clearance) make elderly people potentially more sensitive (IPCS 1993) due to reduced capacity for elimination. However,

elderly people may also be less sensitive to some effects (Kjaergaard et al. 1992, Shusterman et al., 2003) including nasal (Schusterman et al., 2003) and eye irritation (Kjaergaard et al., 1992) indicating that aging may also decrease the susceptibility.

People suffering from cardiovascular diseases are more vulnerable to particles (WHO 2003, Dominici et al., 2006) and persons suffering from asthma and other respiratory diseases are more susceptible to several air pollutants (WHO 2004a, 2005b). For example, sensory irritation may occur at lower exposure level in persons with allergic rhinitis (Shusterman et al., 2003, WHO 2005b).

Currently within Europe and other industrialized countries as well as international bodies, vulnerable groups are considered on a case-by-case basis in general. However, a major gap in identification of vulnerable groups is the lack of data, but several ongoing activities are attempting to extend e.g. reproductive toxicity testing to better address immunological and neurological differences. Additionally, physiological-based pharmacokinetic models can offer insight in intraindividual variability in pharmacokinetics (TNO and RIVM 2006).

The SCHER recommends that the science-based health risk assessment always addresses vulnerable groups; the impact on the evaluation should be based on a case-by-case approach, considering compound's toxicological features and exposure. The SCHER also reminds that the Margin-of-Safety approach already includes specific safety factors to account for intraspecies differences.

### **3.1.3. Mixture/combined effects**

The SCHER was asked to consider the practicality of a risk assessment which takes into account combined exposure and cumulative effects of specific air pollutants. Within this opinion, the SCHER interprets the combined effects as the total effects caused by exposure to all chemical and biological (allergens and microbes) stressors present in indoor air.

At present the (quantitative) risk characterization must mostly be done on a single chemical basis because there are seldom relevant data and established methods to evaluate mixture effects. In indoor environment, exposures are always to complex mixtures of substances from different sources which may jointly contribute to the toxic effects. Due to the complexity of indoor air pollution and its variability with time, estimation of risk associated with exposure of the complex mixture as such and then generalize the obtained results is rarely feasible. This approach has only been used in few cases, when sensory irritation was the end-point (Hempel-Jørgensen et al., 1999, Nielsen et al., 2007b).

The majority of toxicology data refer to exposures to single chemicals. Such data can be used directly if chemicals in a mixture act independently with different endpoints, i.e. the effect of each component of the mixture is not influenced by the presence of the other components ('dissimilar joint action'). The single chemical approach is supported by the results of some studies, which indicate that interactions were unlikely to occur at environmentally relevant concentrations (which often are well below the NOAEL values). Interactive effects giving rise to possible health concern have been reported, starting from concentrations around the LOAEL (Cassee et al, 1998).

However, the single chemical approach is not applicable when the components affect each others response. Such combined effects may be additive (a 'similar joint action'; similar endpoints, similar mechanism of action and /or toxicokinetics properties) or there may be interactions (antagonistic or synergistic effects). Combined effects have been demonstrated e.g. by mixtures of pesticides when potentially harmful effects were observed at concentrations of each single component below or approaching the individual NOAEL value (Cavieses, 2002).

Models to evaluate toxicity of chemical mixtures have focused primarily on quantifying dose addition, as in the EPA assessment of health risk at hazardous waste sites (US EPA, 1986). The methods for dose addition which have been most frequently used are the Relative Potency Factor (RPF), the Toxic Equivalent Factor (TEF) and the Hazard Index (HI). When extensive mechanistic information is not available, the HI is the preferred approach. HI is a dimensionless figure, corresponding to the sum of the ratios between the exposure level and the reference dose (RfD) of each component, representing the relative potency. When HI for the whole mixture is equal to 1, it is supposed that the exposure correspond to the RfD of the mixture; when values are higher than 1, potential health concerns should be considered. HI derivation can be revised in order to be able to incorporate interaction data, when available, introducing a weight of evidence evaluation and an adjustment factor for the relative potency of each component (US EPA, 2001b).

With respect to indoor air pollution, a number of studies have dealt with the combined effects of indoor air pollutants, including effects of fine particles and gases in ambient air. The results have suggested that e.g. particles may behave as carriers for the toxicant into the lungs and that exposure to particulate matter may facilitate airway sensitisation in susceptible individuals (e.g. Hamada et al, 2000). NO<sub>2</sub> has increased the inflammatory effect of aeroallergen exposure in asthmatics (Barck et al., 2005) whereas formaldehyde has not modified the aeroallergen airway effect (Ezratty et al., 2007).

An additive approach has been considered useful for evaluation of mixtures of airborne sensory irritants above the threshold level (e.g. Cometto-Muñiz and Hernández 1990; Hempel-Jørgensen et al. 1999) and may be assumed as a first approximation of sensory irritation effects of mixtures based on animal and human studies (Nielsen et al., 2007b).

Some efforts have also been made to evaluate combined effects of a larger group of indoor air pollutants. The Committee of the Health Council in the Netherlands tentatively evaluated the health impact of volatile organic compounds (VOCs) from building materials (HCN, 2000). The Dutch committee considered the air quality guidelines developed by the WHO for outdoor air (WHO, 2000) and estimated the maximum tolerable pollution of indoor air by VOCs to be between 0.2 and 3.0 mg/m<sup>3</sup>, giving as recommended cumulative limit value of 0.2 mg/m<sup>3</sup> for VOCs not showing carcinogenic, reprotoxic or sensitizing properties. However, because the composition of total VOCs varies from place to place, this may only be used as a very general indicator of indoor air quality. Moreover, the compounds of highest concentrations are not necessarily those with offending effects in indoor air (Wolkoff and Nielsen, 2001).

The main problems encountered in applying the combined effect approach is that few data are available on interactions among more than two chemicals and they usually do not address issues of chronic toxicity at concentrations representative of actual human exposure. The use of PBPK and PBPD modelling may help (ATSDR, 2002, De Rosa et al, 2004). The use of mechanistic data derived from testing with binary mixtures may be extrapolated to more complex mixtures by means of PBPK models, as demonstrated with a mixture of benzene, toluene, ethylbenzene and xylene (BTEX) (Haddad, 2000) and may be very useful for the evaluation of metabolic interactions. In general, the issue of toxicity due to chemical mixture or multiple exposures suffers of the lack of both experimental data on the mode/mechanisms of actions and a generally accepted strategy for the related risk evaluation (McCarty and Borgert, 2006).

At present at the EU level there is not a general recommended approach to conduct the risk assessment for chemical mixtures or for combined effects due to concomitant exposure to different chemicals through different routes.

Altogether, the SCHER considers that the risk assessment which takes into account the combined exposure and cumulative effects of the pollutants in indoor environment is seldom possible. Mostly, there are not enough relevant data and the available methods may not fit

the case. However, the SCHER recommends that the possibility of combined effects is considered in the risk assessment and they are evaluated on a case-by-case approach. Interactions between chemicals and other factors such as microbes are insufficiently known to provide guidance.

### 3.2. Question 2

**To provide a basis for assessment of risks to human health from indoor air quality, and a sound scientific basis for the development and implementation of policies, the SCHER is asked to identify the adequacy of current information and data requirements for filling-in gaps on aspects such as exposure/effect and dose/response relationships, existing measurement standards and gaps in knowledge which will help to guide further research and monitoring efforts.**

#### Answer to Question 2

##### 3.2.1. Adequacy of current information

The SCHER notes that, taking into account all the variability and complexity in the indoor environment, the data for risk assessment are scarce and often insufficient. Recently, the THADE-project (THADE-report, 2004) has summarised several aspects on indoor air pollution in dwellings in Europe, including policies and actions taken in different countries. The evaluation indicates large differences between countries, in all aspects, and also lack of relevant data, both in general and member state specific. The PINCHE-project (PINCHE 2006) on children's health and environment prioritizes risk factors for environmental stressors and gives policy recommendations for research to air pollutants including indoor air.

In relation to exposure, information on concentrations of indoor air pollutants in Europe and information on determinants of personal exposure (e.g. EXPOLIS, 1999, German Environment Surveys (GerES I, GerES II, GerES III), German study on Indoor Factors and Genetics (INGA), the National survey of air pollutants in English homes) is available. These data give indications on the levels for some indoor pollutants and help to identify the compounds with highest concentrations and of highest concern.

Most of this information is on "classical" pollutants, such as carbon monoxide, nitrogen oxide, radon, asbestos, and organic compounds such as VOCs (INDEX 2005). Effects and risks for most of them are known to the extent that strategies to mitigate the problems can be created. But also new sources have emerged for "old" pollutants (e.g. VOCs in air fresheners, lead from candle wicks) and some of them (e.g. terpenes) may react to produce secondary products whose effects are poorly defined (Weschler et al., 2006, Wolkoff et al., 2006a).

Due to privacy of the indoor spaces (e.g. homes), enforceable indoor air standards are not preferred. Their systematic surveillance monitoring would be difficult. Instead, the SCHER supports the development of health-based guideline values and other guidance for key pollutants (as identified in this opinion) to help risk assessment and management. In this context, indicators other than concentrations of the pollutants (i.e. ventilation rate - and CO<sub>2</sub> as a related marker-, general cleanliness, signs of dampness) may also be applicable for monitoring.

At present, outdoor air quality is monitored for some pollutants (e.g., PM<sub>10</sub>, nitrogen oxide, ozone) but the data cannot predict the concentration in buildings, and replace measurement, because several local factors contribute to the access of pollutants indoors (e.g. tightness of the building). The variability in indoor levels has been shown for both organic compounds (e.g. Ilgen et al., 2001, Hodgson et al., 2003, Saarela et al., 2003, Gilbert et al. 2005) and particles (e.g. Lazaridis et al., 2006).

### **3.2.2. Data requirements and gaps of knowledge**

The SCHER has identified the following requirements for a more comprehensive and reliable health risk assessment of indoor air pollution. The needs range from broad and general to specific ones.

The data needs and gaps of knowledge are compiled into two groups, exposure assessment and health effects assessment. The SCHER considers that the items indicated by “++” should have the highest priority.

#### ***Data requirements and gaps in knowledge related primarily to identification and exposure to pollutants:***

##### *Need for compilation of existing data:*

- Comprehensive review of the existing data on the indoor air pollutants; definition of the major pollutants and their concentrations range in each Member State of EU, and set up of a pan European database (++). The database could be a part of a relevant existing information source. The process would compile the existing information on indoor pollutants, including allergens, as background for future work, and would facilitate the use of the data at an EU level, to identify differences among member states and data gaps. This information could drive both possible regular monitoring program and future research.
- Collection and systemization of practical experiences to establish evidence-based risk assessment approaches. Such examples would help to solve similar type problems and decrease the overlapping risk assessment.

##### ***Need for more research***

- Exposure patterns (short and long term in different environments) to indoor air pollutants, in quantitative terms, and identification of the most relevant exposure indicators (++). Description of typical exposure patterns would help to assess the typical levels and variability of exposure.
- Source apportionment of the pollutants in indoor environment, including ambient air, preferably in quantitative terms (++). Identification of the main sources would help their mitigation.
- Emissions of chemicals from consumer products (++). More data on levels of the emissions in realistic use situations is needed in view of the large part of population handling such products.
- Existing indoor source and transport/fate models should be identified, evaluated, validated and harmonized (+). Taxonomy of sources consistent data sharing should also be developed.
- Information on harmful emissions in water damaged buildings, including compounds from decomposing building material, contributing to toxicity (++). See also a detailed answer to question 4c.
- Evaluation of potentially harmful emissions from indoor combustion processes (e.g. halogenated dioxins). Low burning temperature may favour production of halogenated dioxins.

**Data requirements and gaps in knowledge related primarily to health effects of indoor air pollutants:**

*Need for more research:*

- Effects due to combined exposure to indoor air pollutants and objective methods for their evaluation, including development of validated modelling tools (++).
- Adverse health effects of microbes and bioaerosols present in indoor air, especially other than respiratory tract effects; responsible microbes and their components and toxins (++).
- Contribution of indoor air pollutants to childhood respiratory diseases (++).
- Exposure-effect-relationships especially in vulnerable groups (++).
- Effects and risks of products which emit indoor air pollutants that can react in indoor air (+). This is, for example, the case with terpenes that can react with ozone. The true role of such reaction products as indoor air pollutants is not clear.
- Possible effects and risks of man-made nanoparticles in indoor air (+).
- Contribution of coarse, fine and ultrafine particles from indoor sources to adverse health effects.
- Controlled clinical studies (including biochemical markers of effect) among persons suffering symptoms in water damaged buildings to clarify the associations and possibly to identify the most harmful microbes.

**3.2.3. Existing measurement standards**

There are some international measurement standards developed for indoor air quality both from CEN and ISO (often identical standards). Some of the standards developed for ambient air measurements can also be applied for the indoor environment, while methods for workplaces often are developed for higher concentrations of the substance. The SCHER does not see development of new measurement standards as a high priority, but recommends the validation and harmonization of the existing ones, in particular those concerned with indoor material emissions (ECA 2005) and biological agents. Development of passive samplers is in a very active phase and has to be followed, but the technique is not ready for standardisation.

**3.3. Question 4**

**To provide a basis for assessment of risks to human health from indoor air quality, and a sound scientific basis for the development and implementation of policies, the SCHER is asked to identify potential areas of concern in relation to a) specific chemical compounds taking into account the recent outcome of the INDEX report prepared by DG JRC, b) household – chemicals and other products (e.g. decorating materials, cleaners, furnishings, etc.) and c) building dampness/moisture and microbial growth (moulds, bacteria).**

**Answer to Question 4**

### 3.3.1. Concerns in relation to specific chemical compounds

Indoor environment contains a large number of different chemical compounds. Availability of data on exposures to specific chemicals, their toxicity and associated health risks are highly variable. Therefore, a priority ranking of chemicals and exposures which cause concern is difficult and uncertain. However, the SCHER considers that **formaldehyde, carbon monoxide, nitrogen dioxide, benzene, naphthalene, environmental tobacco smoke (ETS), radon, lead** and **organophosphate pesticides** are compounds of concern in indoor environment.

The INDEX project (INDEX 2005) has evaluated health risks of volatile chemicals in indoor air in the European population, as a stepwise procedure, and set up a list of compounds with highest concern on the basis of health impact criteria. After consideration of the quantity and quality of all the data available, 25 compounds were selected for a more detailed analysis and a detailed risk assessment was performed for 14 of them. The highest priority chemicals were **formaldehyde, carbon monoxide, nitrogen dioxide, benzene** and **naphthalene**. SCHER agrees that these are compounds of concern because they have caused adverse health effects as indoor pollutants or have a high potential to cause health effects. However, the concern is not similar in all over in Europe due to different exposure levels. For example, limited data on air fresheners indicate that burning of incense may produce abnormally high benzene and formaldehyde emissions in indoor air (Opinion of SCHER on air fresheners, SCHER 2006).

Though active smoking is excluded from this opinion, the SCHER reminds that tobacco smoking is the primary source of several emissions (benzene, fine and ultrafine particles) indoors and associated health effects. In adults, **ETS** has been associated e.g. with coronary heart disease, sensory irritation and exacerbation of respiratory symptoms, including asthma (IARC 2004). In children, the association with infant sudden death syndrome and middle ear infections and ETS has been observed (Tamburlini et al., 2002, DiFranza et al., 2004). The evidence clearly indicates that ETS requires risk management.

**Radon** in indoor air has been associated with lung cancer (WHO 1998). According to a recent analysis of European epidemiological studies (Darby et al., 2005) radon may be a common problem in Europe. Radon gas diffuses through soil into residences in areas where bedrock contains in excess uranium. Indoor radon concentrations can be decreased by technical means, even in existing buildings. Data on residential radon concentrations should be obtained by measurements in countries where such data do not yet exist and the associated health risk assessed.

Paint-related **lead** still exist in indoor environment in old houses in some EU countries though its use has been restricted or banned in indoor paints. Children are especially exposed through non-dietary ingestion of the dust. Exposure from other sources may be significant (TNO 2005)) and the evidence is increasing that already low level exposure of children to lead is harmful (e.g. Lanphear et al., 2005). Therefore, it is essential to evaluate, if the lead level in indoor environment is still a problem in EU countries. The existing data on lead should be compiled, and thereafter, a need for further research considered.

The indoor use of **organophosphate pesticides** for treatment of cracks and crevices (Byrne et al., 1998) or the use of insect strips (Weis et al. 1998) may lead to high exposures to these compounds by inhalation or ingestion due to accumulation on surfaces including children's toys (Hore et al., 2005) and house dust (Butte and Heinzow, 2002). This uptake may contribute considerably to the overall uptake of organophosphates by children (Gurunathan et al., 1998). The acute toxicity of organophosphate pesticides is well known (WHO 2004b): however, it is very unlikely that indoor levels can result in acute effects. Recently neuro-developmental effects, have been reported both in experimental

animals (Aldridge et al., 2005 ) and humans (Berkowitz et al., 2004) raising concern for possible effects in children from the use of organophosphates in the indoor environment.

Health effects (mainly sensory irritation) of **VOCs** commonly found in indoor air have been investigated in numerous studies. An extensive evaluation of all available controlled human exposure studies by a group of experts (Andersson et al., 1997) found that effect levels for irritation were usually higher than concentrations in indoor air. These studies would not explain possible health effects at much lower concentrations reported in epidemiological studies; however, the exposure was not adequately measured and cause effect relationship could not be proved due to several confounders, such as temperature, ventilation, exposure from other chemicals, or moulds and mites, as well as psychosocial factors. Anderson et al., (1997) stated that the scientific literature is inconclusive with respect to TVOC as a risk index. This conclusion is still valid, when the publications since this review are taken into account. Recent comprehensive controlled human studies at VOC concentrations considerably above those in normal homes show no effects (e.g. Fiedler et al., 2005; Laumbach et al., 2005), epidemiological studies give some indication of health effects (e.g. Hutter et al., 2006; Saijo et al., 2004; Takigawa et al., 2004) but other factors than VOC may play a major role.

Several studies have reported associations between VOCs and asthma symptoms. However, a recent comprehensive review found no consistent association between the commonly measured indoor VOC exposures and onset of new asthma cases (Nielsen et al. 2007a).

Altogether, the available evidence on VOCs in causing health effects in indoor environment is not conclusive; VOCs may also be indicators for the presence of other stressors contributing to health effects.

More recently reaction products formed in indoor air have been investigated. **Terpenes** may react with ozone to produce secondary reaction products (Wolkoff et al., 2006a). Limonene reacts with ozone and has been reported to produce both gaseous reaction products and fine and ultrafine particles (Wainman et al., 2000, Sarwar et al., 2004). The highest terpene concentrations also produced high particle levels (Sarwar et al. 2004). Several other pollutants react in indoor air and on surfaces producing known and as yet unknown reaction products (Weschler et al., 2006). In some studies, the reaction products have shown irritating properties (Clausen et al., 2001, Nøjgaard et al., 2005) and poor perceived air quality (Tamás et al., 2006) at terpene and ozone concentrations that can be present in indoor air. Adverse health effects have not been observed in all studies (Laumbach et al., 2005, Fiedler et al. 2005). Altogether, the concentrations of VOCs and ozone causing mixture effects are as yet poorly known.

In addition to the compounds emitted from the intact materials in the indoor environment there may also be new compounds formed due to **decomposition of the materials**. The glue used to fasten PVC flooring can be hydrolysed by water (dampness) from the underlying material, especially if it is concrete with a high pH. The compounds released from decomposing materials should be identified and their potential health effects evaluated.

**Phthalates** are common contaminants in the indoor environment occurring both in house dust and in indoor air and di(2-ethylhexyl) phthalate (DEHP) is the dominant component (Øie et al., 1997, Rudel et al., 2003, Fromme et al., 2004). The PVC flooring material is an important source for phthalates, but several other sources contribute in indoor environment (Bornehag et al., 2005a). PVC products indoors (different surface materials) have been associated with airway effects in epidemiological studies (Jaakkola et al. 2006) but only in one study has the concentrations of di(2-ethylhexyl) phthalate (DEHP) and butyl benzyl phthalate (BBP) been measured (Bornehag et al., 2004a). In that study DEHP was associated with asthma and BBP with rhinitis in children at the highest exposure quartile (Bornehag et al., 2004a). Long-term exposure to DEHP (Larsen et al 2007) and its

metabolite, mono-2-ethylhexyl phthalate (Hansen et al. 2007), together with a model allergen did not show promoting effects on the development of the allergen specific IgE antibodies. Phthalates are not skin sensitizers for humans and there is no evidence of respiratory sensitization (Medeiros et al., 1999, David et al., 2003, European Union Risk Assessment report, 2007). Based on the lack of mechanistic support and taking into account the low exposure level of phthalates by inhalation (Larsen et al., 2007, Nielsen et al., 2007a), the SCHER does not find consistent scientific evidence which indicate that phthalates should be high concern chemicals in indoor air. The RA report on DEHP (European Union Risk Assessment report, 2007) suggests that the MOSs from exposure in indoor air to reproductive effects, which are the basis for risk characterisation, remain large (over 200 for children, over 1000 for adults).

### **3.3.2. Concerns in relation to household-chemicals and other products (e.g. decorating materials, cleaners, furnishing, etc.)**

Household-chemicals are a large, heterogeneous group containing e.g. cleaners, furnishings, air fresheners, products for laundering, glues, paints, paint strippers, personal care products etc. The products are used mostly as liquids but some are aerosols. Candles and some air fresheners (incense) are burned. These products may emit volatile and semivolatile compounds or release inhalable aerosols and particles. Consumer products, their use and the ensuing emission concentrations in indoor areas may differ much in households across EU.

Compounds emitting from consumer products have been identified mainly in chamber studies but there are little data on their contribution to indoor air pollution. Very little is known about true exposure (in relevant use context) to components of consumer products in indoor air, in quantitative terms. Without such knowledge their health risk(s) can not be properly assessed. At least the range of resulting concentrations in indoor air in typical use situations is needed, as well as validated exposure models. The data is gradually emerging (Singer et al., 2006).

The Danish EPA has investigated the emissions of chemicals from a large number of different consumer product categories and effect on the indoor climate of these emissions has been estimated (Jensen and Knudsen, 2006). Concentrations are predicted using models and assumptions of different products being present in three different model rooms (children's room, kitchen/family room and utility room/hall). The assessment was focussed on eight VOCs (formaldehyde, acetaldehyde, phenol, benzene, toluene, xylenes, styrene and limonene) and three groups of SVOCs (phthalates, brominated flame retardants and perfluoroalkylated compounds). Of the 45 different product categories examined in the project, 33 were found to emit the selected VOCs; the exposure for the SVOCs was mainly estimated from levels in house dust. The highest concentrations were predicted in the children's room, and "typical" levels were in most cases acceptable, while worst case exposures for some of the compounds exceeded accepted limits. The worst emitters of the investigated consumer products were incense (benzene and styrene); spray paint, printed matter and electronic equipment (toluene and xylenes). It is mentioned in the report that also other sources, e.g. building materials, contribute to the total exposure but could not be taken into account, and that no assessment was done for combined exposures from several stressors.

VOCs from consumer products may contribute on average to 10-20 % of total VOCs in different indoor environments, roughly to a similar fraction as transport from outdoors (Edwards et al., 2001, Serrano-Trespalacios et al., 2004), depending on the quality of the outdoor air. Air fresheners, general purpose cleaners and floor care products have been estimated to be the major sources of VOC emissions among house-hold products e.g. in California, USA (Nazaroff and Weschler, 2004). The hazards of selected categories of cleaning agents used in Denmark have been investigated (Wolkoff et al., 1998). In some studies professional domestic cleaning has been associated with asthma (Medina-Ramón et

al., 2003) or its symptoms (Medina-Ramón et al., 2007). There are a limited number of epidemiological studies where associations of adverse health effects with consumer products have been evaluated (e.g. Farrow et al., 2003, Caress and Steinemann, 2003, Scheriff et al., 2005). In most studies the use of consumer products is one qualitative exposure category. Although some associations have been observed, the exact causal relationship remains unclear because the observed effects are associated concomitantly with a number of other factors being able to contribute as well. The heavy use of air fresheners may indicate indoor environment and/or type of living which contain several other risk factors. Therefore, due caution is needed at present in the interpretation of the results. This strengthens the need for a more integrated approach that includes determinants of exposure of different types.

Certain use conditions of consumer products (e.g. facilities in hobbies) also need more attention. Handling of products containing highly volatile components (e.g. organic solvents) in poorly ventilated spaces may result in high VOCs concentrations in air. The research needs related to household-chemicals have been included to Answer to Question 2.

### **3.3.3. Concerns in relation to building dampness/moisture and microbial growth**

Adverse health effects associated with building dampness and moisture problems have been reported since the 1980ies but are a poorly understood phenomenon. The available data about details of this subject have recently been reviewed and summarised as a panel work (IOM 2004).

An association has been shown in numerous epidemiological studies in different environments, and in a number of countries (Bornehag et al., 2001, 2004b). Intervention studies have indicated that renovation of the building either decreases or abolishes the symptoms (Sudakin 1998, Meklin et al., 2005). Furthermore, a dose-response relationship between the extent of damage and health effects has been shown (Haverinen 2002). Dampness and moisture problems in buildings are common in countries where comprehensive studies have been done, and are likely to be an underestimated indoor air problem in EU and should be evaluated more thoroughly.

The associated adverse health effects range from irritation of mucous membranes, respiratory symptoms and infections to permanent diseases, such as asthma and allergy (IOM 2004). However, only a fraction of the symptoms appear to be caused by IgE-mediated allergy, allergic alveolitis (hypersensitivity pneumonitis) or organic dust toxic syndrome, and other, still poorly known patho-physiological mechanisms are involved. General symptoms, such as fever, fatigue, headache and difficulties to concentrate have also been reported. Clusters of cases of sarcoidosis, rheumatoid diseases as well as pulmonary haemorrhage among infants have also been associated with indoor dampness (Nevalainen and Seuri, 2005).

The majority of the health effects associated with dampness and moisture of buildings are those of the respiratory system. Therefore, it is likely that the major route of the exposure to the causative agents is via the airways. There are many types of emission from a microbial growth e.g., particles of microbial origin including spores, vegetative cells and submicron-size fragments (Gorny, 2004) that carry structural components, such as endotoxin and 1,3-beta-glucan, and non-volatile secondary metabolites, e.g. toxins (Croft et al., 1986). Volatile organic compounds emitted from microbial growth include those that are known as odour of mould. Dampness and moisture may initiate chemical degradation of material which may contribute to emissions of degradation products into the indoor air and inadequate ventilation may increase the level (Bornehag et al., 2005b).

Although the association between moisture problems and adverse health effects has been demonstrated, the causative agents/exposures are not defined. This is likely to be due to

great complexity and variability of the contributing factors but also due to lack of basic knowledge. Respiratory inflammation, the most typical symptom, has been verified in laboratory animals by a few microbes typical for moisture problems (Jussila et al., 2001, Huttunen et al., 2003). Still, even in those cases, the principal components causing the effects are not known.

Dampness or moisture may accumulate into the building structures or finishing materials via leaks in roofs, windows or piping; due to condensation as a result of insufficient ventilation or faulty construction, or moisture from the ground may penetrate into the building structure by capillary movement. Attempts to improve the energy efficiency of the building e.g. by adding thermal insulation may alter the functioning of the construction, if not done properly. This may lead to moisture and mould problems. Excess water stimulates microbial growth, usually fungi and bacteria, and in a more advanced damage, also protozoa, nematodes and higher organisms such as mites and insects. The substrate (material) in question and its moisture content regulate the microbial profile that will develop, and also their toxicity (Hyvärinen et al., 2002, Roponen et al., 2001, Murtoniemi et al., 2001). Many of the bacteria and fungal species detected in damp environments are the same as detected in "normal" indoor air but their concentrations may be higher. There are also species which typically exist in water damaged environments (indicator species of the dampness problem). Microbial diversity in various dampness situations varies and one water damaged site (environment) only poorly predicts another (Nevalainen and Seuri, 2005). This may suggest that all dampness is not equally harmful. There is also likely large individual variability in sensitivity to react to those exposures, depending on e.g. the immunological status.

SCHER considers that the adverse health effects associated with building dampness and moisture are a concern. The association between building dampness and the common health effects has been documented, however several other questions, indicated as data gaps in answer to Question 2, are open and need further research before the wideness and seriousness of the problem at EU level can be assessed.

#### **4. CONCLUSIONS**

Indoor environment is a complex issue in terms of toxicology and health risk assessment. There are many different types of pollutants which may give rise to combined effects. The exposed population is the general public including vulnerable groups. Many different factors influence air quality, e.g. ventilation, cleaning conditions, properties of buildings, products used in house-holds, cultural habits, climate, outdoor air etc. Thus, large variations in indoor environments can be expected across the EU.

The SCHER considers that the health risk assessment of the pollutants in indoor environments should be done according to the principles used in the EU for risk assessment of chemicals as this is an evidence based approach. Those principles should be applied on the data available and the specific features related to indoor environment taken into account. The risk assessment paradigm should be used flexibly, taking into account that complaints and diseases related to indoor exposures may have a complex cause-effect relationship.

The SCHER considers that the data base for indoor air risk assessment is in general limited. Frequently, there are more data available for risk assessment of "classical" indoor air pollutants such, as organic pollutants as compared to particles and microbes. Especially, more data on exposure, in quantitative terms, are required. Available dose-response data seldom cover vulnerable groups. The SCHER has identified several gaps of knowledge, presented in answer to Question 2, which should be addressed by European-wide multidisciplinary research.

As to single known compounds, SCHER considers carbon monoxide, formaldehyde, benzene, nitrogen oxides and naphthalene to be compounds of concern because they have caused adverse health effects as indoor pollutants or have a high potential to cause them. Environmental tobacco smoke, radon, lead and organophosphates are also of concern. For most other pollutants the data available are yet limited for risk assessment as indoor air pollutants.

Consumer products, one source of chemicals in indoor environment, emit mostly volatile organic compounds. Lack of data on true exposure for emissions in consumer products has hampered evaluation of the associations with possible health effects most of which are also caused by other factors. The recent data suggest that some of the emitted products may react further in air and on surfaces producing secondary products, including fine and ultrafine particles. The health effects of those reaction products are poorly known.

Association of adverse health effects with dampness, water damage and mould in buildings is repeatedly shown in epidemiological studies but the causative factors and all health effects and consequences are not known. This is potentially a serious indoor air problem in EU. More research is needed to understand the associations to health effects and seriousness of the problem in EU countries.

Combined and mixture effects of indoor air pollutants can so far only rarely be assessed. There are not enough data on combined effects and also the methodology is limited. The SCHER recommends the production of data in order to make the evaluation of combined effects of indoor air pollutants feasible. In addition, the SCHER recommends taking into account routes of exposure other than inhalation (dermal, oral) in risk assessment and contribution of indoor environment exposure to total exposure from other sources. The risk assessment should be transparent to allow the evaluation of its strengths and weaknesses.

The SCHER recommends the development of health based guideline values for key pollutants and other practical guidance in general to help risk management. The SCHER also recommends that practical experiences are collected and systematized to establish evidence-based risk assessment approaches. Such examples help to solve similar type problems and decrease the overlapping risk assessment.

## **5. GENERAL CONCLUSIONS AND RECOMMENDATIONS**

Indoor air may contain over 900 chemicals, particles, and biological materials with potential health effects. Since their concentrations are usually higher than outdoors and people spend more time indoors than outdoors, the SCHER recommends that any studies to correlate outdoor air concentration with health effects need to consider the impact of indoor exposure.

The composition and concentrations of the different components in indoor air vary widely and are influenced by human activities. Since it is not feasible to regulate all possible scenarios, prevention from possible health effects and protection of sensitive populations is best achieved by reducing exposure. As a consequence the SCHER recommends that all relevant sources that are known to contribute should be evaluated. Such sources include tobacco smoke, any open fires including candles, building materials, furniture, pets and pests, use of household products, as well as conditions that lead to the growth of moulds. Constructors, maintenance personnel and inhabitants should also be aware that appropriate humidity avoids annoyances and sufficient air exchange reduces accumulation of pollutants.

---

## 6. REFERENCES

- Afsari A., Matson U., Ekberg LE. 2005. Characterization of indoor sources of fine and ultrafine particles: a study conducted in a full-scale chamber. *Indoor Air*, 15, 141-150.
- Alarie Y. 1973. Sensory irritation by airborne chemicals. *Crit. Rev. Toxicol.* 2, 299-363.
- Alarie Y, Schaper M, Nielsen GD, Abraham MH. 1998. Structure-activity relationships of volatile organic chemicals as sensory irritants. *Arch. Toxicol.*, 72, 125-140.
- Aldridge JE, Levin ED, Seidler FJ and Slotkin TA. 2005. Developmental exposure of rats to chlorpyrifos leads to behavioral alterations in adulthood, involving serotonergic mechanisms and resembling animal models of depression. *Environ. Health Perspect.* 113, 527-531.
- Andersson K, Bakke JV, Bjørseth O, Bornehag C-G, Clausen G, Hongslo JK, Kjellman M, Kjaergaard S, Levy F, Møhlhave L, Skerfving S, Sundell J. 1997. TVOC and health in non-industrial indoor environments. Report from Nordic scientific consensus meeting at Långholmen in Stockholm, 1996. *Indoor Air* 7, 78-91.
- ATSDR 2002. Guidance Manual for the assessment of joint toxic action of chemical mixture Atlanta GA, ATSDR Available at <http://www.atsdr.cdc.gov>
- Barack C, Lundahl J, Halldén G, Bylin G. 2005. Brief exposures to NO<sub>2</sub> augment the allergic inflammation in asthmatics. *Environ. Res.* 97, 58-66.
- Beasley R, Crane J, Lai CKW, Pearce N. 2000. Prevalence and etiology of asthma. *J. Allergy Clin. Immunol.* 105, S466-S472.
- Berkowitz GS, Wetmure JG, Birman-Deych E, Obel J, Lapinsky RH, Godbold JH, Holzman IR, and Wolff MS. 2004. In utero pesticide exposure, maternal paraoxonase activity, and head circumference. *Environ. Health Perspect.* 112, 388-391.
- Bornehag C-G, Blomquist G, Gyntelberg F, Järholm B, Malmberg P, Nordvall L, Nielsen A, Pershagen G, Sundell J. 2001. Dampness in buildings and health. Nordic interdisciplinary review of the scientific evidence of the associations between exposure to "dampness" in buildings and health effects (NORDDAMP). *Indoor Air* 11, 72-86.
- Bornehag C-G, Sundell J, Bonini S, Custovic A, Malmberg P, Skerfving S, Sigsgaard T, Verhoeff A. 2004b. Dampness in buildings as a risk factor for health effects, EUROEXPO: a multidisciplinary review of the literature (1998-2000) on dampness and mite exposure in buildings and health effects. *Indoor Air* 14, 243-257.
- Bornehag C-G, Sundell J, Hägerhed-Engman L, Sigsgaard T. 2005b. Association between ventilation rates in 390 Swedish homes and allergic symptoms in children. *Indoor Air* 15, 275-280.
- Bornehag CG, Sundell J, Weschler CJ, Sigsgaard T, Lundgren B, Hasselgren M, Hagerhed-Engman L. 2004a. The association between asthma and allergic symptoms in children and phthalates in house dust: a nested case-control study. *Environ. Health. Perspectives* 112, 1393-1397.
- Bornehag CG, Lundgren B, Weschler CJ, Sigsgaard T, Hagerhed-Engman L, Sundell J. 2005a. Phthalates in indoor air and their association with building characteristics. *Environ. Health. Perspectives* 113, 1399-1404.
- Butte W, Heinzow B. 2002. Pollutants in house dust as indicators of indoor contamination. *Rev. Environ. Contam. Toxicol.* 175, 1-46.
- Byrne SL, Shurdut BA, Saunders DG. 1998. Potential chlorpyrifos exposure to residents following standard crack and crevice treatment. *Environ. Health Perspect.* 106, 725-731.

California Environmental Protection Agency, 2005. Indoor air pollution in California. A draft report submitted by California Air Resources Board.

Caress SM and Steinemann AC. 2003. A review of a two-phase population study of multiple chemical sensitivities. *Environm. Health Perspect.* 111, 1490-1497.

Cassee F, Groten J, van Bladeren J, Feron V. 1998. Toxicological evaluation and risk assessment of chemical mixtures *Crit. Rev. Toxicol.* 28, 73-101.

Cavieres MF, Jaeger J, Porter W. 2002. Developmental toxicity of a commercial herbicide mixture in mice: I. Effects on embryo implantation and litter size. *Environ. Health Perspect.* 110, 1081-1085.

Chan-Yeung M, Becker A. 2006. Primary prevention of childhood asthma and allergic disorders. *Curr. Opin. Allergy Clin Immunol.* 6, 146-151.

Clausen PA, Wilkins CK, Wolkoff P, Nielsen GD. 2001. Chemical and biological evaluation of a reaction mixture of R-(+)-limonene/ozone:formation of strong airway irritants. *Environ. Int.* 26, 511-522.

Cometto-Muñiz JE, Hernández SM. 1990. Odorous and pungent attributes of mixed and unmixed odorants. *Percept Psychophys.* 47, 391-399.

Commission Directive 93/67/EEC on Risk Assessment for new notified substances.

Commission Regulation (EC) No 1488/94 on Risk Assessment for existing substances.

Council Regulation (EEC) 793/93 on the evaluation and control of the risks of existing substances

Croft WA, Jarvis BB, Yatawara C. 1986. Airborne outbreak of trichothecene toxicosis. *Atm. Env.* 20, 549-552.

Darby S, Hill D, Auvinen A, Barros-Dios JM, Baysson H, Bochicchio F et al. 2005. Radon in homes and risk of lung cancer: collaborative analysis of individual data from 13 European case-control studies. *BMJ* 330, 223. Epub 2004 Dec 21.

D'Amato G, Spiekma FTM, Liccardi G, Jäger S, Russo M, Kontou-Fili K, Nikkels H, Wüthrich B, Bonini S. 1998. Pollen-related allergy in Europe. *Allergy*, 53, 567-578.

David RM, Lockhart LK, Ruble KM. 2003. Lack of sensitization for trimellitate, phthalate, terephthalate and isobutyrate plastizers in a human repeated insult patch test. *Food Chemical. Toxicol.* 4, 589-593.

De Rosa CT, El-Masri HA, Pohl H, Cibulas W, Mumtaz MM. 2004. Implications of chemical mixtures in public health practice *J. Toxicol. Environ. Health B* 7, 339-350.

DiFranza JR, Aligne A, Weitzman M. 2004. Prenatal and postnatal environmental tobacco smoke exposure and children's health. *Pediatrics* 113, 1007-1015.

Dominici F, Peng RD, Bell ML, Pham L., McDermont A, Zeger SL, Samet JM. 2006. Fine particulate air pollution and hospital admission for cardiovascular and respiratory diseases. *JAMA* 295, 1127-1134.

Doty RL, Cometto-Muñiz JE, Jalowayski AA, Dalton P, Kendal-Reed M, Hodgson M. 2004. Assessment of upper respiratory react and ocular irritative effects of volatile chemicals in humans. *Crit. Rev. Toxicol* 34, 85-142.

Dourson M, Charnley G, Scheuplein R. 2002. Differential sensitivity of children and adults to chemical toxicity. II. Risk and regulation. *Regulat. Toxicol. Pharmacol.* 35, 448-467.

Draft RA report on DEHP. 2006 European Union Risk Assessment Report. Bis(2-ethylhexyl) phthalate.

ECA 2000. Risk Assessment in relation to indoor air quality. Report No. 22. European Collaborative Action. Urban air, indoor environment and human exposure. Environment and Quality of Life. European Commission; Joint Research Centre, Environment Institute.

ECA 2005. Harmonisation of indoor material emission labelling systems in the EU. Inventory of existing schemes. Environment and Quality of Life. Report 24. European Commission; Joint Research Centre, Institute for Health and Consumer Protection, Physical and Chemical Exposure Unit.

ECA 2006. Strategies to determine and control the contributions of indoor air pollution to total inhalation exposure (STRATEX). Environment and Quality of Life. Report 25. European Commission; Joint Research Centre, Institute for Health and Consumer Protection, Physical and Chemical Exposure Unit.

Edwards RD, Jurvelin J, Koistinen K, Saarela K, Jantunen M: 2001. VOC source identification from personal and residential indoor, outdoor and workplace microenvironment samples in EXPOLIS-Helsinki, Finland. *Athmos. Environ.* 35, 4829-4841.

Edwards RD, Schweizer C, Jantunen M, Lai HK, Bayer-Oglesby L, Katsoyanni K, Niuwenhuijsen M, Saarela K, Sram R, Künzli N. 2005. Personal exposures to VOC in the upper end of the distribution – relationships to indoor, outdoor and workplace concentrations. *Athmos. Environ.* 39, 2299-2307.

Eggleston PA. 2001. Ecology and elimination of cockroaches and allergens in the home. *J. Allergy Clin. Immunol.*, 1007, S422-429.

Elliott L, Longnecker MP, Kissling GE, London SL. 2006. Volatile organic compounds and pulmonary function in the Third National Health and Nutrition Examination Survey, 1988-1994. *Environ. Health Perspect.* 114, 1210-1214.

European Union Risk Assessment Report. 2007. Bis(2-ethylhexyl) phthalate. <http://ecb.jrc.it/existing-chemicals/>

Ezratty V, Bonay M, Neukirch C, Orset-Gulliossou G, Dehoux M, Koscielny S, Carbanes P-A, Lambrozo J, Aubier M. 2007. Effect of formaldehyde on asthmatic response to inhaled allergen challenge. *Environ. Health Perspect.* 115, 210-214.

EXPOLIS. 1999. Air Pollution Exposure in European Cities: the EXPOLIS Study. Final report. Kansanterveyslaitoksen julkaisu B 16.

EXPOLIS-INDEX. 2004. Human Exposure Patterns for Health Risk Assessment: Indoor determinants of personal exposures in the European EXPOLIS population in Athens, Basel, Grenoble, Milan, Helsinki, Oxford, and Prague. Final Report. [http://www.ktl.fi/expolis/EXPOLIS-INDEX2004/expolis\\_index.html](http://www.ktl.fi/expolis/EXPOLIS-INDEX2004/expolis_index.html)

Farrow A, Taylor H, Northstone K, Golding J. 2003. Symptoms of mothers and infants related to total volatile organic compounds in household products. *Arch. Environm. Health* 58, 633-641.

Fiedler N, Laumbach R, Kelly-McNeil K, Liroy P, Fan Z-H, Zhang J, Ottenweller J, Ohman-Strickland P, Kipen H. 2005. Health effects of a mixture of indoor volatile organics, their ozone oxidation products, and stress. *Environ. Health Perspect.* 113, 1542-1548.

Fromme H, Lahrz T, Piloty M, Gebhart H, Oddoy A, Rüden H. 2004. Occurrence of phthalates and musk fragrances in indoor air and dust from apartments and kindergartens in Berlin (Germany). *Indoor Air* 14, 188-195.

German Environmental Survey (GerES I, 1985/86; GerES II, 1990/92, GerES III, 1998). <http://www.umweltbundesamt.de/survey-e/pub/index.htm>

Gilbert NL, Guay M, Miller JD, Judek S, Chan CC, Dales RE. 2005. Levels and determinants for formaldehyde, acetaldehyde, and acrolein in residential indoor air in Prince Edward Island, Canada. *Environm. Res.* 99, 11-17.

Ginsberg G, Hattis D, Miller R, Sonawane B. 2004. Pediatric pharmacokinetic data: Implications for environmental risk assessment for children. *Pediatrics* 113, 973-983.

Gorny RL 2004. Filamentous microorganisms and their fragments in indoor air--a review. *Ann. Agric. Environ. Med.* 11, 185-97.

Grandjean P, Landrigan PL. 2006. Developmental neurotoxicity of industrial chemicals. *Lancet* 368, 2167-2178.

Gurunathan S, Robson M, Freeman N, Buckley B, Roy A, Meyer R, Bukowski J, Liroy PJ. 1998. Accumulation of chlorpyrifos on residential surfaces and toys accessible to children. *Environ. Health Perspect.* 106, 9-16.

Haddad S, Charest-Tardif G, Krishnana K. 2000. Validation of a physiological modelling framework for simulating toxicokinetics of chemicals in mixtures *Toxicol. Appl. Pharmacol.* 161, 249-257.

Hamada K, Goldsmith CA, Goldman A, Kobzik L. 2000. Resistance of very young mice to inhaled allergen sensitization is overcome by coexposure to an air pollutant aerosol. *Am. J. Respiratory Crit.Care Med.* 161, 1285-1293.

Hansen JS, Larsen ST, Poulsen LK, Nielsen GD. 2007. Adjuvant effects of inhaled mono-2-ethylhexyl phthalate in BALB/cJ mice. *Toxicology* 232, 79-88.

Hardman R. 2006. A toxicologic review of quantum dots: Toxicity depends on physicochemical and environmental factors. *Environ. Health Perspect.* 114, 165-172.

Haverinen U. 2002 Modeling moisture damage observations and their associations with health symptoms. PhD Thesis. Publications of National Public Health Institute A10/2002. Kuopio, Finland.

HCN 2000. Volatile Organic Compounds in Indoor Environment. Publication No. 2000/10 Health council of the Netherlands The Hague.

Healy JD. 2003. Excess winter mortality in Europe: a cross country analysis identifying key risk factors. *Journal of Epidemiology and Community Health* 57,784-789.

Hempel-Jørgensen A, Kjærgaard SK, Mølhav L, Hudnell KH. 1999. Sensory eye irritation in humans exposed to mixtures of volatile organic compounds. *Arch. Environ. Health* 54, 416-424.

Hodgson AT, Faulkner D, Sullivan DP, DiBartolomeo DL, Russell ML, Fisk WJ. 2003. Effect of outside air ventilation rate on volatile organic compound concentrations in a call center. *Atmospheric Environm.* 37, 5517-5527.

Hore P, Robson M, Freeman N, Zhang J, Wartenberg D, Ozkaynak H, Tulve N, Sheldon L, Needham L, Barr D, Lioy PJ. 2005. Chlorpyrifos accumulation patterns for child-accessible surfaces and objects and urinary metabolite excretion by children for 2 weeks after crack-and-crevice application. *Environ. Health Perspect.* 113, 211-219.

Hutter HP, Moshhammer H, Wallner P, Damberger B, Tappler P, Kundi M. 2006. Health complaints and annoyances after moving into a new office building: a multidisciplinary approach including analysis of questionnaires, air and house dust samples. *Int J Hyg Environ Health* 209, 65-68.

Huttunen K, Hyvärinen A, Nevalainen A, Komulainen H, Hirvonen M-R. 2003. Production of proinflammatory mediators by indoor air bacteria and fungal spores in mouse and human cell lines. *Environ. Health Perspect.* 111, 85-92

Hyvärinen A, Meklin T, Vepsäläinen A, Nevalainen A. 2002. Fungi and actinobacteria in moisture-damaged building materials – concentrations and diversity. In: *Biodeter. Biodegr.* 49, 27-37.

IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. 2004. Tobacco smoke and involuntary smoking, Vol. 83.

Ilgen E., Levsen K, Angerer K, Schneider P, Joachim H, Wichmann H-E. 2001. Aromatic hydrocarbons in the atmospheric environment – Part II: univariate and multivariate analysis and case studies of indoor concentrations. *Atmospheric Environ.*, 35, 1253-1264.

INDEX project, 2005. Critical appraisal of the setting and implementation of indoor exposure limits in the EU. EUR 21590 EN. European Commission, Directorate General, Joint Research Centre.

IOM, Institute of Medicine, National Academies of Science. 2004. Damp Indoor Spaces and Health. The National Academies Press, Washington D.C.

IPCS 1993. Principles for evaluating chemical effects on the aged population. *Environmental Health Criteria* 144, WHO.

Jaakkola JJK, Leromnimon A, Jaaakkola MS. 2006. Interior surface materials and asthma in adults: a population-based incident case-control study. *Am. J. Epidemiol.* 164, 742-749.

Jensen AA, Knudsen HN. 2006. Total health assessment of chemicals in indoor climate from various consumer products, Danish Ministry of the Environment, Survey of Chemical Substances in Consumer Products, No 75. [http://www2.mst.dk/common/Udgivramme/Frame.asp?pg=http://www2.mst.dk/Udgiv/publications/2006/87-7052-214-6/html/default\\_eng.htm](http://www2.mst.dk/common/Udgivramme/Frame.asp?pg=http://www2.mst.dk/Udgiv/publications/2006/87-7052-214-6/html/default_eng.htm)

Jussila J, Komulainen H, Huttunen K, Roponen M, Hälinen A, Hyvärinen A, Kosma V-M, Pelkonen K, Hirvonen M-R. 2001. Inflammatory responses in mice after intratracheal instillation of spores of *Streptomyces californicus* from indoor air of moldy house. *Toxicol. Appl. Pharmacol.* 171, 61-691.

Kjaergaard S, Pedersen OF, Mølhav L. 1992. Sensitivity of the eyes to airborne irritant stimuli: influence of individual characteristics. *Arch. Environ. Health* 47, 45-50.

Kephalopoulos S., Bruinen de Bruin J., Arvanitis A., Hakkinen P., Jantunen M. Consumer exposure modeling framework issues: towards harmonization at global scale. 2007. *J. Exp. Analysis and Environm. Epidemiol.*, in press.

Kosatsky T. 2005. The 2003 European heat waves. *Eurosurveillance* 10, 148-149.

Lam C-W, James JT, McCluskey R, Arepalli S, Hunter RL. 2006. A review of carbon nanotube toxicity and assessment of potential occupational and environmental health risks. *Critical Reviews in Toxicology*, 36, 189-217.

Lanphear BP, Hornung R, Khoury J, Yolton K, Baghurst P, Bellinger DC, Canfield RL, Dietrich KN, Bornschein R, Greene T, Rothenberg SJ, Needleman HL, Schnaas L, Wasserman G, Graziano J, Roberts R. 2005. Low-level environmental lead exposure and children's intellectual function: an international pooled analysis. *Environ. Health Perspect.* 113, 894-899.

Larsen ST, Hansen JS, Hansen EW, Clausen PA, Nielsen GD. 2007. Airway inflammation and adjuvant effect after repeated airborne exposures to di-(2-ethylhexyl) phthalate and ovalbumin in BALB/c mice. *Toxicology* 235, 119-129.

Laumbach RJ, Fiedler N, Gardner CR, Laskin DL, Fan Z-H, Zhang J, Weschler CJ, Liroy PL, Devlin RB, Ohman-Srickland P, Kelly-McNeil K, Kipen HM. 2005. Nasal effects of a mixture of volatile organic compounds and their ozone oxidation products. *J. Occup. Environ. Med.* 47, 1182-1189.

Lazaridis M, Aleksandropoulou V, Smolik J, Hansen JE, Glytsos T, Kalogerakis N, Dahlin E. 2006. Physico-chemical characterization of indoor/outdoor particulate matter in two residential houses in Oslo, Norway: measurements overview and physical properties – URBAN-AEROSOL Project. *Indoor Air*, 16:282-295.

Maertens RM, Bailey J, White PA. 2004. The mutagenic hazards of settled house dust: a review. *Mutat. Res.* 567, 401-425.

McCarty LS, Borgert CJ. 2006. Review of the toxicity of chemical mixtures: theory, policy and regulatory practice. *Reg. Toxicol. Pharmacol.* 45, 119-145.

Medeiros AM, Devlin DJ, Keller LH. 1999. Evaluation of skin sensitization response of dialkyl (C<sub>6</sub>-C<sub>13</sub>) phthalate esters. *Contact Dermatitis*, 41:287-289.

Medina-Ramón M, Zock JP, Kogevinas M, Sunyer J, Antó JM. 2003. Astma symptoms in women employed in domestic cleaning: a community based study. *Thorax* 58, 950-954.

Medina-Ramón M, Zock JP, Kogevinas M, Sunyer J, Torralba A., Borrel A, Burgos F, Antó JM. 2005. Astma, chronic bronchitis, and exposure to irritant agents in occupational domestic cleaning: a nested case-control study. *Occup. Environ. Med.* 62, 598-606.

Meklin T, Putus T, Pekkanen J, Hyvärinen A, Hirvonen M-R, Nevalainen A. 2005. The effects of moisture damage repairs on microbial exposure and symptoms in schoolchildren. *Indoor Air* 15 (Suppl.10), 40-47.

Murtoniemi T, Nevalainen A, Suutari M, Toivola M, Komulainen H, Hirvonen MR. 2001. Induction of cytotoxicity and production of inflammatory mediators in RAW264.7 macrophages by spores grown on six different plasterboards. *Inhal. Toxicol.* 13, 233-247.

Nazaroff WW and Weschler CJ. 2004. Cleaning products and air fresheners: exposure to primary and secondary pollutants. *Athmos. Environ.* 38:2841-2865.

- Nevalainen A, Seuri M. 2005. Of microbes and men. *Indoor Air* 15 (Suppl. 9), 58-64
- Nielsen GD. 1991. Mechanisms of activation of the sensory irritant receptor by airborne chemicals. *Crit. Rev. Toxicol.* 21, 183-208.
- Nielsen GD, Hansen JS, Lund RM, Bergqvist M, Larsen ST, Clausen SK, Thygesen P, Poulsen OP. 2002. IgE-mediated asthma and rhinitis I: A role of allergen exposure? *Pharmacol. Toxicol.*, 90, 231-242.
- Nielsen GD, Larsen ST, Olsen O, Løvik M, Poulsen LK, Glue C, Wolkoff P. 2007a. Do indoor chemicals promote development of airway allergy? *Indoor Air*, in press.
- Nielsen GD, Wolkoff P, Alarie Y. 2007b. Sensory irritation: Risk assessment approaches. *Regulatory Toxicology and Pharmacology*, in press.
- Nøjgaard JK, Christensen KB, Wolkoff P. 2005. The effect on human eye blink frequency by exposure to limonene oxidation products and methacrolein. *Toxicol. Lett.* 156 241-251.
- Pediatrics 2004. The vulnerability, sensitivity, and resiliency of the developing embryo, infant, child, and adolescent to the effects of environmental chemicals, drugs, and physical agents as compared to adults. *Pediatrics*, 114 (4), Supplement.
- Phipatanakul W, Eggleston PA, Wright EC, Wood PA and the National Cooperative Inner-City Asthma Study. 2000a . Mouse allergen. I. The prevalence of mouse allergen in inner-city homes. *J Allergy Clin. Immunol.*, 106, 1070-1074.
- Phipatanakul W, Eggleston PA, Wright EC, Wood PA and the National Cooperative Inner-City Asthma Study. Mouse allergen. 2000b. II. The relationship of mouse allergen exposure to mouse sensitization and asthma morbidity in inner-city children with asthma. *J Allergy Clin. Immunol.*, 106, 1075-1080.
- PINCHE. 2006. Policy Interpretation Network on Children's Health and Environment, QLK4-2002-02395. Report WP7, Summary PINCHE policy recommendations, [http://www.pinche.hvdgm.nl/resource/pdf/documents/final/PINCHE\\_WP7\\_final\\_110106.pdf](http://www.pinche.hvdgm.nl/resource/pdf/documents/final/PINCHE_WP7_final_110106.pdf)
- Platts-Mills TAE, Rakes G, Heyman PW. 2000. The relevance of allergen exposure to the development of asthma in children. *J. Allergy Clin. Immunol.*, 105, S503-S508.
- Reinikainen L, Jaakkola JJK. 2001. Effects of temperature and humidification in the office environment. *Arch. Environm. Health.* 56, 365-368.
- Reinikainen L, Jaakkola JJK. 2003. Significance of humidity and temperature on skin and upper airway symptoms. *Indoor Air* 13, 344-352.
- Roponen M, Toivola M, Meklin T, Ruotsalainen M, Komulainen H, Nevalainen A, Hirvonen MR. 2001. Differences in inflammatory responses and cytotoxicity in RAW264.7 macrophages induced by *Streptomyces anulatus* grown on different building materials. *Indoor Air* 11, 179-184.
- Rudel RA, Camann DE, Spengler JD, Korn LR, Brody JG. 2003. Phthalates, alkylphenols, pesticides, polybrominated diphenyl ethers, and other endocrine-disrupting compounds in indoor air and dust. *Environ. Sci. Technol.* 37, 4543-4553.
- Saarela K, Tirkkonen T, Laine-Ylijoki J, Jurvelin J, Nieuwenhuijsen MJ, Jantunen M. 2003. Exposure of population and microenvironmental distributions of volatile organic compound concentrations in the EXPOLIS study. *Atmospheric Environ.* 37, 5563-5575.

Saijo Y, Kishi R, Sata F, Katakura Y, Urashima Y, Hatakeyama A, Kobayashi S, Jin K, Kurahashi N, Kondo T, Gong Y, Umemura T. 2004. Symptoms in relation to chemicals and dampness in newly built dwellings. *Int. Arch. Occup. Environ. Health* 77, 461-470 .

Sarwar G, Olson DA, Corsi RL. 2004. Indoor particles: The role of terpene emission from consumer products. *J. Air & Waste Manage. Assoc.* 54, 367-377.

SCALE 2004a. Baseline report on research needs in the framework of the European Environment and Health Strategy ((COM 2003)338 final. [http://www.brussels-conference.org/Download/Baseline\\_report\\_TWG\\_Research\\_Needs\\_fin.pdf](http://www.brussels-conference.org/Download/Baseline_report_TWG_Research_Needs_fin.pdf)

SCALE. 2004b. Research needs in the framework of the European Environment and Health Strategy. [http://europa.eu.int/comm/environment/health/pdf/040330research\\_needs.pdf](http://europa.eu.int/comm/environment/health/pdf/040330research_needs.pdf)

Schaub B, Lauener R, von Mutius E. 2006. The many faces of the hygiene hypothesis. *J. Allergy Clin. Immunol.*, 117, 969-977.

SCHER 2006. Opinion on the report "Emission of chemicals by air fresheners. Tests on 74 consumer products sold in Europe. January 2005".  
[http://ec.europa.eu/health/ph\\_risk/committees/04\\_scher/docs/scher\\_o\\_026.pdf](http://ec.europa.eu/health/ph_risk/committees/04_scher/docs/scher_o_026.pdf)

Scheriff A, Farrow A, Golding J, the ALSPAC Study Team, Henderson J. 2005. Frequent use of chemical household products is associated with persistent wheezing in pre-school age children. *Thorax* 60,45-49.

Scheuplein R., Charnley G, Dourson M. 2002. Differential sensitivity of children and adults to chemical toxicity.I. Biological basis. *Regulat. Toxicol. Pharmacol.* 35, 448-467.

Schneider K., Gerdes H, Hassauer M, Oltmans J, Schulze J. 2002. Berücksichtigung der Riskogruppe Kind bei der Ableitung gesundheitsbezogener Umweltsstandards. (Consideration of children as a risk group of health-based environmental standards). Forschungs- und Beratungsinstitut Gefarstoffe (FoBiG) GmbH.  
<http://www.apug.de/archiv/pdf/kinderempfindlichkeit.pdf>

Seppänen OA, Fisk WJ, Mendell MJ. 1999. Association of ventilation rates and CO<sub>2</sub> concentrations with health and other responses in commercial and institutional buildings. *Indoor Air*, 9, 226-252.

Seppänen OA, Fisk WJ, Lei QH. 2005. Ventilation and performance in office work. *Indoor Air*. 16, 28-36.

Serrano-Trespalacios PI, Ryan L, Spengler JD. 2004. Ambient, indoor and personal exposure relationships of volatile organic compounds in Mexico City Metropolitan Area. *J. Exp. Anal. Env. Epidemiol.* 14, S118-S132.

Shaughnessy RJ, Haverinen-Shaughnessy U, Nevalainen A, Moschandreas D. 2006. The effects of classroom air temperature and outdoor air supply rate on the performance of school work by children. *Indoor Air* 16, 465-468.

Shoeib M, Harner T, Ikonomidou M and Kannan K. 2004. Indoor and outdoor air concentrations and phase partitioning of perfluoroalkyl sulfonamides and polybrominated diphenyl ethers. *Environ. Sci. Technol.* 38, 1313-1320.

Shusterman D. 2002. Review of upper airway, including olfaction, as mediator of symptoms. *Environ. Health Perspect.* 110 (suppl. 4), 649-653.

Shusterman D, Murphy MA, Balmes J. 2003. Differences in nasal irritant sensitivity by age, gender, and allergic rhinitis status. *Int. Arch. Occup. Environ. Health* 76, 577-583

Singer BC, Destailats H, Hodgson AT, Nazaroff WW. 2006. Cleaning products and air fresheners and resulting concentrations of glycol ethers and terpenoids. *Indoor Air* 16, 179-191.

Stapleton HM, Dodder NG, Offenberg JH, Schantz MM and Wise SA. 2005. Polybrominated diphenyl ethers in house dust and clothes dryer lint, *Environ. Sci. Technol.* 39, 925-931.

Sudakin DL. 1998. Toxigenic fungi in water damaged building: An intervention study. *Am J Ind Med* 34,183-190.

Takigawa T, Horike T, Ohashi Y, Kataoka H, Wang DH, Kira S. 2004. Were volatile organic compounds the inducing factors for subjective symptoms of employees working in newly constructed hospitals? *Environ. Toxicol.* 19, 280-290.

Tamás G, Weschler CJ, Toftum J, Fanger PO. 2006. Influence of ozone-limonene reactions on perceived air quality. *Indoor Air* 16, 168-178.

Tamburlini G, von Ehrestein OS., Bertollini R. 2002. Children's health and environment: A review of evidence. Environmental issue report 29. WHO and EEA. <http://www.who.dk/document/e75518.pdf>

Thade Report. 2004. Towards healthy air in dwellings in Europe. <http://www.efanet.org/activities/documents/THADEReport.pdf>

TGD 2003. Technical Guidance Document on Risk Assessment. 2003. Part I. European Chemicals Bureau.

TNO 2005. Risks to health and the environment related to the use of lead in products. TNO report. STB-01-39 /Final). <http://ec.europa.eu/enterprise/chemicals/docs/studies/tno-lead.pdf>

TNO and RIVM. 2006. Tender ENV.D.4/ETU/2005/0074r. "Study on the treatment of vulnerable groups in EU risk assessment" FINAL REPORT. TNO Quality of Life, Food and Chemical Risk Analysis Department, Zeist, The Netherlands and National Institute of Public Health and Environment (RIVM), Bilthoven, The Netherlands.

US EPA Guidelines for the health risk assessment of chemical mixtures 1986. EPA 630/R-98/002. Washington, DC: US Environmental Protection Agency.

US EPA Supplementary guidance for conducting health risk assessment of chemical mixtures 2001. EPA 630/R-00/002. Washington, DC: US Environmental Protection Agency, Risk Assessment Forum.

Wainman T, Zhang J, Weschler CJ, Liou P. 2000. Ozone and limonene in indoor air: A source of submicron particle exposure. *Environ. Health Perspect.* 108 1139-1145.

Van den Berg M, Birnbaum L, Bosveld B, Brunstrom B, Cook P, Feeley L et al. 1998. Toxic equivalency factors (TEFs) for PCBs, PCDDs, and PCDFs for humans and wildlife. *Environ. Health Perspect.* 106, 775-792.

Weis N, Stolz P, Krooß J, Meierhenrich U. 1998. Dichlorvos-Insektenstrips in Innenräumen: Belastung und Risikoabschätzung. *Gesundheitswesen* 60, 445-449.

Wargocki P, Wyon DP. 2007. The effects of outdoor air supply rate and supply air filter condition in classrooms on the performance of schoolwork by children. HVAC&R Research 13, 165-191.

Weschler CJ, Wells JR, Poppendieck D, Hubbard H, Pearce TA. 2006. Workgroup report: Indoor air chemistry. Environ. Health Perspect. 114, 442-446.

WHO 1997. Assessment of exposure to indoor air pollutants. Jantunen M, Jaakkola JJK and Krzyzanowski M (editors). WHO Regional Publications, European Series, No. 78.

WHO 1998. International Agency for Research of Cancer. Man-made mineral fibres and radon. IARC Monographs on the evaluation of carcinogenic risks to humans. Vol. 43.

WHO 2000. Air quality guidelines. 2<sup>nd</sup> edition, Regional Office for Europe, 2000. [http://www.euro.who.int/air/activities/20050223\\_4](http://www.euro.who.int/air/activities/20050223_4)

WHO 2003. Health aspects of air pollution with particulate matter, ozone, and nitrogen dioxide. Report on a WHO working group. Copenhagen, WHO Regional Office for Europe (document EUR/o4/5042688). <http://www.euro.who.int/document/e79097.pdf>

WHO 2004a. Health aspects of air pollution. Results from the WHO project "Systematic review of health aspect of air pollution in Europe" WHO Regional Office for Europe, Copenhagen. <http://www.euro.who.int/document/E83080.pdf>

WHO. 2004b. WHO specifications and evaluations for public health pesticides. Chlorpyrifos. O,O-diethyl-O-3,5,6-trichloro-2-pyridyl phosphorothioate. World Health Organization, Geneva. [www.who.int/entity/whopes/quality/en](http://www.who.int/entity/whopes/quality/en)

WHO 2005a. Principles of characterising and applying human exposure models. Harmonization Project Document No. 3. , WHO, Geneva.

WHO 2005b. Effects of air pollution on children's health and development. Report on a WHO working group. European Centre for Environment and Health . Bonn office. <http://www.euro.who.int/document/E86575.pdf>

Wilford BH, Shoeib M, Harner T, Zhu J and Jones KC. 2005. Polybrominated diphenyl ethers in indoor dust in Ottawa, Canada: Implications for sources and exposure, Environ. Sci. Technol., 39, 7027-7035.

Wolkoff, P, Nielsen, GD. 2001. Organic compounds in indoor air – their relevance for perceived indoor air quality? Atmos. Environ. 35, 4407-4417.

Wolkoff P, Nøjgaard JK, Franck C, Skov P. The modern office desiccates the eyes? 2006b. Indoor Air 16, 258-265.

Wolkoff P, Schneider T, Kildeso J, Degerth R, Jarozewski M, Schunk H. 1998. Risk in cleaning: chemical and physical exposure. Sci. Tot. Environment 215, 135-156.

Wolkoff P, Wilkins CK, Clausen PA, Nielsen GD. 2006a. Organic compounds in office environments – sensory irritation, odor, measurements and the role of reactive chemistry. Indoor Air 16, 7-19.

Øie L., Hersoug L-G, Madsen JØ. 1997. Residential exposure to plasticizers and its possible role in the pathogenesis of asthma. Environm. Health Perspect., 105, 972-978.

## 7. LIST OF ABBREVIATIONS

ATSDR	Agency for Toxic Substances and Disease Registry
DEHP	Di(2-ethylhexyl) phthalate
ETS	Environmental tobacco smoke
HI	Hazard Index
IPCS	International Programme on Chemical Safety
LOEAL	Lowest-Observed-Adverse-Effect-Level
MEHP	Mono (2-ethylhexyl) phthalate
MOS	Margin of Safety
NIOSH	National Institute of Occupational Safety, Health
NOAEL	No-Observed-Adverse-Effect-Level
NOEL	No-Observed-Effect-Level
ODTS	Organic dust toxic syndrome
OEHHA	Office of Environmental Hazard Assessment of Californian Environmental Protection Agency
PAH	Polyaromatic hydrocarbon
PBPD	Physiology-based pharmacodynamic model
PBPK	Physiology-based pharmacokinetic model
PCB	Polychlorinated Biphenyl
PVC	Polyvinyl Chloride
RfD	Reference Dose
RPF	Relative Potency Factor
SVOC	Semi volatile Organic Compound
TGD	Technical Guidance Document
TEF	Toxic Equivalent Factor
TVOC	Total Volatile Organic Compounds
UBA	Federal Environmental Agency of Germany (Umweltbundesamt)
VOC	Volatile Organic Compound